Disks large homolog 3 (DLG3) is differentially expressed in the tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab (Herceptin), a monoclonal antibody targeting the extracellular domain of the human epidermal growth factor receptor 2 (HER2)¹, is utilized for the treatment of adjuvant and metastatic breast cancer² in the United States and worldwide. We mined published microarray and gene expression data^{3,4} to discover in an unbiased fashion the most significant transcriptional changes associated with treatment with trastuzumab in patients with breast cancer. We identified disks large homolog 3 (DLG3) as among the genes most differentially expressed in the primary tumors of patients with breast cancer treated with trastuzumab. The primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of DLG3 than did patients not treated with trastuzumab, and a single administration of trastuzumab was sufficient to drive differential expression of DLG3 in primary tumors of the breast, demonstrating that a gene whose expression is associated with decreased survival in patients with breast cancer⁵ is transcriptionally induced in primary tumors of the breast as a result of treatment with trastuzumab.

Keywords: trastuzumab, breast cancer, DLG3, disks large homolog 3, systems biology of breast cancer, targeted therapeutics in breast cancer.

1 The mechanism of action ascribed to trastuzumab function is thought to be binding of the F_{ab}
2 domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2
3 (HER2)¹, but the transcriptional consequences resulting from signaling downstream of trastuzumab
4 binding to HER2 is less well understood. By performing unbiased, systems-level comparative
5 transcriptome analysis of primary tumors of the breast in patients treated with trastuzumab and those not
6 treated with trastuzumab using published microarray and gene expression data^{3,4}, we report here that the
7 tyrosine kinase DLG3 is a differentially expressed gene in the tumors of trastuzumab-treated patients: the
8 primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of DLG3 than
9 did those of patients not treated with trastuzumab.

10 **Methods**

11 We used datasets GSE38057³ and GSE114082⁴ this differential gene expression analysis of the
12 transcriptional response to trastuzumab treatment in patients with breast cancer. GSE38057 was
13 generated using DASL Human Cancer Panel by Probe technology with $n=6$ primary breast tumors from
14 patients treated with induction chemotherapy and/or hormone therapy without trastuzumab and $n=35$
15 primary breast tumors from patients treated with induction chemotherapy and/or hormone therapy in
16 addition to trastuzumab, both of which developed brain metastases; analysis was performed using
17 platform GPL5826. GSE114082 was generated using Illumina HumanHT-12 WG-DASL V4.0 R2
18 expression beadchip technology with $n=17$ primary breast tumors from patients before trastuzumab
19 treatment and $n=17$ primary tumors from patients after trastuzumab treatment, analysis performed using
20 platform GPL14951. The Benjamini and Hochberg method of p -value adjustment was used for ranking
21 of differential expression but raw p -values were used to assess statistical significance of global or
22 multiplexed differential expression. Log-transformation of data was auto-detected, and the NCBI
23 generated category of platform annotation was used. A statistical test was performed to evaluate whether
24 DLG3 gene expression was significantly between primary breast tumors of patients receiving trastuzumab
25 using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of
26 differential gene expression in human breast cancer (Version 8.4.0)(455).

27 **Results**

28 We harnessed the power of blind, comparative transcriptome analysis using published microarray
and gene expression data^{3,4} to identify the most significant transcriptional features of the primary breast
tumors of patients treated with trastuzumab (Herceptin).

DLG3 is differentially expressed in the tumors of patients treated with trastuzumab.

We identified the transcription factor DLG3 as among the genes whose expression changed most
significantly in the tumors of breast cancer patients treated with trastuzumab in a multiplexed gene
expression dataset³ (Table 1). When sorting each of the genes expressed in the primary tumors of breast
cancer patients treated with trastuzumab based on significance of change as compared to that of patients
not treated with trastuzumab, DLG3 ranked 47 out of 1506 total transcripts (Table 1), equating to 96.9%
differential expression. DLG3 differential expression in the tumors of patients treated with trastuzumab
was statistically significant (Table 1; $p=0.04232$).

The first dataset, although measuring the expression of over 1500 genes, was still limited. We
turned to a second microarray dataset⁴, this measuring the expression of genes across the transcriptome -
29377 genes in total. The tumors compared in this dataset were sampled from the same patient before
and after a single exposure to trastuzumab, not comparing tumors from different patients treated with a
course of trastuzumab or not. Thus, while differentially expressed genes when comparing tumors based
on trastuzumab treatment can result from a variety of sources, including gene expression changes

1 resulting from tumor evolution over the course of treatment, changes identified in this dataset, occurring
2 after only a single treatment of trastuzumab, are more likely to be a direct transcriptional result of
3 trastuzumab binding events. We found again that DLG3 was among the genes most differentially
4 expressed in the tumors of patients administered a single treatment of trastuzumab (Table 2). When
5 sorting each of the genes expressed in the tumors of patients administered a single treatment of
6 trastuzumab, DLG3 ranked 2127 out of 29377 total transcripts (Table 2), equating to 92.8% differential
7 expression. Differential expression of DLG3 in the primary tumors of breast cancer patients following
8 treatment with trastuzumab was statistically significant (Table 2; $p=5.72E-04$).

9 **DLG3 is expressed at higher levels in the tumors of breast cancer patients treated with**
10 **trastuzumab than in the tumors of those not treated with trastuzumab.**

11 We obtained exact mRNA expression levels for DLG3 from tumors of patients treated with
12 trastuzumab and from the tumors of patients not treated with trastuzumab to directly evaluate magnitude
13 and change of expression in DLG3 based on history of trastuzumab treatment. DLG3 was expressed at
14 higher levels in the tumors of patients treated with trastuzumab as compared to those of patients not
15 treated with trastuzumab, but this difference was not deemed statistically significant (Figure 1; $p=0.1695$).
16 We calculated a mean fold change of 1.39 ± 0.55 in DLG3 expression in the primary tumors of patients
17 treated with trastuzumab relative to that of patients not treated with trastuzumab.

18 Thus, we found using published multiplexed gene expression data³ and a microarray dataset⁴ that
19 DLG3 was among the genes most differentially expressed in the tumors of patients treated with
20 trastuzumab, that DLG3 was expressed at higher levels in the tumors of patients treated with trastuzumab
21 as compared to those not treated with trastuzumab, and that this transcriptional change was likely a direct
22 result of trastuzumab treatment as opposed to a gene expression change associated with clonal selection
23 and evolution of the tumor across the course of treatment.

24 **Discussion**

25 We provided evidence here that the disks large homolog 3, also known as SAP102 or neuronal
26 and endocrine DLG3 is a differentially expressed gene in the tumors of patients treated with trastuzumab,
27 and that the primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of
28 DLG3 in their primary tumors than did patients not treated with trastuzumab, likely a direct result of
trastuzumab administration. Thus, trastuzumab treatment results in expression of a gene whose
expression in patients with breast cancer is associated with inferior survival⁵, a molecule that has multiple
functions in the nervous system, including clearance of NMDA receptors⁶, as well as regulation of
dendritic spine morphology⁷ and cortical synapse development⁸.

References

1. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
2. Herceptin (Trastuzumab). Indications. <https://www.herceptin.com/hcp/indications.html>.
3. Duchnowska, R., Jassem, J., Goswami, C.P., Dundar, M., Gökmen-Polar, Y., Li, L., Woditschka, S., Biernat, W., Sosińska-Mielcarek, K., Czartoryska-Arłukowicz, B. and Radecka, B., 2015. Predicting early brain metastases based on clinicopathological factors and gene expression analysis in advanced HER2-positive breast cancer patients. *Journal of Neuro-oncology*, 122(1), pp.205-216.
4. Triulzi, T., Regondi, V., De Cecco, L., Cappelletti, M.R., Di Modica, M., Paolini, B., Lollini, P.L., Di Cosimo, S., Sfondrini, L., Generali, D. and Tagliabue, E., 2018. Early immune modulation by single-agent trastuzumab as a marker of trastuzumab benefit. *British journal of cancer*, 119(12), pp. 1487-1494.
5. Liu, J., Li, P., Wang, R., Li, J., Zhang, M., Song, Z. and Liu, P., 2019. High expression of DLG 3 is associated with decreased survival from breast cancer. *Clinical and Experimental Pharmacology and Physiology*, 46(10), pp.937-943.
6. Chen, B.S., Gray, J.A., Sanz-Clemente, A., Wei, Z., Thomas, E.V., Nicoll, R.A. and Roche, K.W., 2012. SAP102 mediates synaptic clearance of NMDA receptors. *Cell reports*, 2(5), pp.1120-1128.
7. Müller, B.M., Kistner, U., Kindler, S., Chung, W.J., Kuhlendahl, S., Fenster, S.D., Lau, L.F., Veh, R.W., Huganir, R.L., Gundelfinger, E.D. and Garner, C.C., 1996. SAP102, a novel postsynaptic protein that interacts with NMDA receptor complexes in vivo. *Neuron*, 17(2), pp.255-265.
8. Murata, Y. and Constantine-Paton, M., 2013. Postsynaptic density scaffold SAP102 regulates cortical synapse development through EphB and PAK signaling pathway. *Journal of neuroscience*, 33(11), pp. 5040-5052.

Rank	ID	p-value	t	B	FC	Gene
47	GI_10863920-S-2	0.04232	2.092746	-4.33	1.39 ± 0.55	DLG3

Table 1: DLG3 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in DLG3 between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Rank	ID	p-value	t	B	Gene
2127	ILMN_1811515	5.72E-04	-3.7801874	-0.35997	DLG3

Table 2: A single exposure to trastuzumab is sufficient to drive differential expression of DLG3 in the primary tumors of patients with breast cancer.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: DLG3 is expressed at higher levels in the primary tumors of patients treated with trastuzumab.

The mRNA expression level of DLG3 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between DLG3 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.