hnRNP U is differentially expressed in whole blood from patients with sepsis.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY, USA

We probed published and public microarray datasets^{1,2} to discover the most significant gene expression changes in the blood of patients with sepsis. We identified significant differential expression of the heterogenous nuclear ribonucleoprotein hnRNP U in whole blood from patients with sepsis.

Keywords: sepsis, septic shock, hnRNP U, systems biology of septic shock.

1 Septic shock is a leading cause of mortality in the United States and worldwide³. We
2 used published and public microarray datasets^{1,2} to identify differentially expressed genes in the
3 blood of patients with sepsis. We found significant changes in expression of the heterogenous
4 nuclear ribonucleoprotein hnRNP U in the whole blood of patients with sepsis.
5

6 **Methods**

7
8
9 We utilized microarray datasets GSE100159¹ and GSE26440² for this differential gene
10 expression analysis of blood cells during sepsis. GSE100509 was generated with whole blood
11 using Illumina HumanWG-6 v3.0 expression beadchip technology with $n=12$ whole blood from
12 control subjects and $n=33$ whole blood from sepsis patients. GSE26440 was generated using
13 Affymetrix Human Genome U133 Plus 2.0 Array technology with $n=32$ control subjects and
14 $n=98$ sepsis patients. The Benjamini and Hochberg method of p-value adjustment was used for
15 ranking of differential expression but raw p-values were used for assessment of statistical
16 significance of global differential expression. Log-transformation of data was auto-detected, and
17 the NCBI generated category of platform annotation was used. A statistical test was performed
18 to evaluate whether hnRNP U expression was significantly different when comparing whole
19 blood from control subjects and from patients with sepsis using a two-tailed, unpaired t-test with
20 Welch's correction. GEO2R provides information on differential expression for every transcript
21 measured by microarray but exact mRNA expression values only for the 250 most differentially
22 expressed genes. We used PRISM for all statistical analyses (Version 8.4.0)(455).
23
24
25
26
27
28

Results

We mined published and public microarray datasets to identify differentially expressed genes in patients with sepsis in the whole blood.

hnRNP U is differentially expressed in whole blood from patients with sepsis

We identified the heterogenous ribonucleoprotein hnRNP U as among the most differentially expressed genes in the sepsis transcriptome. When sorting each transcript measured by significance of change in expression between whole blood from control subjects and whole blood from patients with sepsis, hnRNP U ranked 83 out of 48803 total transcripts, or differential expression greater than 99.8% of the transcripts measured globally (Table 1). Differential expression of hnRNP U in whole blood from patients with sepsis was statistically significant (Table 1; $p=6.37E-12$).

hnRNP U is differentially expressed in whole blood from patients with sepsis

We probed additional datasets to determine if we could observe differential expression of hnRNP U gene in any independent microarray datasets. Indeed, we observed significant differential expression of hnRNP U in whole blood from patients with sepsis in a second microarray dataset (Table 2). When sorting each transcript measured by significance of change in expression between whole blood from control subjects and whole blood from patients with sepsis, hnRNP U ranked 2315 out of 54675 total transcripts, or differential expression greater than 95.8% of the transcripts measured globally (Table 2). Differential expression of hnRNP U

1 in whole blood from patients with sepsis was statistically significant (Table 2; $p=1.37E-10$).

2 3 hnRNP U is expressed at significantly lower levels in whole blood from patients with sepsis

4 We obtained exact mRNA expression values for the differentially expressed hnRNP U
5 transcript in whole blood from patients with sepsis and from control subjects to directly compare
6 hnRNP U expression. We also performed a statistical test to determine if this difference was
7 significantly different. hnRNP U was expressed at lower levels in whole blood from patients
8 with sepsis and this difference was statistically significant (Figure 1; $p<0.0001$). We calculated a
9 2.99 ± 0.99 fold change in hnRNP U gene expression when comparing whole blood from
10 patients with sepsis to whole blood from healthy controls (Table 1).
11
12
13

14 Thus, we found that in the blood of patients with sepsis, hnRNP U was among the genes
15 whose expression changed most significantly; expression of hnRNP U was markedly repressed
16 in the whole blood during sepsis.
17
18

19 **Discussion**

20
21 Heterogenous nuclear ribonucleoproteins, hnRNPs are a family of at least twenty nucleic
22 acid-binding proteins that make up approximately 80% of the mass of an hnRNP particle and
23 function in processing of immature RNA transcripts (pre-mRNA) present in heterogenous
24 nuclear RNA (hnRNA) into processed messenger RNA (mRNA) during splicing of pre-mRNAs⁴.
25 hnRNP U, also known as scaffold attachment factor A (SAF-A), is among the largest of the
26 hnRNPs^{5,6}. hnRNP U can interact with pre-mRNA and DNA *in vivo*^{5,7} and bind both
27
28

1 single- and double-stranded DNA and RNA *in vitro*^{5,6}. Unlike other hnRNPs, hnRNP U does not
2 have a consensus RNA binding-domain (CS-RBD) / RNP motif. hnRNP U possesses an RNA-
3 binding RGG box rich in arginine-glycine-glycine repeats⁵. hnRNP U can bind a large
4 population of non-coding RNAs including small nuclear RNAs (snRNAs) important for intron
5 splicing is bound by hnRNP U⁸. An important hnRNP U interaction partner is actin; hnRNP U
6 can be found together with actin at the phosphorylated C-terminal domain (CTD) of RNA
7 Polymerase II (Pol II); nuclear microinjection of a peptide representing the actin-binding
8 sequence of hnRNP U can block BrUTP incorporation *in vivo*⁹. However, a separate study
9 suggested hnRNP U might function as an inhibitor of transcriptional elongation, joining the pre-
10 initiation complex through interactions with the Pol II holoenzyme; hnRNP U could bind TFIIH
11 and impair phosphorylation of the C-terminal domain (CTD) of Pol II by TFIIH *in vitro*¹⁰.
12 Support for a pro-transcriptional function of hnRNP U interactions at the Pol II CTD is
13 disruption of the interaction between actin and hnRNP U by the long non-coding RNA H19
14 resulting in impairment of transcription¹¹. hnRNP U can also interact with DNA topoisomerase
15 II β , acting to enhance relaxation of DNA supercoiling by DNA topoisomerase II β ¹².

22 Long non-coding RNAs are key interaction partners of hnRNP U. hnRNP U can interact
23 with the long non-coding RNA Xist through its RNA-binding domain, where hnRNP U functions
24 in the recruitment of Xist to the inactive X chromosome (Xi) (“painting”)^{13,14}. Depletion of
25 hnRNP U results in detachment of Xist from the Xi, and embryonic stem cells deficient in
26
27
28

1 hnRNP U do not generate Xi¹³. The long non-coding RNA FIRRE is another interaction partner
2 of hnRNP U¹⁵. A positive regulator of inflammatory gene expression in macrophages, FIRRE
3 can, in response to lipopolysaccharide stimulation, regulate the stability of mRNAs involved in
4 the inflammatory response like VCAM1 and IL12p40 through interactions with RNA messages
5 at AU-rich elements in conjunction with hnRNP U¹⁵.
6
7

8 Transcription factors are other important hnRNP U interaction partners. hnRNP U can
9 bind the Yes-associated protein YAP; interaction of YAP and hnRNP U occurs through a proline-
10 rich hnRNP U binding site at the amino-terminus of YAP. Together, YAP and hnRNP U could
11 promote transcription of a p73-driven Bax reporter construct¹⁶. One study demonstrated
12 interaction of hnRNP U with the c-Myc-Max heterodimer at E-boxes at the promoter of the
13 ornithine decarboxylase (ODC) gene, and depletion of hnRNP U impaired transcriptional
14 activation of ODC, suggesting co-activator functions for hnRNP U with the c-Myc-Max
15 complex¹⁷. Another nuclear interaction partner of hnRNP U, of particular relevance here, is the
16 glucocorticoid receptor (GR)^{18,19}; nuclear translocation of hnRNP U is dependent on interactions
17 with GR¹⁸. In response to DNA double-strand breaks, serine 59 of hnRNP U is phosphorylated
18 by DNA protein kinase (DNA-PK)²⁰. In HIV-1 infection, the amino-terminus of hnRNP U was
19 identified in a genetic screen designed to discover genes important for induction of resistance
20 against recombinant HIV-1 clones; the recovered fragment of hnRNP U can bind the 3' long
21 terminal repeat of viral RNAs and effectively suppress HIV-1 gene expression²¹.
22
23
24
25
26
27
28

1 hnRNP U has important cytosolic functions. hnRNP U can interact with caspase-9 pre-
2 mRNA (C9/E3) and compete with hnRNP L for binding to C9/E3, and depletion of hnRNP U
3 resulted in alteration of the caspase-9a/caspase-9b mRNA ratio, suggesting a caspase-9 splice-
4 enhancing function for hnRNP U²². Phosphorylation of hnRNP L by protein kinase Akt could
5 disrupt binding of hnRNP U to C9/E3, together demonstrating regulation of an hnRNP U
6 caspase-9 splicing function by hnRNP L phosphorylation²².

9 Mice with conditional deletion of hnRNP U in the muscle driven by skeletal actin Cre
10 recombinase are born in normal Mendelian ratios but develop adult-onset myopathy featuring
11 Akt activation, glycolytic muscle atrophy and changes in splicing of transcripts encoded by
12 genes involved in metabolic function including *Senp1*, *Mff*, *Sorbs1* and *Inpp4a*. Strand-specific
13 paired-end RNA-sequencing analysis of alternative splicing in *Hnrnpu* mutants revealed that
14 skipped exons and mutually exclusive exons were more frequent than usage of alternative 5' and
15 3' splice sites, suggesting hnRNP U might be important for definition of intron/exon boundaries.
16 The regenerative response of young mice deficient in hnRNP U in the muscle resembles that of
17 older wild-type mice²³. Thus, hnRNP U can interact with a host of non-coding RNAs,
18 transcription factors and transcriptional regulators to modulate target gene expression.

23 At this stage, it is difficult to draw conclusions on which of these roles, if any, might be
24 perturbed by changes in hnRNP U mRNA levels during sepsis. We found that changes in
25 expression of the heterogenous nuclear ribonucleoprotein hnRNP U gene were among the most
26 significant transcriptome-wide in whole blood from patients with sepsis.

References

1. GSE100159. Nicole Baldwin. BIIR. Dallas, TX, USA. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE100159>.
2. Wong, H.R., Cvijanovich, N., Lin, R., Allen, G.L., Thomas, N.J., Willson, D.F., Freishtat, R.J., Anas, N., Meyer, K., Checchia, P.A. and Monaco, M., 2009. Identification of pediatric septic shock subclasses based on genome-wide expression profiling. *BMC medicine*, 7(1), p. 34.
3. Stevenson, E.K., Rubenstein, A.R., Radin, G.T., Wiener, R.S. and Walkey, A.J., 2014. Two decades of mortality trends among patients with severe sepsis: a comparative meta-analysis. *Critical care medicine*, 42(3), p.625.
4. Dreyfuss, G., Swanson, M.S. and PifioI-Roma, S., 1988. Heterogeneous nuclear ribonucleoprotein particles and the pathway of mRPIA formation.
5. Kiledjian, M. and Dreyfuss, G., 1992. Primary structure and binding activity of the hnRNP U protein: binding RNA through RGG box. *The EMBO journal*, 11(7), pp.2655-2664.
6. Fackelmayer, F.O. and Richter, A., 1994. Purification of two isoforms of hnRNP-U and characterization of their nucleic acid binding activity. *Biochemistry*, 33(34), pp. 10416-10422.
7. Göhring, F. and Fackelmayer, F.O., 1997. The scaffold/matrix attachment region binding protein hnRNP-U (SAF-A) is directly bound to chromosomal DNA in vivo: a chemical cross-linking study. *Biochemistry*, 36(27), pp.8276-8283.
8. Xiao, R., Tang, P., Yang, B., Huang, J., Zhou, Y., Shao, C., Li, H., Sun, H., Zhang, Y. and Fu, X.D., 2012. Nuclear matrix factor hnRNP U/SAF-A exerts a global control of alternative splicing by regulating U2 snRNP maturation. *Molecular cell*, 45(5), pp.656-668.
9. Kukalev, A., Nord, Y., Palmberg, C., Bergman, T. and Percipalle, P., 2005. Actin and hnRNP U cooperate for productive transcription by RNA polymerase II. *Nature structural & molecular biology*, 12(3), pp.238-244.
10. Kim, M.K. and Nikodem, V.M., 1999. hnRNP U inhibits carboxy-terminal domain phosphorylation by TFIIF and represses RNA polymerase II elongation. *Molecular and cellular biology*, 19(10), pp.6833-6844.

- 1 11. Bi, H.S., Yang, X.Y., Yuan, J.H., Yang, F., Xu, D., Guo, Y.J., Zhang, L., Zhou, C.C., Wang, F.
2 and Sun, S.H., 2013. H19 inhibits RNA polymerase II-mediated transcription by disrupting
3 the hnRNP U–actin complex. *Biochimica et Biophysica Acta (BBA)-General Subjects*,
4 1830(10), pp.4899-4906.
- 5 12. Kawano, S., Miyaji, M., Ichiyasu, S., Tsutsui, K.M. and Tsutsui, K., 2010. Regulation of
6 DNA topoisomerase II β through RNA-dependent association with heterogeneous nuclear
7 ribonucleoprotein U (hnRNP U). *Journal of Biological Chemistry*, 285(34), pp.
8 26451-26460.
- 9 13. Hasegawa, Y., Brockdorff, N., Kawano, S., Tsutui, K., Tsutui, K. and Nakagawa, S., 2010.
10 The matrix protein hnRNP U is required for chromosomal localization of Xist RNA.
11 *Developmental cell*, 19(3), pp.469-476.
- 12 14. Sakaguchi, T., Hasegawa, Y., Brockdorff, N., Tsutsui, K., Tsutsui, K.M., Sado, T. and
13 Nakagawa, S., 2016. Control of chromosomal localization of Xist by hnRNP U family
14 molecules. *Developmental cell*, 39(1), pp.11-12.
- 15 15. Lu, Y., Liu, X., Xie, M., Liu, M., Ye, M., Li, M., Chen, X.M., Li, X. and Zhou, R., 2017. The
16 NF- κ B–responsive long noncoding RNA FIRRE regulates posttranscriptional regulation of
17 inflammatory gene expression through interacting with hnRNPU. *The Journal of*
18 *Immunology*, 199(10), pp.3571-3582.
- 19 16. Howell, M., Borchers, C. and Milgram, S.L., 2004. Heterogeneous nuclear ribonuclear
20 protein U associates with YAP and regulates its co-activation of Bax transcription. *Journal*
21 *of Biological Chemistry*, 279(25), pp.26300-26306.
- 22 17. Matsuoka, Y., Uehara, N. and Tsubura, A., 2009. hnRNP U interacts with the c-Myc-Max
23 complex on the E-box promoter region inducing the ornithine decarboxylase gene. *Oncology*
24 *reports*, 22(2), pp.249-255.
- 25 18. Eggert, M., Michel, J., Schneider, S., Bornfleth, H., Baniahmad, A., Fackelmayer, F.O.,
26 Schmidt, S. and Renkawitz, R., 1997. The glucocorticoid receptor is associated with the
27 RNA-binding nuclear matrix protein hnRNP U. *Journal of Biological Chemistry*, 272(45),
28 pp.28471-28478.
19. Eggert, H., Schulz, M., Fackelmayer, F.O., Renkawitz, R. and Eggert, M., 2001. Effects of
the heterogeneous nuclear ribonucleoprotein U (hnRNP U/SAF-A) on glucocorticoid-
dependent transcription in vivo. *The Journal of steroid biochemistry and molecular biology*,
78(1), pp.59-65.

20. Berglund, F.M. and Clarke, P.R., 2009. hnRNP-U is a specific DNA-dependent protein kinase substrate phosphorylated in response to DNA double-strand breaks. *Biochemical and biophysical research communications*, 381(1), pp.59-64.
21. Valente, S.T. and Goff, S.P., 2006. Inhibition of HIV-1 gene expression by a fragment of hnRNP U. *Molecular cell*, 23(4), pp.597-605.
22. Vu, N.T., Park, M.A., Shultz, J.C., Goehe, R.W., Hoeferlin, L.A., Shultz, M.D., Smith, S.A., Lynch, K.W. and Chalfant, C.E., 2013. hnRNP U enhances caspase-9 splicing and is modulated by AKT-dependent phosphorylation of hnRNP L. *Journal of Biological Chemistry*, 288(12), pp.8575-8584.
23. Bagchi, D., Mason, B.D., Baldino, K., Li, B., Lee, E.J., Zhang, Y., Chu, L.K., El Raheb, S., Sinha, I. and Neppl, R.L., 2020. Adult onset myopathy with constitutive activation of Akt following the loss of hnRNP-U. *iScience*, p.101319.

Rank	ID	p-value	t	B	FC	Gene	Gene name
83	ILMN_1743677	6.37E-12	-9.1280127	17.02245	2.99 ± 0.99	HNRNPU	heterogeneous nuclear ribonucleoprotein U

Table 1: hnRNP U is differentially expressed in whole blood from patients with sepsis.

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the fold change of hnRNP U in whole blood from patients with sepsis compared to whole blood from control subjects, the fold change in hnRNP U expression in whole blood from patients with sepsis, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
2315	235603_at	1.37E-10	-6.9734572	13.669543	HNRNPU	heterogeneous nuclear ribonucleoprotein U

Table 2: hnRNP U is differentially expressed in whole blood from patients with sepsis.

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the fold change of hnRNP U in whole blood from patients with sepsis compared to whole blood from control subjects, gene and gene name are listed in this chart.

Figure 1: hnRNP U is expressed at significantly lower levels in whole blood from patients with sepsis.

The mRNA expression of hnRNP U in whole blood from control subjects (left) and in whole blood from patients with sepsis (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in HnRNP U expression between control whole blood and whole blood from patients with sepsis, a p-value, listed above.