Differential expression of heat shock protein family B (small) member 3 in human epithelial ovarian cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Epithelial ovarian cancer (EOC) is the most lethal gynecologic cancer (1). We performed discovery of genes associated with epithelial ovarian cancer and of the high-grade serous ovarian cancer (HGSC) subtype, using published microarray data (2, 3) to compare global gene expression profiles of normal ovary or fallopian tube with that of primary tumors from women diagnosed with epithelial ovarian cancer or HGSC. We identified the gene encoding heat shock protein family B (small) member 3, HSPB3, as among the genes whose expression was most different in epithelial ovarian cancer as compared to the normal fallopian tube. HSPB3 expression was significantly lower in high-grade serous ovarian tumors relative to normal fallopian tube. HSPB3 expression correlated with progression-free survival in patients with p53 mutant ovarian cancer. These data indicate that expression of HSPB3 is perturbed in epithelial ovarian cancers broadly and in ovarian cancers of the HGSC subtype. HSPB3 may be relevant to pathways underlying ovarian cancer initiation (transformation) or progression.

Keywords: ovarian cancer, epithelial ovarian cancer, HGSC, high-grade serous ovarian cancer, systems biology of ovarian cancer, targeted therapeutics in ovarian cancer.

The five-year survival rate for women diagnosed with high-grade serous ovarian cancer is between 30-40% and has not changed significantly in decades (4, 5). The development of novel, targeted therapeutics to treat HGSC can be facilitated by an enhanced understanding of the transcriptional behavior of ovarian tumors relative to that of the normal ovary. We mined published microarray data (2, 3) to compare global gene expression profiles between human ovarian tumors, including that of the HGSC subtype, and that of normal ovarian and fallopian tissue. We identified the gene encoding heat shock protein family B (small) member 3, HSPB3, as among the most differentially expressed in HGSC tumors of the ovary and in epithelial ovarian cancer broadly. HSPB3 may be a gene of interest when prioritizing the study of target genes and pathways for the development of novel therapeutic interventions in epithelial ovarian cancer and specifically in high-grade serous ovarian cancers.

Methods

We used microarray data from datasets GSE124766 (2) and GSE146556 (3) for this differential gene expression analysis of epithelial ovarian cancer in conjunction with GEO2R. The Benjamini and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of global differential expression. Log transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. GSE124766 (2) was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=3$ normal fallopian tube tissue (control) and $n=8$ for tumors from patients with high-grade serous ovarian cancer. GSE119054 (3) was generated using Agilent-067406 CBC lncRNA + mRNA microarray V4.0 with $n=3$ for normal ovarian tissue and $n=6$ for tumors from patients with epithelial ovarian cancer; analysis was performed using platform GPL19615. GEO2R provides mRNA expression levels only for the top 250 most differentially expressed genes transcriptome-wide.

A statistical test was performed to evaluate whether HSPB3 expression was significantly different when comparing normal ovarian tissue from control subjects and primary tumors from women diagnosed with HGSC using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses (Version 8.4.0)(455). For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool (6) for correlation of HSPB3 mRNA expression levels with progression-free survival (PFS) in $n=483$ ovarian cancer patients.

Results

We mined published microarray data (2, 3) to identify differentially expressed genes in epithelial ovarian cancer and specifically in high-grade serous ovarian cancer (HGSC), the type of ovarian cancer responsible for 70-80% of deaths resulting from the most lethal gynecologic malignancy.

HSPB3 is differentially expressed in ovarian tumors from women diagnosed with HGSC

We identified HSPB3 as among the genes whose expression was most different when comparing primary HGSC tumors to normal fallopian tube tissue (2) (Chart 1). When sorting all of the transcripts measured by microarray based on change in expression between HGSC and the normal fallopian tube, in this data set, HSPB3 ranked 29 out of 40481 total transcripts (Chart 1), equating to 99.9% differential expression. Differential expression of HSPB3 in HGSC tumors was statistically significant (Chart 1; $p=8.14E-08$).

We analyzed a second microarray dataset (3) generated using normal ovarian tissues and tumors from women diagnosed with epithelial ovarian cancer (EOC) to determine whether differential expression of HSPB3 could be observed in tumors from a different group of patients, which included HGSC tumors.

We found that HSPB3 was differentially expressed in epithelial ovarian cancers broadly (Chart 2). When sorting all of the transcripts measured by microarray based on change in expression between tumors from patients with epithelial ovarian cancers and the normal ovary, HSPB3 ranked 27325 out of 75217 total transcripts (Chart 2), equating to 63.7% differential expression. Differential expression of HSPB3 in EOC was not deemed statistically significant (Chart 2; $p=1.26E-01$).

HSPB3 is expressed at significantly lower levels in HGSC when compared to the normal fallopian tubes.

We then extracted exact mRNA expression levels for the differentially expressed HSPB3 transcript from both normal fallopian tube tissue and from epithelial ovarian tumors. HSPB3 was expressed at significantly lower levels in high-grade serous ovarian cancers when compared to the normal fallopian tube (Figure 1; $p=0.0048$). We calculated a mean fold change of 0.57 in HSPB3 expression when comparing HGSC tumors to normal fallopian tissue, as HSPB3 was expressed at 11.33 ± 0.88 arbitrary units (AU) in the fallopian tubes but at 6.46 ± 0.57 AU in tumors of the ovary.

HSPB3 tumor expression is correlated with survival outcomes in patients with ovarian cancer.

We performed Kaplan Meier survival analysis (6) to evaluate whether HSPB3 tumor expression correlated with survival outcomes in patients with p53 mutant (p53^{mut}) ovarian cancer. HSPB3 primary tumor expression was correlated with progression-free survival (PFS) in patients with p53^{mut} ovarian cancer, in the lower quartile (Figure 2). HSPB3 expression was a negative prognostic indicator in patients with p53^{mut} ovarian cancer. While median PFS was 21.37 months for ovarian cancer patients with low tumor expression of HSPB3, median PFS was 16.63 months for ovarian cancer patients with high tumor expression of HSPB3 (Chart 3). Correlation of PFS with HSPB3 tumor expression in patients with p53^{mut} ovarian cancer, in the lower quartile, was statistically significant (Figure 2: log rank p-value: 0.0076; hazard ratio: 1.43 (1.1-1.87)).

HSPB3 primary tumor expression was correlated with progression-free survival (PFS) in patients with p53^{mut} ovarian cancer, in the lower tertile (Figure 2). HSPB3 expression was a negative prognostic indicator in patients with p53^{mut} ovarian cancer. While median PFS was 20.47 months for ovarian cancer patients with low tumor expression of HSPB3, median PFS was 16.53 months for ovarian cancer patients with high tumor expression of HSPB3 (Chart 3). Correlation of PFS with HSPB3 tumor expression in patients with p53^{mut} ovarian cancer, in the lower tertile, was statistically significant (Figure 2: log rank p-value: 0.0023; hazard ratio: 1.45 (1.14-1.85)).

HSPB3 primary tumor expression was correlated with progression-free survival (PFS) in patients with p53^{mut} ovarian cancer, in the upper tertile (Figure 2). HSPB3 expression was a negative prognostic indicator in patients with p53^{mut} ovarian cancer. While median PFS was 19.23 months for ovarian cancer patients with low tumor expression of HSPB3, median PFS was 15.67 months for ovarian cancer patients with high tumor expression of HSPB3 (Chart 3). Correlation of PFS with HSPB3 tumor expression in patients with p53^{mut} ovarian cancer, in the upper tertile, was statistically significant (Figure 2: log rank p-value: 0.019; hazard ratio: 1.32 (1.04-1.66)).

HSPB3 primary tumor expression was not correlated with progression-free survival in the upper quartile (Figure 2: log rank p-value: 0.27; hazard ratio: 1.15 (0.9-1.49)) in patients with p53^{mut} ovarian cancer.

Thus, we found using published microarray data (2, 3) that HSPB3 was among the genes most differentially expressed in tumors from women with high-grade serous carcinomas, and in epithelial ovarian cancer broadly. HSPB3 expression was significantly lower in HGSC tumors when compared to the normal fallopian tubes, and HSPB3 tumor expression was correlated with progression-free survival in patients with p53^{mut} ovarian cancer.

Discussion

Transcriptional profiling of HGSC in a Cancer Genome Atlas integrated genomic analysis classified HGSC into four subtypes based on gene clustering: immunoreactive, differentiated, proliferative, and mesenchymal (7). We sought to describe the transcriptional landscape of epithelial ovarian cancers in an unbiased fashion by using published microarray data from primary tumors of women diagnosed with HGSC compared to transcriptome data from normal ovarian and fallopian tube tissue (2, 3). In both datasets analyzed (2, 3), HSPB3 was among the genes whose expression changed most significantly when comparing normal gynecological tissue to ovarian tumors, from women with epithelial ovarian cancer as well as specifically from women with HGSC. HSPB3 may be a molecule of relevance in understanding transformation or disease progression in human epithelial ovarian cancer.

References

1. Guppy, A.E., Nathan, P.D. and Rustin, G.J., 2005. Epithelial ovarian cancer: a review of current management. *Clinical Oncology*, 17(6), pp.399-411.
2. Zhang, W., Klinebiel, D., Barger, C.J., Pandey, S., Guda, C., Miller, A., Akers, S.N., Odunsi, K. and Karpf, A.R., 2020. Global DNA hypomethylation in epithelial ovarian cancer: passive demethylation and association with genomic instability. *Cancers*, 12(3), p.764.
3. Dong, S., Wang, R., Wang, H., Ding, Q., Zhou, X., Wang, J., Zhang, K., Long, Y., Lu, S., Hong, T. and Ren, H., 2019. HOXD-AS1 promotes the epithelial to mesenchymal transition of ovarian cancer cells by regulating miR-186-5p and PIK3R3. *Journal of Experimental & Clinical Cancer Research*, 38(1), pp.1-13.
4. Bowtell, D.D., Böhm, S., Ahmed, A.A., Aspuria, P.J., Bast Jr, R.C., Beral, V., Berek, J.S., Birrer, M.J., Blagden, S., Bookman, M.A. and Brenton, J.D., 2015. Rethinking ovarian cancer II: reducing mortality from high-grade serous ovarian cancer. *Nature reviews Cancer*, 15(11), pp.668-679.
5. Vaughan, S., Coward, J.I., Bast, R.C., Berchuck, A., Berek, J.S., Brenton, J.D., Coukos, G., Crum, C.C., Drapkin, R., Etemadmoghadam, D. and Friedlander, M., 2011. Rethinking ovarian cancer: recommendations for improving outcomes. *Nature Reviews Cancer*, 11(10), pp.719-725.
6. Györffy, B., Lánckzy, A. and Szállási, Z., 2012. Implementing an online tool for genomewide validation of survival-associated biomarkers in ovarian-cancer using microarray data from 1287 patients. *Endocrine-related cancer*, 19(2), pp.197-208.
7. Cancer Genome Atlas Research Network, 2011. Integrated genomic analyses of ovarian carcinoma. *Nature*, 474(7353), p.609.

Probe ID: A_23_P92730

p-value: 8.14E-08

t: 11.6628348

B: 8.33329

Gene: HSPB3

Gene name: heat shock protein family B (small) member 3

Chart 1: HSPB3 is differentially expressed in high-grade serous ovarian carcinomas (HGSC).

The rank of differential expression, the probe/transcript ID, the *p*-value with respect to global differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Probe ID: A_23_P45999

p-value: 1.26E-01

t: -1.6633045

B: -5.30177

Gene: HSPB3

Gene name: heat shock protein family B (small) member 3

Chart 2: HSPB3 is differentially expressed in epithelial ovarian cancers, including high-grade serous ovarian carcinomas (HGSC).

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

HSPB3

Figure 1: HSPB3 is expressed at significantly lower levels in HGSC tumors when compared to normal fallopian tube.

The mRNA expression of HSPB3 in normal fallopian tube tissues (left) and in the primary tumors of patients with HGSC (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in HSPB3 expression between the normal fallopian tube and primary tumors from patients with HGSC, a *p*-value, listed above.

Figure 2: HSPB3 tumor expression correlates with progression-free survival in patients with ovarian cancer.

Depicted in this Kaplan-Meier plot is the probability of progression-free survival (PFS) for n=483 patients, in the lower quartile (top left), in the lower tertile (top right), in the upper tertile (bottom left), and in the upper quartile (bottom right), based on primary tumor expression of HSPB3 (low and high). The log rank *p*-value denoting statistical significance of difference in progression-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on HSPB3 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Lower quartile

1 Low HSPB3 expression: 21.37 months

2 High HSPB3 expression: 16.63 months

3 **Lower tertile**

4 Low HSPB3 expression: 20.47 months

5 High HSPB3 expression: 16.53 months

6 **Upper tertile**

7 Low HSPB3 expression: 19.23 months

8 High HSPB3 expression: 15.67 months

9 Chart 3: Progression-free survival is superior in ovarian cancer patients with low tumor expression of
10 HSPB3.

11 The median progression-free survival of $n=483$ ovarian cancer patients based on stratification into low or
12 high tumor expression of HSPB3 is listed in this chart, in the lower quartile, in the lower tertile, and in the
13 upper tertile.