Introduction

Multiple proxies indicate that during the last two million years the East African climate changed, triggering a shift in its plant landscape from forested ecosystems to open woodland/grassland mosaics. These proxies include paleosol carbonates (Cerling et al., 1988; Levin et al., 2004; Quinn et al., 2007; Levin et al., 2011), vertebrates (Bibi and Kiessling, 2015; Bibi et al., 2018; Prassack et al., 2018), palaeobotanical remains (Bonnefille, 1984, 1995), and stable isotopes (Magill et al., 2013a, b; Uno et al., 2016; Lupien et al., 2018). This environmental transformation led to the widespread occurrence of a plant landscape characterized by *Acacia-Commiphora* woodland (White, 1983), and corresponds with key changes in hominin evolution and ecology (e.g. Ravelo et al 2004; Lepre et al, 2011, Beyene et al 2013).

Over the last few decades, different investigations have attempted to identify modern ecosystems as referential correlates for Pleistocene East African vegetation communities within the Somalia-Masai floristic region; a center of endemism that extends over 2 million km² in parts of Ethiopia, Sudan, Uganda, Kenya, and Tanzania. For example, lakes such as Manyara and Makat are often considered correlates for paleo-Lake Oldupai as they have vegetation types not in equilibrium with the regional climate (Barboni, 2014; Copeland, 2007). Peters and Blumenschine (1995) developed a paleolandscape model that predicted possible hominin land use patterns based on reconnaissance surveys in eastern and southern Africa. Copeland (2007) examined habitats in northern Lake Manyara, Ngorongoro, and the Serengeti Plain as baseline to understand the paleolandscape inhabited by early *Homo* in terms of climate, land forms, and soil types.

At present, there is no extensive, ecosystem-wide phytolith analog for the *Acacia-Commiphora* landscape that was the backdrop for human evolution in some regions of East Africa

over the last two million years (Bobe and Behrensmeyer, 2004; Plummer et al., 2009; Cerling et al., 2011; Blumenthal et al., 2017). Researchers have focused instead on a variety of microhabitats such as groundwater-fed woodlands (e.g. Ashley et al., 2010; Barboni et al., 2010), mosaic fluvial systems (e.g. Arráiz et al., 2017), and lake-shore palm groves (Albert et al., 2009, 2015).

In Quaternary palaeoecology, phytolith analysis has successfully addressed evolutionary and environmental questions (see review in Strömberg et al., 2018). Phytoliths are microscopic silicifications mirroring the morphometric characteristics of the plant organs where they precipitate, inside and between plant cells of varied groups (Rovner 1971, Piperno 1988; Hodson et al., 2005; Hodson, 2016; Strömberg et al., 2016). Phytogenic silica is released into soils after litter decomposes and discharges both amorphous silica and phytoliths. This process creates a synchronous phytolith record of the plant communities that grew in the soil (Piperno and Becker, 1996; Kerns et al., 2001; Clarke, 2003; Piperno, 2006; Thorn, 2006; Honaine et al., 2009; Morris et al., 2010; Mercader et al., 2011; Novello et al., 2012, 2018; Blinnikov et al., 2013; An et al., 2015; Watling et al., 2016; Esteban et al., 2017; Fick and Evett, 2018).

We explore the soil phytolith assemblages from *Acacia-Commiphora* woodlands to identify similar environmental contexts in the fossil record, undertaking a concurrent study of phytoliths from local plants to aid in taxonomic classification (Wallis, 2003; Carnelli et al., 2004; Gallego and Distel, 2004; Bremond et al., 2005; Tsartsidou et al., 2007; Barboni and Bremond, 2009; Mercader et al., 2009, 2010; Gao et al., 2018). The comparison of soil and plant phytoliths can assist in tracking the boundary between woodlands and grasslands (Kerns et al., 2001; Morris et al., 2010; Mercader et al., 2011). Elsewhere, this comparison has been helpful to assess phenomena such as time averaging (Alexandre et al., 1997; Blecker et al., 2006; Mercader et al., 2011; Hyland et al., 2013), catchment (Fredlund and Tieszen, 1994; Blinnikov, 2005), and

ecological succession (Leakey 1979; Niboye, 2010) that are hereby studied at the regional and local level.

Study area

Oldupai Gorge falls within the Northern Tanzania Divergence Zone along the East African Rift System (Dawson, 1992; Foster et al., 1997) (Fig. 1a). It represents the boundary between central Tanzania's Archaean craton and the north-south trending Mozambique belt, a product of the neo-Proterozoic Pan-African orogeny (Holmes, 1951; Cahen et al., 1984). Oldupai Gorge sits on the western flank of the Gregory Rift in the Ngorongoro Conservation Area (2°59'46.87"S, 35°21'7.50"E) between the volcanic highlands to the south and east, the metamorphic complexes to the north, and the Serengeti Plains to the west. Paleolake Oldupai was part of an endorheic basin formed > 2 Ma as a result of tectonic subsidence and concomitant uplift in the Ngorongoro Volcanic Highlands (Hay, 1976), where numerous volcanoes existed (Mollet and Swisher, 2012). Oldupai Gorge boasts well-dated archaeological and fossil records (Leakey et al., 1972; Hay, 1976; Deino, 2012). Its sedimentary beds span the Pleistocene (Beds I – IV, Masek, Nduku, and Naisiusiu) and situate two million years of evolutionary history, from the Oldowan to the Late Stone Age (Leakey, 1971; Leakey and Roe, 1994).

The Eastern Serengeti Plains (Fig. 1) offer a varied topography between 1350 m – 1550 m above sea level with a central plateau around Oldupai Gorge and highlands toward Endulen, enduring wide temperature oscillations and high winds (Anderson and Talbot, 1965; EAMD, 1967; Norton-Griffiths et al., 1975; White, 1983). Annual rainfall ranges from 380 mm to 803 mm per year, but at the Gorge it fluctuates from 331 mm – 531 mm (Herlocker and Dirschl, 1972). The

dry months in Oldupai Gorge extend from June to October, with the wettest phase lasting from December to April. Rainfall patterns and localized soil conditions influence the vegetation communities observed in the Serengeti (Anderson and Talbot, 1965). The dominant soils are calcimorphic, yellowish brown, with low humic values ($< 1\%$ organic carbon) and a hard pan (Anderson and Talbot, 1965).

Materials and Methods

The Tanzania Commission for Science and Technology (permit 2018-112-NA-2018-36), and the Tanzanian Ministry of Natural Resources and Tourism, through its Antiquities Division (permit 14/2017/2018) approved this work. Authorities at the Ngorongoro Conservation Area allowed us to enter the protected area (BE.504/620/01/53). The export license for the materials presented in this study were obtained from the Antiquities Division (EA.150/297/01: 5/2018/2019) and the Tanzanian Executive Secretary from the Mining Commission (00001258).

Vegetation Survey

The greater Serengeti area falls within the Somalia-Masai floristic region in which *Acacia-Commiphora* deciduous bushland is the climax vegetation (Kindt et al., 2011). This ecoregion is transitional to miombo woodlands to the south (Frost 1996) and the Zanzibar Inhambane coastal forest mosaic to the east (Lovett and Wasser, 1993). The published plant inventory for Oldupai Gorge totals 32 genera (Herlocker and Dirschl, 1972). We studied 29 species representing 20 of these genera and 15 families. Characteristic Somalia-Masai taxa analysed here comprise the

dominant members of the canopy (*Acacia mellifera*, *A. nilotica*), emergents (*Boscia angustifolia*), bushes (*Barleria eranthemoides*, *Maerua triphylla*), the herbaceous layer (*Hypoestes forskalii*, *Ocimum* spp.), succulents (*Sansevieria robusta*, *Aloe secundiflora*), and the grasses (*Sporobolus consimilis*, *Cynodon dactylon* -Chloridoideae, *Aristida adoensis* -Aristidoideae).

Vegetation in the greater Oldupai region has been monitored for decades (Niboye 2010). The average perennial cover today is 22% (Niboye, 2010), which is comparable to that observed several decades ago (Anderson and Talbot, 1965), with bare terrain exposing > 45% of the area. Herlocker and Dirschl (1972) noticed an increased plant variety in the eastern sector of the gorge, introduced by the topographic changes from the canyon walls, and thus a higher diversity of bushes and grasses. They also noticed commonalities with the communities from the western gorge.

Ecologically, the plant community that typifies the natural vegetation at Oldupai Gorge is short *Acacia* and *Commiphora* woodland. Several tree species from these two genera co-dominate along with bushes such as *Barleria eranthemoides*. Succulents abound in ravines and depressions where several species of *Sansevieria*, *Euphorbia*, and *Sarcostemma* are common. Herbaceous plants are variably scattered across the landscape but, overall, they are low in numbers. Importantly, this domain is characterized by a paucity of grass taxa, which are typically inconspicuous and occasional (Kindt et al., 2011).

At the regional level (Fig. 1b) the vegetation was surveyed using qualitative criteria, aimed at investigating the range of species present in the greater Oldupai region, learning about their phytolith production and morphotype characteristics, and comparing the botanical dataset with the soil group. We did not attempt to collect plant specimens in numbers proportionate to frequency in each of their respective ecological communities, nor to other physiognomic variables or phylogenetic affiliation. Instead, we collected representative plants from the western and eastern

gorge at the end of the wet season into the dry season in 38 sampling locations to target 29 species along an E-W transect (25 km long) from Granite Falls to Olbabal. Field samples were identified in the National Herbarium of Tanzania at the Tropical Pesticides Research Institute. When species level identification was uncertain and could belong to several species within the genus, we listed it as spp. Grouping and nomenclature come from the index of accepted names for the flowering plants of Sub-Saharan Africa (Klopper et al., 2006).

At the local level, the vegetation was surveyed using quantitative criteria. First, an aerial image covering 1 km² (Fig. 1c) taken at the end of the wet season in the eastern half of Oldupai Gorge, was georeferenced via ArcGIS - ArcMAP 10.5.1. A grid consisting of 100 cells, each 100 m², was superimposed. The cover-abundance value was digitized in each cell as a polygonal shape file, and classified it in four classes or ranks:

- 1) Rank zero, or 'No-cover': when plant cover constituted < 10% within the spatial polygon (45.64% of the total area).
- 2) Rank one, or 'Sparse': when plant growth was 10% - 20% (20.32% of the total).
- 3) Rank two, or 'Open': when plant growth was 20% – 60% (26.41% of the total).
- 4) Rank three, or 'Closed': when plant growth was > 60% (7.63% of the total).

This rank system was subjected to groundtruthing in 300 spots (Mercader et al 2017: Fig. 2), and then we proceeded to conduct a qualitative vegetation survey of the entire 1 km² plot (Fig. 1c). After this, we zoomed into a 62,500 m² quadrat (Fig. 1d) to conduct a full inventory, plotting, and collection of all plants present within this smaller plot. According to plant cover rank (Fig. 2), thirteen samples (37.14%) were recovered from rank zero; eight (22.86%) from rank one; eight (22.86%) from rank two; and six (17.14%) from rank three. Samples were each assigned to one of two topographical categories: promontory or ravine. Ravines are at a mean elevation of 1,422.66

m with a range between 1,412 m and 1,441 m (29 m). Promontories are at a mean elevation of 1,452.41 m with a range between 1,416 m and 1,469 m (53 m). Most samples ($n = 29$, 82.86%) were recovered from promontory locations (Supplemental Figure 1).

Particle Size Analysis

We carried out particle size analysis of selected soil samples ($n = 10$) via laser diffraction (Malvern Mastersizer 2000: Sperazza et al., 2004). A soil sample (1.0 g) was deflocculated with 0.1% sodium hexametaphosphate. Ultrasonics dislodged clays for 5 minutes. Sample was left on orbital shaker overnight at 200 rpm. We cleansed the sample of carbonates, phosphate, and organics via hydrochloric acid (HCl 37%), nitric acid (HNO₃ 60%) and hydrogen peroxide (H₂O₂ 30%) successively, rinsing the sample three times via centrifuging in between each treatment (3,000 rpm, 5 min). When measuring particle size in the Malvern 2000, we used obscuration rates above 15% but below 30%. The measurement of pH in all soils was through a ThermoFisher Orion 320 PerpHecT LogR meter, calibrated daily. The Ag/AgCl electrode was rinsed after calibration and in between samples. Sediment color was estimated dry under constant light conditions by two observers relative to the Munsell color system. Water loss on drying is the balance between soil field moisture in three grams of soil and the mass of the sample after drying at 110 °C for 24 h.

Soil Phytoliths

Sampling was never conducted on archaeological sediments, nor within any of the well-known and mapped lacustrine and fluvial outcrops that constitute the ancient beds (Leakey et

al., 1972; Hay, 1976); our work was restricted to purely surficial sandy loam (Anderson and Talbot, 1965). Prior to collecting soil samples, and to better understand local soils and visualize their horizons, several trenches were dug reaching the C-horizon: a characteristic underlying hard pan described by earlier soil analysts (Anderson and Talbot, 1965; Jager 1982). In section, we observed a thin A-horizon, restricted to the rhizosphere and litter zone, while a B-horizon seems absent. We scraped 25 cm² off the surface (5 ml) to a maximum depth of 1cm within the A-horizon (Fishkis et al., 2009; Mercader et al., 2010), excluding litter and not mixing or generating composite samples. Thirty-five loci were selected for stratified sampling, so that a comparable number of samples were collected from each plant cover type. The median distance between all soil assemblages is 516 m (measured distances, n = 522) over a range of 13 m – 1,252 m. Geographical coordinates were documented as point features. The full dataset of soil samples including provenance, elevation, coordinates, processing weights, and individual counts are available at the Federated Research Data Repository (doi:10.20383/101.0122)

Detailed phytolith extraction protocols are available with the Federated Research Data Repository (doi 10.20383/101.0123), which are modified from Albert et al., 1999. In brief, extraction from sediment used a 3.0 g aliquot that received 10 ml of 0.1% sodium hexametaphosphate for clay dispersion, followed by sonication (15 minutes) and overnight orbital shaking plus decantation of clays. Samples were rinsed and dried over two days (> 70 °C). Soil samples received 10 ml equal parts 3N hydrochloric and nitric acids; then, rinsed and centrifuged at 3,000 rpm. Drying followed for two days (70 °C). Samples were weighed and transferred to Petri dishes on a hot plate (70 °C) whereby 10 ml of hydrogen peroxide (30%) was added to remove organics. Rinsing, drying, and weighing followed. Phytolith separation was done by transferring this fraction to 15 ml centrifuge tubes, to which 5 ml sodium polytungstate (~2.4 s.g.) was added.

Samples were vortexed for five minutes at 3,000 rpm. The supernatant was removed. A gradual decrease in specific gravity over the course of three subsets (subfractions 1, 2, 3) concentrated phytoliths in the third extract. Samples were rinsed with de-ionized water before centrifuging twice more, followed by two cycles of high centrifugation (4,500 rpm). Samples dried at 70 °C another time and a final weight obtained. The aliquot extracted for phytolith analysis weighed approximately 0.001 g. It was mounted with Entellan New and inspected while fresh; allowing particles to be rotated in three dimensions under the microscope. Phytoliths from soil samples were counted on one slide over 600 mm² to the total amount, reaching average counts of 250. Diatoms were tallied concurrently. Carbonate content was calculated as the loss on acid treatment relative to the original mass. Extraction weights are the balance between the original mass and that of the extract after defloculation, carbonate removal, destruction of phosphate, elimination of organics, rinsing, and drying.

Plant Phytoliths

Extraction followed Albert and Weiner (2001), with modifications. Botanical specimens were sorted into anatomical parts, and each was individually weighed and measured. Full datasets with species, plant part studied, provenance, elevation, coordinates, processing weights, and individual counts are available at the Federated Research Data Repository (doi:10.20383/101.0122). All samples were cleansed by ultrasonication, then rinsed and dried overnight > 80 °C. Dry weight was obtained, and samples placed in a muffle furnace (24 h, 400 °C). The resulting ashes were weighed and transferred to Petri dishes to which 10 ml of 3N hydrochloric and nitric acids were added to a hot plate at 70 °C for 30 minutes. Samples were

transferred to 15 ml tubes and rinsed of remaining acids in three wash and centrifugation cycles at 3000 rpm, with excess water being boiled off. Hydrogen peroxide (10 ml, 30%) was added to the Petri dishes heated on hot plates (70 °C), transferred to 15 ml centrifuge tubes, rinsed, and centrifuged (3,000 rpm) for three cycles. Samples were dried > 70 °C overnight, and then weighed. An aliquot of approximately 0.001 g was analysed after mounting the sample for microscopy with Entellan New. Inspection captured three-dimensional characteristics during rotation conducted within 48h of mounting and before the resin dried. Frequent polarization of the field of view under the microscope ensured that no anisotropic particles and crystals were confused for phytoliths. Neither fresh nor weather volcanic glass could be observed. We counted one full slide and only discrete phytoliths (average = 106 phytoliths), while articulated shapes were counted as one. As for very rich extracts, we tallied a minimum of 200 phytoliths and/or short cells for the grasses.

Phytolith Classification

There are two approaches to environmental reconstruction in phytolith research: the indices (Barboni et al., 2007; Bremond et al., 2008) and the general approach (Strömberg, 2004). Different researchers adopt one approach or another depending on their work's practice and scope. We followed the general approach here to detect plant variations at the spatial level. Quantitative analysis involved exploration aimed at discovering relationships amongst the soil and plant phytoliths, which, when combined and filtered, isolated groups, classes, and morphotypes unique to each set and those in common. Further investigation involved comparison of phytoliths from soils and plants to explore their relative distributions through Chi-square independence tests,

establishing the statistical significance of the relationship between categorical variables within the two populations using a 95% confidence value ($p = 0.05$).

Two system, polarizing microscopes (Olympus BX51, Motic BA410E) allowed us to confirm the opal silica's isotropy and tally phytoliths at 400x, following the International Code for Phytolith Nomenclature 1.0 to name and describe specimens (Madella et al., 2005), with exceptions duly noted. Geographically pertinent reference collections and published soil phytolith assemblages were consulted (Albert et al., 2006, 2015; Bamford et al., 2006; Fahmy, 2008; Mercader et al., 2009, 2010, 2011; Barboni and Bremond, 2009; Eichorn et al., 2010; Novello et al., 2012; Collura and Neumann, 2017; Neumann et al., 2017). Full datasets with individual counts for each morphotype are available at the Federated Research Data Repository (doi:10.20383/101.0122). We classified discrete phytolith shapes in four large groups (woody dicot, grass short cells, generic Poaceae, and rare) and 17 classes that encompass 76 types. The woody dicot group totalled the blocky class (Mercader et al., 2009; Novello et al., 2012; Collura and Neumann, 2017), cylindrical types from bark (Mercader et al., 2009), varied globular phytoliths (Runge, 1999; Mercader et al., 2009), tabular types from woody dicots (Mercader et al., 2009; Collura and Neumann, 2017), clavates (Mercader et al., 2009), and sclereid phytoliths (Collura and Neumann, 2017). Phytoliths from Poaceae short cells were subdivided in three classes: lobate (Fahmy, 2008; Neumann et al., 2017), rondels (Mercader et al., 2010; Novello et al., 2012; Neumann et al., 2017) and chloridoid saddles (Twiss et al., 1969; Mercader et al., 2010). Two additional groups comprised generic grass phytoliths and rare morphotypes.

It should be noted that the tabular class, with 12 shape variants, contains a confuser morphotype called here tabular *sensu stricto*, and defined as a narrow, thin parallelepiped shape with straight edges that may occur in both monocots and dicots. This confuser morphotype differs

from the tabular psilate observed in woody dicots (this study) in that the latter is a wide and thick tabular shape with a very slight sinuosity to the outline.

Results

Soil Phytoliths

Soil Characteristics and Phytolith Abundance

Local topsoils (Table 1) have a mean particle size of 344 μm , with pH of 8.7 and 5% water content. Unimodal particle size distribution is apparent in the poorly sorted soil matrices of most samples, while the coarse sand component in samples from the eastern half of our plot generates a bimodal population in which the poorly sorted fines represent the soil, and the well-sorted, coarse sands originate from easterly aeolian import (Fig. 3). Silt and very fine sands average 34% per sample, while carbonates are 69% by mass. The extracted phytolith-bearing fraction, which includes both biogenic and geogenic silica, is 54.53% ($M = 53.85\%$; range = 45.54%, 25.23% – 70.77%). Dissolution and mechanical breakage were rarely observed, and caution must be used not to confuse the effect of dissolution causing holes randomly through a phytolith versus the cavate textures that heavily decorate some blocky and tabular morphotypes (Fig. 4q,w) from woodland taxa (cf. Mercader et al., 2009: Fig. 6g,k,l,m).

No soil sample was void of phytoliths. In total, 10,745 phytoliths were tallied ($\bar{X} = 307$, $M = 285$ per sample; range = 839, 16 – 855) containing 64 morphotypes that can be grouped into 15 major classes (Fig. 4, Table 2). Of these, six are prominent classes (Fig. 5, 6) and account for 83.78% of all soil phytoliths: tabular: 41.74%, blocky: 13.11%, globular: 8.52%, rondel: 6.98%,

cylindric: 6.96%, and lobate: 6.47%. Within the tabular class, the three most prominent contributors represent 82.16 % of the subtotal: scrobiculate: 45.40%; sinuate: 20.47%; and thick lacunate: 16.30%. Two types of blocky phytoliths represent 91.12% of the class; blocky: 62.38% and ridged: 28.74%. One single type of globular phytolith represents 85% of all globular types: a globular to hemispherical, facetate body with a centric cavity; possibly from mesophyll cells found in the leaves from several taxa in the Lauraceae (Fig. 4ad; Thorn, 2001; Murungi, 2017). In contrast, the classic indicator of arboreal cover in the African tropics, the globular granulate, amounts to only 9% of the globular class. Two globular morphotypes were recovered from soils only: globulose bisected and segmented. Within the cylindric class, two phytoliths reach 91.44%: scrobiculate, 76% and sinuate, 15.37%.

Two types dominate the rondels (97.33%): Towers, 63.33% and rondels *sensu stricto*, 34%. Lastly, for the lobate class, 86.33% of the variability is represented by small bilobates with concave ends (61.58%), small bilobates with flat ends (16.04%), and small bilobates with convex ends (8.34%). Four lobate types were found exclusively in soils: small bilobate with notched ends, large bilobate with flat ends, large bilobate with notched ends, and cross. The saddle class is uncommon (5.87%) and the short saddle dominates it (74.32% of the subtotal), while squat saddles are rare. Generic tabulars (3.88% of the total), bulliforms and lanceolates (2.89% of the total), generic cylindric phytoliths (1.23% of the total), sclereids (1.1% of the total), fusiforms / gutiforms (0.74% of the total) (nomenclature not per ICPN 1.0), and clavates (0.25% of the total) are scarce. The remainder includes composite epidermal tissue, polygonal prisms with subrounded top (cf. *Commelinaceae*), and papillae; and they all are extremely uncommon.

Soils also produced 1,564 diatoms in addition to the phytolith complement summarized above. The mean number of diatoms per sample is 44.68 and the median recovery is 9.50 with a

range of zero – 301. The specimens are mostly from the Nitzschoid group (Fig 4af-ah), which grows in wet soils. The diatom count per plant cover class is inversely proportional to density of plant cover today, with the highest frequency in soil samples without vegetation and/or sparse growth probably because of pooling water in these areas (Albert et al 2006).

Extraction Weight and Plant Cover

Conventionally, extraction weights are reported as a proxy of phytolith concentration. Like others before us (Blinnikov et al., 2013), however, we did not see a discernible pattern between plant cover, phytolith count, and extraction weight (Supplemental Figure 1) with values randomly distributed. Thus, plant cover rank zero varied between 43.6% - 69.5%, for rank one between 40.96% - 70.77%, rank two between 25.23% - 63.56%, and rank three between 41.1% - 56.16%. The scores for plant cover ranks overlap, and so do those for extraction weight versus terrain elevation.

We explored the relationships between plant cover rank and phytoliths from woody dicots versus grasses. The woody phytolith group dominates all terrain ranks, and appears to increase with denser plant cover, while grass phytoliths peak in sparsely vegetated terrain. The null hypothesis posits that if plant cover has no influence on the relative proportions of woody and grass phytoliths, there will be no significant difference between overall proportions and plant cover rank proportions. For plant cover rank zero, vegetation density has not significantly influenced the relative proportions of woody and grass phytoliths present in the topsoil (Chi-square = 2.382, not significant at $p < 0.10$). There are significant differences in all other plant landscapes: Rank one has significantly more grass phytoliths than expected from the null hypothesis (Chi-square = 9.0812, $p < 0.01$) while ranks two (Chi-square = 7.2288, $p < 0.01$) and three (Chi-square = 8.0625,

$p < 0.01$) have many more woody phytoliths. Statistically, therefore, sparse, open, and closed terrain influenced the relative proportions of woody and grass phytoliths released into the soil (Table 3), with sparse land containing more grass phytoliths, and open woodland / closed terrain supporting more woody dicot types.

For short cell classes (Table 4), rank zero yields rondels followed by lobates and saddles; rank one: lobates, saddles, and rondels; rank two and three: lobates, rondels, and saddles. These class fluctuations across plant cover, however, are artifacts of differing sample numbers per plant zone (Supplemental Figure 2-4). Chi-square testing revealed no significant changes in short cell class related to plant cover rank. By morphotype, the most frequent phytoliths are towers, short saddles, and small bilobates with concave ends. In ranks zero and one, this order is preserved, but it reverses in ranks two and three. That is, the tandem of towers and short saddles are more frequent in areas of sparse to no plant cover (rank zero, one), while short saddles and small bilobates with concave ends are more frequent in areas of increased plant cover (rank two, three). However, Chi-square testing shows no significant change in the relative frequencies of short cell morphotypes related to plant cover today. In fact, where one phytolith type increases in frequency, so do the other two: These morphotypes vary positively with each other.

Plant Phytoliths

Production

All plant samples contained phytoliths, except for the woody tissue of *Commiphora kua*. A total of 4,310 phytoliths were classified (Table 5, 6) in morphotypes ($n = 52$) and classes ($n = 13$). The median number of phytoliths per species is 130 (range: zero – 539) and were found mostly

in leaf material (62.01%) with specimens from woody tissue making up the second largest group (23.52%) (Fig. 7, Supplemental Table 1). The number of samples from leaf and woody tissue was identical at nineteen each. Leaf samples produced on average two and a half times as many phytoliths, with the exception of *Balanites aegyptiaca*, *Melhania parviflora*, and *Ocimum* spp. in which woody tissue generates more phytoliths. Where extraction weights of phytoliths can be compared between leaf and woody tissue from the same species ($n = 13$), leaf samples have a slight tendency to produce larger extraction percentages (8/13).

We do not see a correlation between high phytolith production and the number of morphotypes per taxon (Table 7). The average yield per morphotype is 83 phytoliths. Three types top the highest frequency with figures six times higher than the average: oblong granulate, small bilobate with concave ends, and rondel ovate. In second place, tabular, globulose, vessels, and short saddles. Average richness is noticed for tabular psilate, globular granulates, epidermal cells, hair complex phytoliths, mesophyll cells, and large bilobates with convex ends. Also included in this group are rondel tower, rondel horned, and blocky. All other morphotypes appear at below average frequency. Thus, highly polymorphic species generate average phytolith amounts, and low polymorphic taxa can be top silica accumulators; however, all but one species among the poor phytolith producers (*Balanites aegyptiaca*) fall within the low polymorphism group, probably an artifact of small sample size.

Richness

Five taxonomic groups can be distinguished:

1. Top producers rank at median values almost three times higher than the rest of the group (~332 phytoliths per mount). The lead type is *Boscia angustifolia* (Capparaceae), along

with three members of the Poaceae representing three subfamilies, Panicoideae, Chloridoideae, and Aristidoideae. In this area, grassy species produce many more phytoliths than most other plants, outproducing other functional classes by factors ranging between 1.63 and 27.83.

2. High values are also reported for a group of five species with median phytolith number of 218. Four of these species belong with the Chloridoid grasses. One member is with the Olacaceae (*Ximenia caffra*).
3. Average production was noticed in two members of the genus *Acacia* (*tortilis*, *nilotica*) (Fabaceae). Other notable producers include *Melhania* (Malvaceae), *Maerua* (Capparaceae), *Cissus* (Euphorbiaceae), and *Barleria* (Acanthaceae).
4. Low producers with a median of 104 phytoliths include *Ocimum* (Lamiaceae) and *Commiphora* (Burseraceae), and members of the Acanthaceae, Asphodelaceae, and Asparagaceae.
5. Very poor phytolith producers (median value = 19) comprise some of the most characteristic taxa in the region; namely, *Acacia mellifera*, *Balanites aegyptiaca*, three *Commiphora* species, the most emblematic taxon of Oldupai Gorge (*Sansevieria robusta*), *Salvadora persica*, and the genus *Sarcostemma* (Apocynaceae): The lowest absolute rank in the region is that of *Sarcostemma viminale*.

Morphotypes

The morphotypes from woody plants are led by the globular class through three phytoliths: Oblong shapes with a granulate texture, globulose, and globular granulate. The tabular class includes three morphotypes in abundance: scrobiculate, sinuate, and ellipsoidal; while the

cylindrical class peaks with cylinder sinuates and psilates. Blocky morphotypes are relatively common, but four blocky morphotypes could not be found in the local plants we studied: corniculate, ridged, sinuate, and thick lacunate. The Poaceae produce various rondel types, dominated by ovates, towers, and horned towers. Lobate phytoliths comprise six variants but are mostly manifested through two types: small bilobates with concave ends (e.g. Panicoid type) and large bilobates with convex ends (e.g. Aristidoid type). The saddle class comprises two types largely dominated by short saddles both being very common in *Cynodon dactylon*. Several cylindrical phytoliths were documented in thirteen local dicots but were absent or extremely infrequent among the Poaceae: cylindrical crenate, psilate, scrobiculate, and sinuate.

Discussion

The Acacia-Commiphora Analog

In East Africa, the transition to arid-adapted woodlands and grasslands over the last two million years has been documented in pedogenic carbonates (Cerling et al., 1988; Levin et al., 2004, 2011; Quinn et al., 2007), mammal, bird, and fish fossils (Bibi and Kiessling, 2015; Bibi et al., 2018; Prassack et al., 2018), micro- and macrobotanical remains like pollen, phytoliths, and wood (Bonnefille, 1982, 1995; Barboni et al., 1999), and stable isotopes of hydrogen and carbon from leaf wax lipid biomarkers (Magill et al., 2013a, b; Uno et al., 2016; Lupien et al., 2018).

The phytoliths from present-day vegetation and soil surfaces are the baseline to interpret palaeolandscapes and ecological niches that might have been occupied by early hominins. To study the influence of plant community structure and cover on human ecology, soil phytolith

assemblages under known vegetation density and composition can provide a referential to interpret key palaeoecological aspects of early Pleistocene floristic regions and their main vegetation types. We provide a phytolith analog for *Acacia-Commiphora* mosaics that may represent the trend for more open landscapes recorded in East Africa since the early Pleistocene. In this analog, the cohort of phytolith morphologies that identifies semiarid woodland include tabular scrobiculate, tabular sinuate, blocky, globular facetate, tabular thick lacunate, cylinder scrobiculate, tabular sulcate, tower, short saddle, small bilobate with concave ends, and blocky ridged. We did not find unique identifiers or highly diagnostic shapes for the most typical species that today make up the *Acacia-Commiphora* ecosystem, but this is hardly surprising and confirms similar observations in adjacent regional centers of plant endemism, such as the Zambezian woodland, where woody dicot types have very little taxonomic power (Mercader et al., 2009) because of redundancy across orders (Bozarth, 1992). Also in common with phytolith spectra from plants and soils in Zambezian woodlands (Mercader et al., 2009, 2010, 2011), there is severe overrepresentation of robust morphotypes such as tabulars, blockies, and globulars in addition to lobate short cells.

Do Soil Phytoliths Mirror Physiognomy and Composition of Vegetation Aboveground?

We address this question at two levels, regional (plot = 100 ha) and local (quadrat = 6.25 ha). When considering the larger regional plot, all soil samples are dominated by phytoliths from woody dicots (Supplemental Table 2-4) (Albert et al., 2006):

The blocky morphotypes that could not be found in local plants (corniculate, ridged, sinuate, and thick lacunate) were partly tracked by geo-referencing: Blocky ridged and thick lacunate associate with areas of sparse plant growth (Supplemental Figure 5-6). Large numbers of

these forms in soils suggest that plants not represented in our inventory might have produced them or that the species producing these types have disappeared in past decades or not seasonally available when we collected our samples.

Cylindrical phytoliths were common in many of the local but were either absent or extremely infrequent among the grasses. Cylinder corniculates and thick lacunates occur only in the soils, while crenates occur only in plants. Cylinder crenates appear in *Acacia mellifera* and *Ocimum* spp. Similarly, the few corniculates recovered from soils may not be inferentially important. However, the relatively large number of cylinder thick lacunates in soils is noteworthy, as plants from the Miombo woodlands (Mercader et al., 2009) produce them. Fusiforms, guttiforms, clavates, and sclereids are found in plants and soils.

The globular class is common in both soil and plant samples. The globular facetate (mesophyll cells) is very abundant, and geo-referencing shows an association with areas covered by plants and absence from bare places (Supplemental Figure 7). Two globulose morphotypes were recovered from soils only: globulose bisected and segmented. The former is a unique phytolith that, in adjacent Zambezian woodlands, is restricted to the Solanaceae (Mercader et al 2009: Fig 3 ae).

Tabular phytoliths are relatively uncommon in the plants but very common in the soils. Four morphotypes could not be found amongst the plants, but were in soil samples: corniculate, ellipsoidal large, facetate, and pilate. Their georeferencing shows no pattern for three of them, but one type (tabular ellipsoidal large) appears solely in locations with plants growing today.

Poaceae phytoliths are skewed in their soil expression (Table 8, Supplemental Table 5): the local grasses are demonstrated overproducers (Twiss et al., 1969; Novello et al., 2012) but they are not overrepresented in these soils, rather the opposite. We infer that topsoils embody plant

communities, past and present, in a time-averaged palimpsest dominated by woodlands (Supplemental Table 3). It is likely that the heterogeneous landscape and the changing grazing history, erosion, and fire regime by pastoralists over the last 60 years (Sinclair et al 2008) have created a phytolith record of ecological succession underscored by a reversion towards a dominance of woody vegetation, as independently recorded by studies of land cover change (Sinclair et al 2007, Niboye 2010) that denote a low human impact on ecosystem dynamics since the 1960s.

Within the short cells, three major classes were identified in plants and soils: lobates, rondels, and saddles; albeit saddles are overrepresented in soils. Ten phytolith morphotypes were recovered within the lobate class. Of these, six were found in both plant and soil samples and four were found only in soils (small bilobate with notched ends; large bilobate with flat ends; large bilobate with notched ends; cross). The lead lobate type (Fahmy, 2008; Neumann et al., 2017), small bilobate with concave ends, is produced abundantly in one local and one regional taxon: *Pennisetum mezianum* and *Sporobolus africanus* respectively. Aristidoid, large bilobates with convex ends are documented in *Aristida adoensis* while small bilobates with flat ends appear in *Pennisetum mezianum*. Rondel class phytoliths are common in both plants (*Sporobolus* specimens) and soils. Rondel ovates are unique to the plant sample, while truncated rondels occur only in soils. Rondels and horned rondels appear in *Sporobolus* spp. The absence of rondel ovates in the soils is notable given their preponderance in the genus *Sporobolus*. Saddles were recovered from three plants. Saddle squats are present in *Cynodon dactylon* and the abundance of this morphotype in the soil samples may suggest correlation. Short saddles are found in plants and soils while saddle ovates are from soils but not plants. In brief, short cell phytoliths from soils do not always reflect the grass species growing in them today.

Catchment

We explore the notion of catchment by zooming into one local quadrat covering 6.25 ha to conduct a side-by-side comparison of phytolith assemblages from topsoils ($n = 6$) and plants ($n = 18$) (Table 8, Supplemental Table 4-5). To do this, we tallied all phytoliths by class. Of the thirteen phytolith classes represented in this quadrat, eight classes are severely underrepresented in the soil, including all three Poaceae short cell classes, two woody classes (globular/fusiform), epidermal cells in anatomical connection, and vessels (Bamford et al., 2006). Specifically, a positive balance exists for cylindrical shapes from woody tissue, along with sclereids and clavates. Moreover, dicot tabular morphotypes are overrepresented in the soil by a factor of seven. Several blocky morphotypes from woody tissue are four times more common in soils than in the plants that produced them, and lack of correspondence was similarly noticed for bulliforms and lanceolates. We attribute this imbalance to the phytolith signal from grasses being dampened by an overwhelming accumulation of woody phytoliths.

Rich phytolith producers such as the Poaceae display differential representation in soils, especially at sampling loci where they were growing. For instance, *Cynodon dactylon* produces short and squat saddles. In the quadrat where it grows today the average frequency of short saddles is half ($n = 7.5$) of the local average ($n = 13$), and the same applies to squat saddles (this quadrat = 1.8 versus 3.5 in the rest of the collection). *Sporobolus panicoides* preferentially generates rondel ovates, but none could be found in any of the six soil samples surrounding it. *Aristida adoensis* supports a majority of large bilobates with convex edges, and the average representation in the plant ($n = 0.2$) is similar to that in the soils nearby ($n = 0.3$). Importantly, one of the highest ranked

short cells from both collections, soils and plants, the small bilobate with concave ends, attains lower than average representation in the soils under scrutiny ($n = 4.8/6$) perhaps because the main producers of this morphotype (*Pennisetum mezianum*, *Sporobolus africanus*) were not close.

Considering that all phytolith classes found in the living tissues of the eighteen plant taxa growing within this quadrat are recorded in the underlying soil surface it can be suggested that the catchment area for these soil phytolith assemblages is smaller than five hectares (Blinnikov, 1994; Fredlund and Tieszen, 1994; Blinnikov et al., 2002).

Ecological Disturbance, Time Averaging, Heterogeneity, and Sampling Methods

The soil and plant phytolith assemblages are clearly distinguishable from one another. The presence of rare group phytoliths influences ratios in plants (26.98% of the total assemblage) and is much less an aspect of the soils (4.11%). However, even if we were to remove rare and delicate morphotypes, and consider short cells and woody dicot phytoliths only, the differences between the two datasets are obvious. In the plants, the two major groups are near parity (ratio = 1.031; Poaceae $n = 1556$ / woody $n = 1509$) while in the soils the two groups are quite disparate in relative frequency (ratio = 0.263; Poaceae $n = 1509$ and woody $n = 7914$).

Soil samples are not homogenous relative to one another. We indicated statistically significant relationships between the soil phytolith population in samples collected under a given plant cover and the rank of that zone, although not all individual samples show this relationship. The ratio between Poaceae short cells and woody phytoliths varies widely from 1:1 to 0:1. Moreover, soil samples with high number of Poaceae short cells have the highest number of woody phytoliths. Even in the most open parts of the landscape phytoliths from woody tissue outnumber

those from grasses (Gao et al., 2018) and some woody types abound in soils but are absent from the local plants sampled (Albert et al., 2015).

We are suggesting decadal inheritance of soil phytoliths, as evidenced by:

- 1) The highest phytolith recovery is from areas without plant cover or Rank Zero.
- 2) No significant change in the relative frequencies of short cell morphotypes related to plant cover today.
- 3) Multiplicity of soil phytolith types, woody and grassy, that cannot be traced back to local plants.
- 4) Several phytolith classes are overrepresented in the soils.

That is, although soil phytoliths might represent ‘decay-in-place’ (Piperno, 2006), as well as transport by wind, water, fire and herbivory (Fredlund et al., 2001), the two phytolith populations reported here, from plants and soils, will always differ from each other because soil assemblages are the sum of phytolith accumulations and losses over time (Fredlund and Tieszen, 1994; Kerns et al., 2001).

Conclusion

Over the last few decades, different environmental proxies have been utilized in Africa to identify present-day ecosystems that could be used as referentials to frame the emergence of the genus *Homo* (Cerling et al., 1988; Barboni et al., 1999, 2010; Quinn et al., 2007; Albert and Marean, 2012; Barboni 2014; Esteban et al., 2017; Novello et al., 2018; Prassack et al., 2018). One of these analogs is the *Acacia-Commiphora* biome, where we studied a phytolith reference collection of characteristic plants that aided in the taxonomic identification of phytoliths from the

underlying soils. This paper also studied the resolution of phytoliths from present-day surfaces along the *Acacia-Commiphora* landscape to reflect woody dicot versus grassy plant cover, detecting the shifting boundary between woodlands and grasslands accurately: sparsely vegetated areas supported a larger pool of grass phytoliths while terrain covered by woody taxa yielded more woody phytoliths. In addition, soil phytolith assemblages recorded decadal ecological succession and changes in vegetation communities.

The comparison of soil phytoliths with local plants suggested that differential production, accumulation, and preservation interplay at the soil level. Well-known silica accumulators such as grasses seem localized and minor because of the cyclical, long-term succession to woodland. Although grasses boomed and busted through the waxing and waning of anthropogenic disturbance by fire and grazing, the long-term accumulation of woody types counterbalanced Poaceae representation. Therefore, phytoliths from local soils are capable of identifying woodlands as the climactic land cover across the landscape and they have value as a long-term proxy to detect ecosystem change and regulation that become apparent over a period of decades. The areal catchment recorded in the soils of the eastern Serengeti Plains encompasses several hectares. Furthermore, a heterogeneous record calls for research design that is able to capture spatial variability, documented here through large-scale soil sampling. The identified phytolith analog will let future work along the East African Rift System frame the environmental context of human evolution whenever the phytolith assemblages from associated sediments and paleosols yield a cohort of co-dominant types featuring the phytolith classes from dicot woody tissue and grasses identified here as typical of *Acacia-Commiphora* mosaics.

Acknowledgements

We thank the Tanzania Commission for Science and Technology, the Tanzanian Ministry of Natural Resources and Tourism (Division of Antiquities), the University of Dar es Salaam (DVC: Academic, Research), UDSM's College of Humanities and Department of Archaeology and Heritage, as well as the Oldupai Palaeoanthropology and Palaeoecology Project. We are grateful to the Tropical Pesticides Research Institute in Arusha. This work could not have been accomplished without the local Masai communities that welcomed us to their homeland and shared with us their expertise. A preprint of this manuscript has been made available online with the Open Science Framework (doi 10.31219/osf.io/3ed7f). This work was supported by the Canadian Social Science and Humanities Research Council Partnership Grant 895-2016-1017. We thank two anonymous reviewers and Dr. L. Wallis for their input.

References

- Albert R.M., Ofer L., Estroff L., Weiner S., Tsatskin A., Ronen A., Lev-Yadun S. 1999. Mode of occupation of Tabun Cave, Mt Carmel, Israel during the Mousterian period: a study of the sediments and phytoliths. *Journal of Archaeological Science* 26, 1249-1260.
- Albert R.M., Weiner S. 2001. Study of opal phytoliths in prehistoric ash layers using a quantitative approach. In: Meunier J, Coline F. (Eds.), *Phytoliths: applications in earth sciences and human history*. Balkema: Lisse, pp. 251-266.
- Albert R.M., Marean C. 2012. The exploitation of plant resources by early Homo sapiens: the phytolith record from Pinnacle Point 13B Cave, South Africa. *Geoarchaeology* 27, 363-384.
- Albert R.M., Bamford M.K., Cabanes D. 2006. Taphonomy of phytoliths and macroplants in different soils from Olduvai Gorge (Tanzania) and the application to Plio-Pleistocene palaeoanthropological samples. *Quaternary International* 148, 78-94.
- Albert, R.M., Bamford, M.K., Cabanes D. 2009. Palaeoecological significance of palms at Olduvai Gorge, Tanzania, based on phytolith remains. *Quaternary International* 193, 41-48.

Albert R.M., Bamford M.K., Esteban I. 2015. Reconstruction of ancient palm vegetation landscapes using a phytolith approach. *Quaternary International* 369, 51-66.

Aleman J.C., Canal-Subitani S., Favier C., Bremond L. 2014. Influence of the local environment on lacustrine sedimentary phytolith records. *Palaeogeography, Palaeoclimatology, Palaeoecology* 414, 273-283.

Alexandre A., Meunier J.D., Lézine A.M., Vincens A., Schwartz D. 1997. Phytoliths: indicators of grassland dynamics during the late Holocene in intertropical Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 136, 213-229.

An X., Lu H., Chu G. 2015. Surface soil phytoliths as vegetation and altitude indicators: a study from the southern Himalaya. *Scientific reports* 5, 15523.

Anderson G.D., Talbot L.M. 1965. Soil factors affecting the distribution of the grassland types and their utilization by wild animals on the Serengeti Plains, Tanganyika. *The Journal of Ecology*, 33-56.

Andrews P., Bamford M. 2008. Past and present vegetation ecology of Laetoli, Tanzania. *Journal of Human Evolution* 54, 78-98.

Arráiz H., Barboni D., Ashley G.M., Mabulla A., Baquedano E., Dominguez-Rodrigo, M. 2017. The FLK Zinj paleolandscape: reconstruction of a 1.84 Ma wooded habitat in the FLK Zinj-AMK-PTK-DS archaeological complex, Middle Bed I (Olduvai Gorge, Tanzania). *Palaeogeography, Palaeoclimatology, Palaeoecology* 488, 9-20.

Ashley G.M., Barboni D., Dominguez-Rodrigo M., Bunn H.T., Mabulla A.Z.P., Diez-Martin F., Barba R., Baquedano E. 2010. A spring and wooded habitat at FLK Zinj and their relevance to origins of human behavior. *Quaternary Research* 74: 304-314.

Bamford M.K., Albert R.M., Cabanes D. 2006 Plio–Pleistocene macroplant fossil remains and phytoliths from Lowermost Bed II in the eastern palaeolake margin of Olduvai Gorge, Tanzania. *Quaternary International* 148, 95-112.

Barboni D. 2014. Vegetation of Northern Tanzania during the Plio-Pleistocene: a synthesis of the paleobotanical evidences from Laetoli, Olduvai, and Peninj hominin sites. *Quaternary International* 322, 264-276.

Barboni D., Bremond L. 2009. Phytoliths of East African grasses: an assessment of their environmental and taxonomic significance based on floristic data. *Review of Palaeobotany and Palynology* 158, 29-41.

Barboni D., Ashley G.M., Dominguez-Rodrigo M., Bunn H.T., Mabulla A.Z., Baquedano E. 2010. Phytoliths infer locally dense and heterogeneous paleovegetation at FLK North and surrounding localities during upper Bed I time, Olduvai Gorge, Tanzania. *Quaternary Research* 74, 344-354.

Barboni D., Bonnefille R., Alexandre A., Meunier J.D. 1999. Phytoliths as palaeoenvironmental indicators, West Side Middle Awash Valley, Ethiopia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 152, 87-100.

Barboni D., Bremond L., Bonnefille R. 2007. Comparative study of modern phytolith assemblages from inter-tropical Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 246, 454-470.

Beyene Y., Katoh S., WoldeGabriel G., Hart W.K., Uto K., Sudo M., Kondo M., Hyodo M., Renne P.R., Suwa G., Asfaw B. 2013. The characteristics and chronology of the earliest Acheulean at Konso, Ethiopia. *Proceedings for the National Academy of Sciences* 110(5), 1584-1591.

Bibi F., Kiessling W. 2015. Continuous evolutionary change in Plio-Pleistocene mammals of eastern Africa. *Proceedings of the National Academy of Sciences* 112, 10623-10628.

Bibi F., Pante M., Souron A., Stewart K., Varela S., Werdelin L., Boissarie J.R., Fortelius M., Hlusko L., Njau J., de la Torre I. 2018. Paleoecology of the Serengeti during the Oldowan-Acheulean transition at Olduvai Gorge, Tanzania: The mammal and fish evidence. *Journal of Human Evolution* 120, 48-75.

Biswas O., Ghosh R., Paruya D.K., Mukherjee B., Thapa K.K., Bera S. 2016. Can grass phytoliths and indices be relied on during vegetation and climate interpretations in the eastern Himalayas? Studies from Darjeeling and Arunachal Pradesh, India. *Quaternary Science Reviews* 134, 114-132.

Blecker S.W., McCulley R.L., Chadwick O.A., Kelly E.F. 2006. Biologic cycling of silica across a grassland bioclimate sequence. *Global Biogeochemical Cycles* 20.

Blinnikov M.S. 1994. Phytolith analysis and Holocene dynamics of alpine vegetation. Veröffentlichungen des Geobotanischen Institutes der ETH. Zürich, *Stiftung Rübél*. 115, 23-40.

Blinnikov M.S. 2005. Phytoliths in plants and soils of the interior Pacific Northwest, USA. *Review of Palaeobotany and Palynology* 135, 71-98.

Blinnikov M.S., Bagent C.M., Reyerson P.E. 2013. Phytolith assemblages and opal concentrations from modern soils differentiate temperate grasslands of controlled composition on experimental plots at Cedar Creek, Minnesota. *Quaternary International* 287, 101-113.

Blinnikov M., Busacca A., Whitlock C. 2002. Reconstruction of the late Pleistocene grassland of the Columbia basin, Washington, USA, based on phytolith records in loess. *Palaeogeography, Palaeoclimatology, Palaeoecology* 177, 77-101.

Blumenthal S.A., Levin N.E., Brown F.H., Brugal J.P., Chritz K.L., Harris J.M., Jehle G.E., Cerling T.E. 2017. Aridity and hominin environments. *Proceedings of the National Academy of Sciences* 114, 7331-7336.

Bobe R., Behrensmeyer A.K. 2004. The expansion of grassland ecosystems in Africa in relation to mammalian evolution and the origin of the genus Homo. *Palaeogeography, Palaeoclimatology, Palaeoecology* 207, 399-420.

Bonnefille R., Lobreau D., Rioulet G. 1982. Fossil pollen of *Ximenia* (Olacaceae) in the Lower Pleistocene of Olduvai, Tanzania: palaeoecological implications. *Journal of Biogeography*, 169-186.

Bonnefille R. 1984. Palynological Research at Olduvai Gorge. *National Geographic Society Research Reports* 17, 227-242.

Bonnefille B. 1995. A reassessment of the Plio-Pleistocene pollen record of East Africa. In: E. Vrba, G. Denton, T. Partridge and L. Burckle (Eds.) *Paleoclimate and Evolution, with emphasis on human origins*. Yale University Press, New Haven. pp 299-310.

Bozarth S. 1992. Classification of Opal Phytoliths Formed in Selected Dicotyledons Native to the Great Plains. In: Rapp G.J., Mulholland S.C. (Eds.), *Phytolithic Systematics: Emerging Issues*. Plenum Press: New York, pp. 193-214.

Bremond L., Alexandre A., Hély C., Guiot, J. 2005. A phytolith index as a proxy of tree cover density in tropical areas: calibration with Leaf Area Index along a forest–savanna transect in southeastern Cameroon. *Global and Planetary Change* 45, 277-293.

Bremond L., Alexandre A., Wooller M.J., Hély C., Williamson D., Schäfer P.A., Majule A., Guiot J. 2008. Phytolith indices as proxies of grass subfamilies on East African tropical mountains. *Global and Planetary Change* 61, 209-224.

Cabanes D., Shahack-Gross R. 2015. Understanding fossil phytolith preservation: the role of partial dissolution in paleoecology and archaeology. *PloS One* 10, e0125532.

Cabanes D., Weiner S., Shahack-Gross R. 2011. Stability of phytoliths in the archaeological record: a dissolution study of modern and fossil phytoliths. *Journal of Archaeological Science* 38, 2480-2490.

Cahen L., Snehing N.J., Delhal J., Vail J.R., Bonhomme M., Ledent D. 1984. *The Geochronology and Evolution of Africa*. Clarendon Press, Oxford.

Carnelli A.L., Theurillat J.P., Madella M. 2004. Phytolith types and type-frequencies in subalpine–alpine plant species of the European Alps. *Review of palaeobotany and palynology* 129, 39-65.

Cerling T.E., Bowman J.R., O’Neil J.R. 1988. An isotopic study of a fluvial-lacustrine sequence: the Plio-Pleistocene Koobi For a sequence, East Africa. *Palaeogeography, palaeoclimatology, palaeoecology* 63, 335-356.

Cerling T.E., Wynn J.G., Andanje S.A., Bird M.I., Korir D.K., Levin N.E., Macharia A.N., Quade J., Remien, C.H. 2011. Woody cover and hominin environments in the past 6 million years. *Nature* 476, 51.

Clarke J. 2003. The occurrence and significance of biogenic opal in the regolith. *Earth-Science Reviews* 60, 175-194.

Collura L.V., Neumann K. 2017. Wood and bark phytoliths of West African woody plants. *Quaternary International* 43, 142-159.

- Copeland S.R. 2007. Vegetation and plant food reconstruction of lowermost Bed II, Olduvai Gorge, using modern analogues. *Journal of Human Evolution* 53(2), 146-175.
- Dawson J.B. 1992. Neogene tectonics and volcanicity in the North Tanzania sector of the Gregory Rift Valley: contrasts with the Kenya sector. *Tectonophysics* 204, 81-92.
- Deino A.L. 2012. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of Bed I, Olduvai Gorge, Tanzania, and the chronology of early Pleistocene climate change. *Journal of Human Evolution* 63, 251-273.
- EAMD East African Meteorological Department. 1967. The weather of East Africa. National Government Publication, Nairobi.
- Eichhorn B., Neumann K., Garnier A. 2010. Seed phytoliths in West African Commelinaceae and their potential for palaeoecological studies. *Palaeogeography, Palaeoclimatology, Palaeoecology* 298, 300-310.
- Esteban I., De Vynck J.C., Singels E., Vlok J., Marean C.W., Cowling R.M., Fisher E.C., Cabanes D., Albert R.M. 2017. Modern soil phytolith assemblages used as proxies for Paleoscape reconstruction on the south coast of South Africa. *Quaternary International* 434, 160-179.
- ESRI. 2016. ARCGIS ArcMap 10.5.1 [Computer software]. Redlands, CA, Esri Inc.
- Fahmy A.G. 2008. Diversity of lobate phytoliths in grass leaves from the Sahel region, West Tropical Africa: Tribe Paniceae. *Plant Systematics and Evolution* 270, 1-23.
- Fick S.E., Evett R.R. 2018. Distribution modelling of pre-Columbian California grasslands with soil phytoliths: New insights for prehistoric grassland ecology and restoration. *PloS One* 13, e0194315.
- Fishkis O., Ingwersen J., Streck T. 2009. Phytolith transport in sandy sediment: experiments and modeling. *Geoderma* 151, 168-178.
- Foster A., Ebinger C., Mbede E., Rex D. 1997. Tectonic development of the northern Taitaviiian sector of the East African Rift System. *Journal of the Geological Society* 154, 689-700.
- Frayse F., Pokrovsky O.S., Schott J., Meunier J.D. 2006. Surface properties, solubility and dissolution kinetics of bamboo phytoliths. *Geochimica et Cosmochimica Acta* 70, 1939-1951.
- Fredlund G.G., Tieszen L.T. 1994. Modern phytolith assemblages from the North American great plains. *Journal of Biogeography*, 321-335.
- Fredlund G.G., Egan D., Howell E.A. 2001. Inferring vegetation history from phytoliths. *The Historical Ecology Handbook*. Island Press, Washington DC.
- Frost P. 1996. The ecology of miombo woodlands. *The miombo in transition: woodlands and welfare in Africa*, 11-57.
- Gallego L., Distel R.A. 2004. Phytolith assemblages in grasses native to Central Argentina. *Annals of Botany* 94, 865-874.

- Gao G., Jie D., Liu L., Liu H., Li D., Li N., Shi J., Leng C., Qiao Z. 2018. Assessment and calibration of representational bias in soil phytolith assemblages in Northeast China and its implications for paleovegetation reconstruction. *Quaternary Research* 90, 38-49.
- Gao G., Jie D., Wang Y., Liu L., Liu H., Li D., Shi J., Leng C. 2018. Phytolith reference study for identifying vegetation changes in the forest– grassland region of northeast China. *Boreas* 47, 481-497.
- Hay R.L. 1976. *Geology of Olduvai Gorge. A study of sedimentation in a semiarid basin*. University of California Press, Los Angeles.
- Herlocker D.J., Dirchl H.J. 1972. Vegetation of Ngorongoro Conservation area, Tanzania. Ottawa: Canadian Wildlife Service (Canadian Wildlife Service Report Series no. 19). Illustrations, maps Geog, 5, pp. 39.
- Hodson M.J. 2016. The development of phytoliths in plants and its influence on their chemistry and isotopic composition. Implications for palaeoecology and archaeology. *Journal of Archaeological Science* 68, 62-69.
- Hodson M.J., White P.J., Mead A., Broadley M.R. 2005. Phylogenetic variation in the silicon composition of plants. *Annals of botany* 96, 1027-1046.
- Holmes A. 1951. The sequence of Precambrian orogenic belts in south and central Africa. In: Sandiford, K.S., Blondel, F. (Eds). *Proceedings of the 18th International Geological Congress, Association of African Geological Surveys*, London. pp. 254–269.
- Honaine M.F., Zucol A.F., Osterrieth M.L. 2009. Phytolith analysis of Cyperaceae from the Pampean region, Argentina. *Australian Journal of Botany* 57, 512-523.
- Hyland E., Smith S.Y., Sheldon N.D. 2013. Representational bias in phytoliths from modern soils of central North America: implications for paleovegetation reconstructions. *Palaeogeography Palaeoclimatology Palaeoecology* 374, 338–348.
- Jager T. 1982. *Soils of the Serengeti woodlands, Tanzania*. Wageningen: Pudoc.
- Kerns B.K., Moore M.M., Hart S.C. 2001. Estimating forest-grassland dynamics using soil phytolith assemblages and $\delta^{13}\text{C}$ of soil organic matter. *Ecoscience* 8, 478-488.
- Kindt R., Van Breugel P., Lillesø J.B., Bingham M., Demissew S., Dudley C., Friis I., Gachathi F., Kalema J., Mbago F., Minani V., Moshi H., Mulumba J., Namaganda M., Ndangalasi H., Ruffo C., Jamnadass R., Graudal L.O.V. 2011. *Potential Natural Vegetation of Eastern Africa (Ethiopia, Kenya, Malawi, Rwanda, Tanzania, Uganda and Zambia)*. Forest & Landscape, University of Copenhagen.
- Klopper R.R., Chatelain C., Bänninger V., Steyn H.M., De Wet B.C., Arnold T.H., Gautier L., Smith G.F., Spichiger R. 2006. Checklist of the flowering plants of Sub-Saharan Africa. An index of accepted names and synonyms. *South African Biodiversity Network Report* 42, SABONET, Pretoria.

Leakey M.D. 1971. *Olduvai Gorge. Excavations in Bed I and II, 1960–63, Vol. 3*. Cambridge University Press, Cambridge.

Leakey M.D. 1979. *Olduvai Gorge: My Search for Early Man*. HarperCollins.

Leakey M.D., Hay R.L., Thurber D.L., Protsch R., Berger R. 1972. Stratigraphy, archaeology, and age of the Ndutu and Naisiusiu beds, Olduvai Gorge, Tanzania. *World Archaeology* 3, 328-341.

Leakey M.D., Roe D. 1994. *Olduvai Gorge: Volume 5, Excavations in Beds III, IV and the Masek Beds (Vol. 5)*. Cambridge University Press, Cambridge.

Lepre C.J., Roche H., Kent D.V., Harmand S., Quinn R.L., Brugal J.P., Texier P.J., enoble A., Feibel C.S. 2011. An earlier origin for the Acheulean. *Nature* 477(7362), 82.

Levin N.E., Brown F.H., Behrensmeyer A.K., Bobe R., Cerling T. 2011. Paleosol carbonates from the Omo Group: Isotopic records of local and regional environmental change in East Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 307, 75-89.

Levin N.E., Quade J., Simpson S., Semaw S., Rogers M. 2004. Isotopic Evidence for Plio-Pleistocene Environmental Change at Gona, Ethiopia. *Earth and Planetary Science Letters* 219, 93-110.

Lovett J.C., Wasser S.K. 1993. *Biogeography and ecology of the rain forests of eastern Africa*. Cambridge University Press, Cambridge.

Lupien R.L., Russell J.M., Feibel C., Beck C., Castañeda I., Deino A., Cohen A.S. 2018. A leaf wax biomarker record of early Pleistocene hydroclimate from West Turkana, Kenya. *Quaternary Science Reviews* 186, 225-235.

Madella M., Alexandre A., Ball T. 2005. International code for phytolith nomenclature 1.0. *Annals of botany* 96, 253-260.

Madella M., Lancelotti C. 2012. Taphonomy and phytoliths: a user manual. *Quaternary International* 275, 76-83.

Magill C.R., Ashley G.M., Freeman K.H. 2013a. Ecosystem variability and early human habitats in eastern Africa. *Proceedings of the National Academy of Sciences* 110, 1167-1174.

Magill C.R., Ashley G.M., Freeman K.H. 2013b. Water, plants, and early human habitats in eastern Africa. *Proceedings of the National Academy of Sciences* 110, 1175-1180.

Mercader J., Bennett T., Esselmont C., Simpson S., Walde D. 2009. Phytoliths in woody plants from the Miombo woodlands of Mozambique. *Annals of Botany* 104, 91-113.

- Mercader J., Astudillo F., Barkworth M., Bennett T., Esselmont C., Kinyanjui R., Grossman D.L., Simpson S., Walde D. 2010. Poaceae phytoliths from the Niassa Rift, Mozambique. *Journal of Archaeological Science* 37, 1953-1967.
- Mercader J., Bennett T., Esselmont C., Simpson S., Walde D. 2011. Soil phytoliths from miombo woodlands in Mozambique. *Quaternary Research* 75, 138-150.
- Mercader J., Abtosway M., Baquedano E., Brid R.W., Diez-Martin F., Dominquez-Rodrigo M., Favreau J., Itambu M., Lee P., Mabulla A., Patalano R. 2017. Starch contamination landscapes in field archaeology: Olduvai Gorge, Tanzania. *Boreas* 46(4), 918-934.
- Mollet G.F., Swisher III C.C. 2012. The Ngorongoro Volcanic Highland and its relationships to volcanic deposits at Olduvai Gorge and East African Rift volcanism. *Journal of human evolution* 63, 274-283.
- Morris L.R., Ryel R.J., West N.E. 2010. Can soil phytolith analysis and charcoal be used as indicators of historic fire in the pinyon-juniper and sagebrush steppe ecosystem types of the Great Basin Desert, USA? *The Holocene* 20, 105-114.
- Murungi, M. 2017: Phytoliths at Sibudu (South Africa): Implications for vegetation, climate, and human occupation during the MSA. (PhD thesis). University of the Witwatersrand, Johannesburg.
- Neumann K., Fahmy A.G., Müller-Scheeßel N., Schmidt M. 2017. Taxonomic, ecological and palaeoecological significance of leaf phytoliths in West African grasses. *Quaternary International* 434, 15-32.
- Niboye, E.P. 2010. Vegetation Cover Changes in Ngorongoro Conservation Area from 1975 to 2000: The importance of remote sensing images. *The open Geography Journal* 3, 15-27.
- Novello A., Bamford M.K., Van Wijk Y., Wurz S. 2018. Phytoliths in modern plants and soils from Klasies River, Cape Region (South Africa). *Quaternary International* 464, 440-459.
- Novello A., Barboni D., Berti-Equille L., Mazur J.C., Poilecot P., Vignaud, P. 2012. Phytolith signal of aquatic plants and soils in Chad, Central Africa. *Review of Palaeobotany and Palynology* 178, 43-58.
- Norton-Griffiths M., Herlocker D., Pennycuik L. 1975. The patterns of rainfall in the Serengeti ecosystem, Tanzania. *African Journal of Ecology* 13, 347-374.
- Peters C.R., Blumenschine R.J. 1995. Landscape perspectives on possible land use patterns for Early Pleistocene hominids in the Olduvai Basin, Tanzania. *Journal of Human Evolution* 29(4), 321-362.
- Osterrieth M., Madella M., Zurro D., Alvarez M.F. 2009. Taphonomical aspects of silica phytoliths in the loess sediments of the Argentinean Pampas. *Quaternary International* 193, 70-79.
- Piperno, D. (1988). *Phytolith Analysis. An Archaeological and Geological Perspective*. London, Academic Press.

Piperno D.R. 2006. *Phytoliths: a comprehensive guide for archaeologists and paleoecologists*. Rowman Altamira.

Piperno D.R., Becker P. 1996. Vegetational history of a site in the central Amazon basin derived from phytolith and charcoal records from natural soils. *Quaternary research* 45, 202-209.

Plummer T.W., Ditchfield P.W., Bishop L.C., Kingston J.D., Ferraro J.V., Braun D.R., Hertel F., Potts R. 2009. Oldest evidence of toolmaking hominins in a grassland-dominated ecosystem. *PLoS One* 4, e7199.

Prassack K.A., Pante M.C., Njau J.K., de la Torre I. 2018. The paleoecology of Pleistocene birds from Middle Bed II, at Olduvai Gorge, Tanzania, and the environmental context of the Oldowan-Acheulean transition. *Journal of Human Evolution* 120, 32-47.

Quinn R.L., Lepre C.J., Wright J.D., Feibel, C. 2007. Paleogeographic variations of pedogenic carbonate $\delta^{13}\text{C}$ values from Koobi Fora, Kenya: implications for floral compositions of Plio-Pleistocene hominin environments. *Journal of Human Evolution* 53, 560-573.

Ravelo A.C., Andreasen D.H., Lyle M., Lyle A.O., Wara M.W. 2004. Regional climate shifts caused by gradual global cooling in the Pliocene epoch. *Nature* 429(6989), 263.

Rovner, I. 1971. Potential of opal phytoliths for use in paleoecological reconstruction. *Quaternary Research* 1: 343-359.

Runge F. 1999. The opal phytolith inventory of soils in central Africa—quantities, shapes, classification, and spectra. *Review of Palaeobotany and Palynology* 107, 23-53.

Sinclair AR, Mduma SA, Hopcraft JG, Fryxell JM, Hilborn R, Thirgood S. Long-term ecosystem dynamics in the Serengeti: lessons for conservation *Conserv Biol.* 2007 21(3):580-90.

Sinclair A.R.E., Packer C., Mduma S.A., Fryxell J.M. 2008. *Serengeti III: human impacts on ecosystem dynamics*. University of Chicago Press: Chicago.

Strömberg C.A. 2004. Using phytolith assemblages to reconstruct the origin and spread of grass-dominated habitats in the Great Plains of North America during the late Eocene to early Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 207, 239-275.

Strömberg C.A., Di Stilio V.S., Song Z. 2016. Functions of phytoliths in vascular plants: an evolutionary perspective. *Functional Ecology* 30, 1286-1297.

Strömberg C.A., Dunn R.E., Crifo C. Harris E.B. 2018. Phytoliths in paleoecology: analytical considerations, current use, and future directions. In: D.A. Crift, D.F. Su, S.W. Simpson (Eds.) *Methods in Paleoecology: Reconstructing Cenozoic Terrestrial Environments and Ecological Communities*. Springer, Switzerland. pp. 235-287.

- Sperazza M., Moore J.N., Hendrix M.S. 2004. High-resolution particle size analysis of naturally occurring very fine-grained sediment through laser diffractometry *Journal of Sedimentary Research* 74, 736-743.
- Thorn V.C. 2001. Oligocene and Early Miocene phytoliths from CRP-2/2A and CRP-3, Victoria Land Basin, Antarctica. *Terra Antarctica*. 8(4), 407-422.
- Thorn V.C. 2006. Vegetation reconstruction from soil phytoliths, Tongariro National Park, New Zealand. *New Zealand Journal of Botany* 44, 397-413.
- Tsartsidou G., Lev-Yadun S., Albert R.M., Miller-Rosen A., Efstratiou N., Weiner S. 2007. The phytolith archaeological record: strengths and weaknesses evaluated based on a quantitative modern reference collection from Greece. *Journal of Archaeological Science* 34, 1262-1275.
- Twiss P.C., Suess E., Smith R.M. 1969. Morphological Classification of Grass Phytoliths 1. *Soil Science Society of America Journal* 33, 109-115.
- Uno K.T., Polissar P.J., Jackson K.E. 2016. Neogene biomarker record of vegetation change in eastern Africa. *Proceedings of the National Academy of Sciences*, 201521267.
- Wallis L. 2003. An overview of leaf phytolith production patterns in selected northwest Australian flora. *Review of palaeobotany and palynology* 125, 201-248.
- Watling J., Iriarte J., Whitney B.S., Consuelo E., Mayle F., Castro W., Schaan D., Feldpausch T.R. 2016. Differentiation of neotropical ecosystems by modern soil phytolith assemblages and its implications for palaeoenvironmental and archaeological reconstructions II: Southwestern Amazonian forests. *Review of Palaeobotany and Palynology* 226, 30-43.
- White F. 1983. The Vegetation of Africa: A descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa. Natural Resources Research. UNESCO, Paris.

Table 1: Particle size analysis of selected topsoils, textural classification, phytolith-bearing fraction, modality, pH, color, and water content. See Figure 2 for sample location.

Table 2: Topsoil phytolith counts per group, class, and morphotype. See full dataset online (doi 10.20383/101.0122)

Table 3: Woody versus grassy phytoliths per plant cover rank, and short cell breakdown in topsoils.

Table 4: Percent of short cell morphotypes per plant cover rank in topsoils.

Table 5: Plant phytolith counts per group, class, and morphotype.

Table 6: Count of phytoliths by species and major group in descending order. See full dataset online for additional details (doi 10.20383/101.0122).

Table 7: Plant species studied for phytolith reference, with polymorphism per taxon. Figure 1 shows location of these species. Other details in full dataset online (doi 10.20383/101.0122).

Table 8: Topsoil and plant phytolith counts per group and class within the 6.25 ha quadrat.

Figure 1: a) Location map of Oldupai Gorge at the continental and regional levels; b) Plant sampling localities within the Oldupai Gorge region and location of 100 ha transect; c) 100 ha transect with the topsoil sampling localities and location of 6.25 ha quadrat; d) 6.25 ha plant sampling localities. Plant sampling legend: a) *Cynodon dactylon* b) *Sporobolus panicoides* c) *Asparagus africanus* d) *Ocimum* spp. e) *Acacia tortilis* f-h) *Sansevieria robusta* i) *Aristida adoensis* j) *Commiphora africana* k) *Commiphora kua* l) *Ximenia caffra* m) *Acacia mellifera* n) *Barleria eranthemoides* o) *Salvadora persica* p) *Aloe secundiflora* q) *Sarcostemma viminale* r) *Cissus cactiformis* s) *Hypoestes forskoolii* t-u) *Acacia tortilis* v) *Balanites aegyptiaca* w) *Commiphora* spp. x) *Acacia tortilis* y) *Sporobolus consimilis* z) *Pennisetum mezianum* aa) *Boscia angustiflora* ab) *Ximenia caffra* ac) *Commiphora merkeri* ad) *Acacia mellifera* ae) *Cissus quadrangularis* af) *Aloe secundiflora* ag) *Asparagus africanus* ah) *Balanites aegyptiaca* ai) *Acacia mellifera* aj) *Ximenia caffra* ak) *Maerua triphylla* al) *Acacia nilotica*.

Figure 2: Sampling localities for soils and particle size analysis in relation to plant cover rank, with catchment transect in outline. See Table 1 for analytical description of soils.

Figure 3: Particle size distribution examples a) bimodal (Sample 192), and b) unimodal (Sample 229). Refer to Table 1 for complete sample characterization, and Supplementary Materials Figure 1 for sampling location.

Figure 4: Soil phytoliths and diatoms a) bilobate, short concave (#30); b) bilobate, short notched (#166); c) bilobate short convex (#54); d) bilobate, short flat (#112); e) bilobate, long concave (#145); f) bilobate, long convex (#98); g) rondel (#30); h) tower (#112); i) tower, horned (#112); j) tower, tapered (#65); k) lanceolate (#30); l) bulliform (#54); m) saddle, squat (#30); n) saddle (#65); o) sclereid (#191); p) clavate (#65); q) tabular thick lacunate (#54); r) tabular strangulated (#127); s) tabular sinuate (#120); t) tabular scrobiculate (#65); u) tabular sulcate (#30); v) blocky (#30); w) blocky thick lacunate (#2); x) cylindroid sinuate (#65); y) cylindroid scrobiculate (#149); z) cylindroid thick lacunate (#30); aa) globular granulate (#191); ab) guttiform (#30); ab) fusiform (#54); ad) globular facetate (#30); ae) polygonal prism (#131); af-ah) diatoms (#65, 112, 258).

Figure 5: Highest ranking woody phytolith proportions in relation to plant cover rank. Please see Figs. 1 and 2 for location of sampling area and corresponding sample numbers.

Figure 6: Highest ranking short cell phytolith proportions in relation to plant cover rank. Please see Figs. 1 and 2 for location of sampling area and corresponding sample numbers.

Figure 7: Plant phytoliths a) bilobate, short concave (left: *P. mezianum* right: *A. adoensis*); b) bilobate, short convex (*A. adoensis*); c) bilobate, long convex (*A. adoensis*); d) rondel, ovate (*S. panicoides*); e) rondel, horned (*S. spp.*); f) tower (*S. consimilis*); g) saddle, short (*C. dactylon*); h) bulliform (*S. consimilis*); i) globulose (*A. secundiflora*); j) globular granulate (left: *B. eranthemoides* right: *A. secundiflora*); k) oblong granulate (left: *B. angustifolia*; right: *H. forskoolii*); l) blocky (*Ocimum* spp.); m) blocky (*A. tortilis*); n) blocky radiating (*X. caffra*); o) fusiform (*O. spp.*); p) guttiform (*C. cactiformis*); q) tabular sinuate (*A. mellifera*); r) tabular psilate (*A. mellifera*); s) tabular (*A. tortilis*); t) tabular scrobiculate (*M. tryphilla*); u) tabular strangulated (*S. persica*); v) tabular crenate (*S. consimilis*); w) cylinder sinuate (*A. nilotica*); x) cylinder

scrobiculate (*C. cactiformis*); y) sclereid (*A. tortilis*); z) hair (*C. dactylon*); aa) hair base (*X. caffra*)
ab) epidermal cells (*B. eranthemoides*); ac) globular mesophyll (*X. caffra*); ad) vessel (*A. tortilis*).

Figure 1

Figure 2

Particle Size Distribution

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Table 1

Sample	Mean particle size, μm	Particle class	Silt & Very Fine Sand, %	Modality	Sediment pH	Sediment Colour	Sediment H ₂ O content, %
43	294.2	Medium sand	39.62	Unimodal	8.7	Dark yellowish brown	3.9
63	513.8	Coarse sand	22.73	Bimodal	8.3	Moderate yellowish brown	5.4
85	216.9	Fine sand	49.49	Unimodal	8.8	Dark yellowish brown	6.5
138	94.7	Very fine sand	72.86	Unimodal	8.7	Dark yellowish brown	3.6
146	335.8	Medium sand	38.33	Unimodal	9.1	Pale yellowish brown	3.7
192	469.6	Medium sand	37.59	Bimodal	8.6	Moderate yellowish brown	2.7
229	322.1	Medium sand	26.31	Unimodal	8.7	Dark yellowish brown	2.8
236	670.3	Coarse sand	17.75	Bimodal	8.4	Dark yellowish brown	3.7
279	736.8	Coarse sand	6.54	Unimodal	9.2	Dark yellowish brown	2.9
286	300.1	Medium sand	35.52	Bimodal	8.6	Dark yellowish brown	4.0
Mean	344.3	Medium sand	29.55	Unimodal	8.7	Yellowish brown	3.8

Table 2

Major Group	Phytolith Class	Phytolith Morphotype	Count
Poaceae short cell	Lobate	Bilobate, long concave	38
		Bilobate, long convex	8
		Bilobate, long flat	14
		Bilobate, long notched	4
		Bilobate, short concave	428
		Bilobate, short convex	58
		Bilobate, short flat	114
		Bilobate, short notched	28
		Cross	1
		Polylobate	2
	Rondel	Rondel, horned	8
		Rondel, truncated	12
		Rondel, tower	475
		Rondel	255
	Saddle	Saddle, ovate	24
		Saddle, short	469
		Saddle, squat	140
Woody	Blocky	Blocky	879
		Blocky corniculate	2
		Blocky ridged	405
		Blocky scrobiculate	23
		Blocky sinuate	28
		Blocky thick lacunate	72
	Cylindrical	Cylinder corniculate	3
		Cylinder psilate	132
		Cylinder scrobiculate	569
		Cylinder sinuate	115
		Cylinder thick lacunate	61
	Fusiform	Fusiform	11
		Guttiform	68
	Sclereid/Clavate	Clavate	27
		Sclereid	119
	Spherical	Globular bisected	1
		Globular facetate	778
		Globular granulate	83
		Globular granulate, large	3
		Globular granulate oblong	24
		Globular psilate	10
		Globulose	2
		Globulose segmented	1
		Oblong granulate	10
		Hemisphere psilate	3
	Tabular	Tabular corniculate	11
		Tabular ellipsoidal	35
		Tabular ellipsoidal, large	21

		Tabular facetate	8
		Tabular laminate	186
		Tabular oblong	16
		Tabular pilate	3
		Tabular scrobiculate	2036
		Tabular sinuate	918
		Tabular strangulated	11
		Tabular sulcate	509
		Tabular thick lacunate	731
Undetermined grass	Grass, undetermined	Bulliform	47
		Trichome	264
Rare/Unknown	Composite	Epidermal	3
		Vessel	18
	Drum	Polygonal prism	2
	Sedge	Papillae	1
	Short cell, other	Trapeziform sinuate	1
	Tabular undetermined	Tabular	70
		Tabular crenate	36
		Tabular psilate	311

Table 3

Plant Cover Rank	Woody	Grass	Short Cell Class			Grand Total
			Lobate	Rondel	Saddle	
0	74.8	25.2	34.4	36.3	29.3	100.0
1	72.7	27.3	29.0	39.3	31.7	100.0
2	79.3	20.7	35.8	33.3	30.9	100.0
3	80.8	19.2	37.3	33.3	29.4	100.0

Table 4

Morphotype	Plant Cover Rank				Grand Total
	0	1	2	3	
Most frequent short cell					
Rondel, tower	36.7	36.9	30.6	31.9	34.6
Saddle, short	33.5	34.1	35.8	32.6	34.2
Bilobate, short concave	29.9	29.0	33.6	35.4	31.2
Grand Total	100.0	100.0	100.0	100.0	100.0
Saddle					
Saddle, ovate	10.4	0.0	0.6	0.0	3.8
Saddle, short	70.7	71.8	80.2	79.7	74.3
Saddle, squat	18.9	28.2	19.6	21.7	22.1
Grand Total	100.0	100.0	100.0	100.0	100.0
Rondel					
Rondel, horned	1.9	0.9	0.6	0.0	1.1
Rondel, tower	62.5	62.6	63.4	68.7	63.3
Rondel, truncated	2.5	0.9	1.1	1.5	1.6
Rondel	33.1	35.6	34.9	29.9	34.0
Grand Total	100.0	100.0	100.0	100.0	100.0
Bilobate					
Bilobate, long concave	5.4	4.7	5.9	6.7	5.5
Bilobate, long convex	1.9	0.6	1.1	0.0	1.2
Bilobate, long flat	3.5	1.2	1.1	1.3	2.0
Bilobtae, long notched	1.5	0.0	0.0	0.0	0.6
Bilobate, short concave	53.8	66.9	64.9	68.0	61.6
Bilobate, short convex	8.1	7.6	9.0	9.3	8.3
Bilobate, short flat	21.5	16.3	11.2	12.0	16.4
Bilobate, short notched	3.8	1.7	6.9	2.7	4.0
Cross	0.0	0.6	0.0	0.0	0.1
Polylobate	0.4	0.6	0.0	0.0	0.3
Grand Total	100.0	100.0	100.0	100.0	100.0

Table 5

Major Group	Phytolith Class	Phytolith Morphotype	Count
Poaceae short cell	Lobate	Bilobate, long concave	26
		Bilobate, long convex	102
		Bilobate, short concave	491
		Bilobate, short convex	30
		Bilobate, short flat	19
		Polylobate	19
	Rondel	Rondel, horned	82
		Rondel, ovate	464
		Rondel, tower	86
		Rondel	11
	Saddle	Saddle, short	211
		Saddle, squat	15
Woody	Blocky	Blocky	79
		Blocky radiating	9
		Blocky scrobiculate	1
	Cylindrical	Cylinder crenate	8
		Cylinder psilate	35
		Cylinder scrobiculate	6
		Cylinder sinuate	46
	Fusiform	Fusiform	16
		Guttiform	23
	Sclereid/Clavate	Clavate	7
		Sclereid	11
	Spherical	Globular granulate	132
		Globular granulate, large	2
		Globular granulate oblong	11
		Globular psilate	1
		Globular tuberculate	5
		Globulose	239
		Hemisphere psilate	1
		Oblong granulate	596
	Tabular	Tabular ellipsoidal	46
		Tabular laminate	4
		Tabular oblong	9
		Tabular scrobiculate	113
		Tabular sinuate	74
		Tabular strangulated	2
		Tabular subrounded	18
		Tabular sulcate	3
		Tabular thick lacunate	12
Undetermined grass	Grass, undetermined	Bulliform	63
		Trichome	19
Rare/Unknown	Composite	Epidermal cells	131
		Vessel	224
	Epidermal complex	Globular mesophyll	127

	Hair	62
	Hair base	104
	Palisade	6
	Stomata	17
Tabular, undetermined		
	Tabular	279
	Tabular crenate	54
	Tabular psilate	159

Table 6

Species	Phytolith Major Group				Grand Total
	Poaceae short cell	Rare/ Unknown	Undetermined Grass	Woody	
<i>Boscia angustifolia</i>	0	2	0	537	539
<i>Pennisetum mezianum</i>	264	64	6	0	334
<i>Sporobolus africanus</i>	302	16	12	0	330
<i>Aristida adoensis</i>	186	78	15	1	280
<i>Sporobolus consimilis</i>	192	2	43	0	237
<i>Sporobolus</i> spp.	207	20	0	0	227
<i>Cynodon dactylon</i>	208	6	5	0	219
<i>Sporobolus panicoides</i>	197	19	1	0	217
<i>Ximenia caffra</i>	0	183	0	22	205
<i>Melhanian parviflora</i>	0	61	0	122	183
<i>Acacia tortilis</i>	0	94	0	75	169
<i>Maerua tryphilla</i>	0	66	0	102	168
<i>Acacia nilotica</i>	0	96	0	64	160
<i>Cissus cactiformis</i>	0	59	0	81	140
<i>Barleria eranthemoides</i>	0	85	0	45	130
<i>Ocimum</i> spp	0	33	0	89	122
<i>Commiphora</i> spp.	0	121	0	0	121
<i>Hypoestes forskalii</i>	0	2	0	107	109
<i>Aloe secundiflora</i>	0	3	0	97	100
<i>Asparagus africanus</i>	0	54	0	24	78
<i>Acacia mellifera</i>	0	12	0	45	57
<i>Balanites aegyptiaca</i>	0	23	0	34	57
<i>Commiphora africana</i>	0	4	0	25	29
<i>Cissus quadrangularis</i>	0	21	0	4	25
<i>Commiphora merkeri</i>	0	10	0	13	23
<i>Sansevieria robusta</i>	0	13	0	10	23
<i>Salvadora persica</i>	0	10	0	6	16
<i>Sarcostemma viminale</i>	0	6	0	6	12
Grand Total	1556	1163	82	1509	4310

Table 7

Clade (Poaceae)	Species	Morphotype, n
	<i>Aloe secundiflora</i>	8
	<i>Acacia mellifera</i>	9
	<i>Acacia nilotica</i>	11
	<i>Acacia tortilis</i>	20
Aristidoideae	<i>Aristida adoensis</i>	15
	<i>Asparagus africanus</i>	7
	<i>Balanites aegyptiaca</i>	15
	<i>Barleria eranthemoides</i>	15
	<i>Boscia angustiflora</i>	3
	<i>Cissus cactiformis</i>	14
	<i>Cissus quadrangularis</i>	6
	<i>Commiphora africana</i>	8
	<i>Commiphora merkeri</i>	6
	<i>Commiphora kua</i>	0
	<i>Commiphora</i> spp.	6
Chloridoideae	<i>Cynodon dactylon</i>	5
	<i>Hypoestes forskalii</i>	6
	<i>Maerua triphylla</i>	11
	<i>Melhania parviflora</i>	15
	<i>Ocimum</i> spp.	16
Panicoideae	<i>Pennisetum mezianum</i>	7
	<i>Salvadora persica</i>	6
	<i>Sansevieria robusta</i>	7
	<i>Sarcostemma viminale</i>	5
Chloridoideae	<i>Sporobolus africanus</i>	10
Chloridoideae	<i>Sporobolus consimilis</i>	8
Chloridoideae	<i>Sporobolus panicoides</i>	4
Chloridoideae	<i>Sporobolus</i> spp.	6
	<i>Ximenia caffra</i>	14

Table 8

Major Group	Phytolith Class	Plants n=18	Topsoil n=6
Poaceae short cell	Lobate	169	60
	Rondel	197	61
	Saddle	225	78
Woody	Blocky	69	290
	Cylindrical	65	80
	Fusiform	30	21
	Sclereid/Clavate	11	15
	Spherical	185	59
	Tabular	166	1087
Undetermined grass	Grass, undetermined	21	65
Rare/Unknown	Composite	195	2
	Epidermal complex	201	0
	Tabular, undetermined	263	94

Supplemental Figure 1: Extraction weight per sample in relation to plant cover rank and altitude.

Supplemental Figure 2: Highest ranking rondel phytolith proportions in relation to plant cover rank.

Supplemental Figure 3: Highest ranking bilobate phytolith proportions in relation to plant cover rank.

Supplemental Figure 4: Highest ranking saddle phytolith proportions in relation to plant cover rank.

Supplemental Figure 5: Topsoil samples with presence of blocky ridged morphotype.

Supplemental Figure 6: Topsoil samples with presence of blocky thick lacunate morphotype.

Supplemental Figure 7: Topsoil samples with presence of globular facetate morphotype.

Supplemental Table 1: Count of phytoliths by modern plant species and plant part.

Species	Plant Part				Woody Tissue	Grand Total
	Culm	Leaf	Thorn	Whole		
<i>Acacia mellifera</i>	0	0	0	0	57	57
<i>Acacia nilotica</i>	0	130	0	0	30	160
<i>Acacia tortilis</i>	0	107	0	0	62	169
<i>Aloe secundiflora</i>	0	93	0	0	7	100
<i>Aristida adoensis</i>	0	280	0	0	0	280
<i>Asparagus africanus</i>	0	40	0	0	38	78
<i>Balanites aegyptiaca</i>	0	21	0	0	36	57
<i>Barleria eranthemoides</i>	0	113	0	0	17	130
<i>Boscia angustifolia</i>	0	304	0	0	235	539
<i>Cissus cactiformis</i>	0	0	0	0	140	140
<i>Cissus quadrangularis</i>	0	0	0	0	25	25
<i>Commiphora africana</i>	0	0	0	0	29	29
<i>Commiphora merkeri</i>	0	0	23	0	0	23
<i>Commiphora spp</i>	0	121	0	0	0	121
<i>Cynodon dactylon</i>	0	219	0	0	0	219
<i>Hypoestes forskalii</i>	0	107	0	0	2	109
<i>Maerua tryphilla</i>	0	109	0	0	59	168
<i>Melhania parviflora</i>	0	44	0	0	139	183
<i>Ocimum spp.</i>	0	44	0	0	78	122
<i>Pennisetum mezianum</i>	50	284	0	0	0	334
<i>Salvadora persica</i>	0	2	0	0	14	16
<i>Sansevieria robusta</i>	0	0	0	0	23	23
<i>Sarcostemma viminale</i>	0	0	0	0	12	12
<i>Sporobolus africanus</i>	0	0	0	330	0	330
<i>Sporobolus consimilis</i>	0	237	0	0	0	237
<i>Sporobolus panicoides</i>	217	0	0	0	0	217
<i>Sporobolus spp.</i>	0	227	0	0	0	227
<i>Ximenia caffra</i>	0	194	0	0	11	205
Grand Total	267	2676	23	330	1014	4310

Supplemental Table 2: Topsoil sample ratio of Poaceae short cell versus woody morphotypes in descending order of value.

Sample	Poaceae	Woody	Ratio	Overall Sample Size
112	278	284	0.9789	600
98	197	347	0.5677	573
145	141	262	0.5382	430
65	132	270	0.4889	422
277	102	269	0.3792	388
30	201	596	0.3372	855
258	48	144	0.3333	218
183	70	211	0.3318	304
240	37	125	0.2960	171
176	19	66	0.2879	95
160	84	302	0.2781	415
153	128	473	0.2706	642
131	98	367	0.2670	498
271	28	109	0.2569	146
96	60	267	0.2247	351
292	36	171	0.2105	213
191	56	293	0.1911	356
200	52	301	0.1728	413
54	40	236	0.1695	293
149	18	113	0.1593	151
195	26	164	0.1585	204
172	17	110	0.1545	142
2	40	263	0.1521	350
166	17	120	0.1417	143
260	33	260	0.1269	315
121	14	115	0.1217	140
170	26	232	0.1121	277
66	8	98	0.0816	138
120	24	296	0.0811	339
227	18	246	0.0732	272
230	22	437	0.0503	491
53	4	82	0.0488	96
97	4	89	0.0449	103
245	0	14	0.0000	16
287	0	182	0.0000	185

Supplemental Table 3: Major group representation by sample type.

Major Group	Sample Origin		Grand Total
	Modern Plant	Soil	
Poaceae short cell	1556	2078	3634
Rare/Unknown	1163	442	1605
Undetermined Grass	82	311	393
Woody	1509	7914	9423
Grand Total	4310	10745	15055

Supplemental Table 4: Blocky, cylindrical, spherical, tabular morphotypes by sample origin.

Morphotype	Sample Origin		Grand Total
	Modern Plant	Soil	
Blocky			
Blocky	79	879	958
Blocky corniculate	0	2	2
Blocky radiating	9	0	9
Blocky ridged	0	405	405
Blocky scrobiculate	1	23	24
Blocky sinuate	0	28	28
Blocky thick lacunate	0	72	72
Grand Total	89	1409	1498
Cylindrical			
Cylinder corniculate	0	3	3
Cylinder crenate	8	0	8
Cylinder psilate	35	132	167
Cylinder scrobiculate	6	569	575
Cylinder sinuate	46	115	161
Cylinder thick lacunate	0	61	61
Grand Total	95	880	975
Spherical			
Globular bisected	0	1	1
Globular facetate	0	778	778
Globular granulate	132	83	215
Globular granulate large	2	3	5
Globular granulate oblong	11	24	35
Globular psilate	1	10	11
Globular tuberculate	5	0	5
Globulose	239	2	241
Globulose segmented	0	1	1
Hemisphere psilate	1	3	4
Oblong granulate	596	10	606
Grand Total	987	915	1902
Tabular			
Tabular corniculate	0	11	11
Tabular ellipsoidal	46	35	81
Tabular ellipsoidal large	0	21	21
Tabular facetate	0	8	8
Tabular laminate	4	186	190
Tabular oblong	9	16	25
Tabular pilate	0	3	3
Tabular scrobiculate	113	2036	2149

Tabular sinuate	74	918	992
Tabular strangulated	2	11	13
Tabular subrounded	18	0	18
Tabular sulcate	3	509	512
Tabular thick lacunate	12	731	743
Grand Total	281	4485	4766

Supplemental Table 5: Poaceae short cell group and class morphotypes by sample origin.

Morphotype	Sample Type		Grand Total
	Modern Plant	Soil	
Lobate			
Bilobate, long concave	26	38	64
Bilobate, long convex	102	8	110
Bilobate, long flat	0	14	14
Bilobate, long notched	0	4	4
Bilobate, short concave	491	428	919
Bilobate, short convex	30	58	88
Bilobate, short flat	19	114	133
Bilobate, short notched	0	28	28
Cross	0	1	1
Polylobate	19	2	21
Grand Total	687	695	1382
Rondel			
Rondel	11	255	266
Rondel, horned	82	8	90
Rondel, ovate	464	0	464
Rondel, tower	86	475	561
Rondel, truncated	0	12	12
Grand Total	643	750	1393
Saddle			
Saddle, ovate	0	24	24
Saddle, short	211	469	680
Saddle, squat	15	140	139
Grand Total	226	633	859
Poaceae short cell class			
Lobate	687	695	1382
Rondel	643	750	1393
Saddle	226	633	859
Grand Total	1556	2078	3634