Some Remarks and Propositions on Riemann Hypothesis

Jamal Salah

Department of Basic Science, College of Health and Applied Sciences

A' Sharqiyah University, Ibra post code 400 Oman

drjamals@asu.edu.om

Riemann Hypothesis

$$\zeta(s) = 0 \rightarrow \operatorname{Re}(s) = \frac{1}{2}$$

There are roughly 10^{13} nontrivial zeros lying on the critical line **$\operatorname{Re}(s) = 1/2$**

Abstract

In **1859**, **Bernhard Riemann**, a German mathematician, published a paper to the Berlin Academy that would change mathematics forever. The mystery of **prime numbers** was the focus. At the core of the presentation was indeed a concept that had not yet been proven by Riemann, one that to this day baffles mathematicians. The way we do business could have been changed **if** the Riemann hypothesis holds **true**, that is because prime numbers are the **key element** for **banking** and **e-commerce security**. It will also have a significant influence, impacting quantum mechanics, chaos theory, and the future of computation, on the cutting edge of science. In this article we look at some well know results of Riemann Zeta function in a different light. We explore the **proofs of Zeta integral Representation, Analytic continuity and the first functional equation**. Initially, we observe omitting a logical undefined term in the integral representation of Zeta function by the means of **Gamma function**. For that we propose **some modifications** in order to reasonably justify the location of the non-trivial zeros on the critical line, by assuming that $\text{Zeta}(s)$ and $\text{Zeta}(1-s)$ simultaneously **equal zero**. Consequently, we **conditionally** prove Riemann Hypothesis.

Euler's Approach

□ Zeta Function:

$$\zeta(x) = \sum_{n=1}^{\infty} n^{-x}, \quad x > 1$$

□ Euler's Observation:

$$\zeta(x) = \prod_{p \text{ Prime}} \frac{p^x}{p^x - 1}$$

□ Euler's Analytic Continuity:

$$(1 - 2^{1-x})\zeta(x) = \sum_{n=1}^{\infty} (-1)^{n+1} n^{-x}$$

$$\zeta(x) = \frac{1}{1 - 2^{1-x}} \sum_{n=1}^{\infty} (-1)^{n+1} n^{-x}, \quad 0 < x < 1$$

Euler's Approach

Euler's omitting restrictions to compute $\zeta(-1)$

□ Given:

$$\zeta(-1) = -\frac{1}{3} \sum_{n=1}^{\infty} (-1)^{n+1} n = -\frac{1}{3} (1 - 2 + 3 - 4 + \dots)$$

□ Derive and plug -1:

$$1 + x + x^2 + \dots + x^n + \dots = \frac{1}{1-x}, \quad |x| < 1$$

$$1 + 2x + 3x^2 + \dots = \frac{1}{(1-x)^2} \rightarrow 1 - 2 + 3 - 4 + \dots = \frac{1}{4}$$

$$\zeta(-1) = -\frac{1}{3} \times \frac{1}{4} = -\frac{1}{12}$$

Integral Representation: Riemann's Approach

Complex Zeta Function:

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s}, \quad \operatorname{Re}(s) > 1$$

Integral Representation (Multiplying Zeta by Gamma):

Gamma Function:

$$\Gamma(s) = \int_0^{\infty} e^{-\tau} \tau^{s-1} d\tau, \quad \operatorname{Re}(s) > 1$$

$$\tau = nt \rightarrow d\tau = n dt$$

$$\Gamma(s) = n^s \int_0^{\infty} e^{-nt} t^{s-1} dt$$

$$\tau = \infty \times 0 \rightarrow t \neq 0$$

$$\zeta(s)\Gamma(s) = \sum_{n=1}^{\infty} \int_0^{\infty} e^{-nt} t^{s-1} dt = \int_0^{\infty} \left(\sum_{n=1}^{\infty} e^{-nt} \right) t^{s-1} dt = \int_0^{\infty} \frac{t^{s-1}}{e^t - 1} dt, \quad \operatorname{Re}(s) > 1.$$

Modified Integral Representation

$$\zeta(s)\Gamma(s) = \int_0^{\infty} \frac{t^{s-1}}{e^t - 1} dt, \quad \operatorname{Re}(s) > 1 \text{ and } t \neq 0$$

□ Theorem 1:

$$\zeta(1-s)\Gamma(1-s) = \int_0^{\infty} \frac{t^{-s}}{e^t - 1} dt, \quad \operatorname{Re}(s) < 0, \quad t \neq 0$$

$$\zeta(s) = \zeta(1-s) = 0 \rightarrow \operatorname{Re}(s) = \frac{1}{2}$$

Proof of Theorem 1

□ Proof.

$$\zeta(s)\Gamma(s) + \zeta(1-s)\Gamma(1-s) = \int_0^{\infty} \frac{t^{s-1} + t^{-s}}{e^t - 1} dt, \quad t \neq 0$$

$$\int_0^{\infty} \frac{t^{s-1} + t^{-s}}{e^t - 1} dt = 0, \quad t \neq 0$$

$$t^{s-1} + t^{-s} = 0, \quad t \neq 0.$$

$$s = \frac{1}{2} + i \frac{(2m+1)\pi}{2\ln t}, \quad t \neq 0, \quad \ln t \neq 0 \text{ and } m \in \mathbb{Z}$$

Functional Equation (Riemann's Approach)

□ Hankel Contour:

$$t = \epsilon e^{i\theta}, 0 \leq \theta \leq 2\pi, \quad \text{then let } \epsilon \rightarrow 0$$

$$\oint_C \frac{(-t)^{s-1}}{e^t - 1} dt = \int_{\rho_1} \frac{(-t)^{s-1}}{e^t - 1} dt + \int_{\rho} \frac{(-t)^{s-1}}{e^t - 1} dt + \int_{\rho_2} \frac{(-t)^{s-1}}{e^t - 1} dt$$

0

$$\zeta(s) = \frac{\Gamma(1-s)}{2\pi i} \oint_C \frac{(-t)^{s-1}}{e^t - 1} dt$$

Functional Equation (Riemann's Approach)

□ The modified Contour:

$$\oint_{\gamma} \frac{(-t)^{s-1}}{e^t - 1} dt = \lim_{R \rightarrow \infty} \left(\int_{C_R} \frac{(-z)^{s-1}}{e^z - 1} dz + \int_{D_R} \frac{(-z)^{s-1}}{e^z - 1} dz \right) = -(2\pi i) \sum_{n \in \mathbb{Z}^+} \operatorname{Res}_{z=2\pi i n} \frac{(-z)^{s-1}}{e^z - 1}$$

$$\zeta(s) = \frac{\Gamma(1-s)}{2\pi i} \oint_{\gamma} \frac{(-t)^{s-1}}{e^t - 1} dt = \frac{\Gamma(1-s)}{2\pi i} \left(2i 2^s \pi^s \zeta(1-s) \sin\left(\frac{\pi}{2}s\right) \right)$$

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

Modifications: Functional Equation

□ Remark:

$$t = \epsilon e^{i\theta}, \quad 0 \leq \theta \leq 2\pi, \quad |t| = \epsilon \rightarrow \epsilon \neq 0$$

□ Modifications:

$$\zeta(s) = \frac{\Gamma(1-s)}{2\pi i} \oint_C \frac{(-t)^{s-1}}{e^t - 1} dt$$

$$\zeta(s) = \frac{\Gamma(1-s)}{2\pi i} \epsilon^{s-1} + \frac{\Gamma(1-s)}{2\pi i} \oint_C \frac{(-t)^{s-1}}{e^t - 1} dt$$

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

$$\zeta(s) = \epsilon^{s-1} + 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

Proof of Riemann Hypothesis

□ Proof.

$$\zeta(s) = \epsilon^{s-1} + 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

$$\zeta(s) - A(s)\zeta(1-s) + \zeta(1-s) - A(1-s)\zeta(s) = \epsilon^{s-1} + \epsilon^{-s}$$

$$\zeta(1-s) = \epsilon^{-s} + 2^{1-s} \pi^{-s} \sin\left(\frac{\pi(1-s)}{2}\right) \Gamma(s) \zeta(s)$$

$$s = \frac{1}{2} + i \frac{(2m+1)\pi}{2 \ln \epsilon}, \quad \epsilon \neq 0, \quad \ln \epsilon \neq 0 \text{ and } m \in \mathbb{Z}$$

References

- ❶ Jamal Salah, Some Remarks and Proposition on Riemann Hypothesis, Mathematics and Statistics, Vol. 9, No. 2, pp. 159 - 165, 2021. DOI: 10.13189/ms.2021.090210.
- ❷ Jamal Y. Mohammad Salah An Alternative perspective to Riemann Hypothesis, PSYCHOLOGY AND EDUCATION (2020) 57(9): 1278-1281.
- ❸ Jamal Y. Mohammad Salah, The consequence of the analytic continuity of Zeta function subject to an additional term and a justification of the location of the non-trivial zeros, International Journal of Science and Research (IJSR). Volume 9, Issue 3, Pages 1566 – 1569. 2020.
- ❹ Jamal Y. Mohammad Salah, Two Conditional proofs of Riemann Hypothesis, International Journal of Sciences: Basic and Applied Research (IJSBAR). Volume 49, Issue 1, Pages 74-83. 2020.