Differential expression of GATM in cancer of the vulva.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

In these brief notes we document work using published microarray data (1, 2) to pioneer integrative transcriptome analysis comparing vulvar carcinoma to its tissue of origin, the vulva. We report the differential expression of glycine amidinotransferase, encoded by GATM, in cancer of the vulva. GATM may be of pertinence to understanding transformation and disease progression in vulvar cancer (3).

Keywords: vulvar cancer, vulvar carcinoma, GATM, glycine amidinotransferase.

Gynecologic cancers include cervical cancer, ovarian cancer, endometrial (uterine) cancer, and vulvar cancer (4, 3). Vulvar cancer is thought to be the least prevalent of these, with 6190 cases reported in the United States in 2018 (5). We integrated separate published microarray datasets to understand, at the systems level, the transcriptional landscape of vulvar carcinoma.

Methods

We utilized microarray datasets GSE63678 (1) and GSE38228 (2) for this integrated, transcriptome-level differential gene expression analysis of human vulvar cancers in conjunction with GEO2R. GSE63678 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=7$ normal vulvar tissues and $n=6$ vulvar carcinoma tissues; analysis was performed using platform GPL571. GSE38228 was generated using Illumina HumanHT-12 V4.0 expression beadchip technology with $n=5$ normal vulvar tissues and $n=13$ vulvar SCC (squamous cell carcinoma) tissues; analysis was performed using platform GPL10558. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether GATM expression was significantly different between normal vulva and vulvar cancer using a two-tailed t-test.

Results

We mined published microarray data (1, 2) to understand the most significant transcriptional changes between the normal vulva and cancer of the vulva.

GATM is differentially expressed in vulvar cancer.

We found that glycine amidinotransferase, encoded by GATM, was among the most differentially expressed genes, transcriptome-wide, in cancer of the vulva (1). When sorting each transcript whose expression was measured across the transcriptome based on significance of change between the normal vulva and vulvar cancer tissues, GATM ranked 80 out of 22277 total transcripts, equating to 99.6% differential expression (Chart 1) (1). Differential expression of GATM was statistically significant (Chart 1; $p=7.81E-06$).

We validated our findings by performing this same analysis using a second published microarray dataset (2). When sorting each transcript whose expression was measured across the transcriptome based on significance of change between the normal vulva and vulvar cancer tissues, GATM ranked 1177 out of 47323 total transcripts, equating to __% differential expression (Chart 2) (2). Differential expression of GATM was again statistically significant (Chart 2; $p=7.70E-05$).

GATM expression is higher in cancer tissue as compared to the vulva.

We obtained exact mRNA expression data from the first microarray dataset we mined to directly compare expression levels between control and cancer tissues. GATM was expressed at significantly higher levels in cancer of the vulva as compared to normal vulva. GATM was expressed at 8.16 ± 0.94 arbitrary units (A.U.) in normal, but at 5.13 ± 0.70 A.U. in vulvar cancer tissues (Figure 1). We calculated

a mean fold change of 0.63 in GATM expression when comparing vulvar carcinoma to vulva.

Discussion

We documented here that GATM is differentially expressed at the transcriptome level in cancer of the human vulva (1, 2). GATM is a candidate gene of interest in understanding the transcriptional biology of vulvar cancer.

References

1. Pappa, K.I., Polyzos, A., Jacob-Hirsch, J., Amariglio, N., Vlachos, G.D., Loutradis, D. and Anagnou, N.P., 2015. Profiling of discrete gynecological cancers reveals novel transcriptional modules and common features shared by other cancer types and embryonic stem cells. *PLoS One*, 10(11), p.e0142229.
2. Micci, F., Panagopoulos, I., Haugom, L., Dahlback, H.S.S., Pretorius, M.E., Davidson, B., Abeler, V.M., Tropé, C.G., Danielsen, H.E. and Heim, S., 2013. Genomic aberration patterns and expression profiles of squamous cell carcinomas of the vulva. *Genes, Chromosomes and Cancer*, 52(6), pp.551-563.
3. Canavan, T.P. and Cohen, D., 2002. Vulvar cancer. *American family physician*, 66(7), p.1269.
4. Grimes, D.A. and Economy, K.E., 1995. Primary prevention of gynecologic cancers. *American journal of obstetrics and gynecology*, 172(1), pp.227-235.
5. Weinberg, D. and Gomez-Martinez, R.A., 2019. Vulvar cancer. *Obstetrics and Gynecology Clinics*, 46(1), pp.125-135.

Rank: 80
Probe ID: 203178_at
p-value: 7.81E-06
t: 6.995402
B: 4.02336
Gene: GATM
Gene name: glycine amidinotransferase

Chart 1: Differential expression of GATM in vulvar cancer.

Rank of differential expression, probe ID, *p*-value of global differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Rank: 1177
Probe ID: ILMN_1729180
p-value: 7.70E-05
t: 5.0611134
B: 1.50592
Gene: GATM
Gene name: glycine amidinotransferase

Chart 2: Differential expression of GATM in vulvar cancer.

Rank of differential expression, probe ID, *p*-value of global differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Figure 1: GATM is expressed at significantly higher levels in vulvar cancer as compared to normal vulva.

The mRNA expression level of GATM in the normal vulva (left) and in vulvar carcinoma (right) is graphically depicted with the result of a statistical test evaluating significance of difference in GATM expression between the two groups, a *p*-value, listed above.