Differential expression of fibronectin type III domain containing 5 in human epithelial ovarian cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Epithelial ovarian cancer (EOC) is the most lethal gynecologic cancer (1). We performed discovery of genes associated with epithelial ovarian cancer and of the high-grade serous ovarian cancer (HGSC) subtype, using published and public microarray data (2, 3) to compare global gene expression profiles of normal ovary or fallopian tube with that of primary tumors from women diagnosed with epithelial ovarian cancer or HGSC. We identified the gene encoding fibronectin type III domain containing 5, FNDC5, as among the genes whose expression was most different in epithelial ovarian cancer as compared to the normal fallopian tube. FNDC5 expression was significantly lower in high-grade serous ovarian tumors relative to normal fallopian tube. These data indicate that expression of FNDC5 is perturbed in epithelial ovarian cancers broadly and in ovarian cancers of the HGSC subtype. FNDC5 may be relevant to pathways underlying ovarian cancer initiation (transformation) or progression.

Keywords: ovarian cancer, epithelial ovarian cancer, HGSC, high-grade serous ovarian cancer, systems biology of ovarian cancer, targeted therapeutics in ovarian cancer.

The five-year survival rate for women diagnosed with high-grade serous ovarian cancer is between 30-40% and has not changed significantly in decades (4, 5). The development of novel, targeted therapeutics to treat HGSC can be facilitated by an enhanced understanding of the transcriptional behavior of ovarian tumors relative to that of the normal ovary. We mined published and public microarray data (2, 3) to compare global gene expression profiles between human ovarian tumors, including that of the HGSC subtype, and that of normal ovarian and fallopian tissue. We identified the gene encoding fibronectin type III domain containing 5, FNDC5, as among the most differentially expressed in HGSC tumors of the ovary and in epithelial ovarian cancer broadly. FNDC5 may be a gene of interest when prioritizing the study of target genes and pathways for the development of novel therapeutic interventions in epithelial ovarian cancer and specifically in high-grade serous ovarian cancers.

Methods

We used microarray data from datasets GSE124766 (2) and GSE126308 (3) for this differential gene expression analysis of epithelial ovarian cancer in conjunction with GEO2R. The Benjamini and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of global differential expression. Log transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. GSE124766 (2) was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=3$ normal fallopian tube tissue (control) and $n=8$ for tumors from patients with high-grade serous ovarian cancer. GSE126308 (3) was generated using Affymetrix Human Transcriptome Array 2.0 with $n=20$ tumors from high-grade serous adenocarcinoma (HGSC) patients classified as early in disease progression and $n=30$ for tumors from HGSC patients classified as “late” in disease progression; analysis was performed using platform GPL17586.

A statistical test was performed to evaluate whether FNDC5 expression was significantly different when comparing normal ovarian tissue from control subjects and primary tumors from women diagnosed with HGSC using a two-tailed, unpaired t-test with Welch’s correction. We used PRISM for all statistical analyses (Version 8.4.0)(455).

Results

We mined published and public microarray data (2, 3) to identify differentially expressed genes in epithelial ovarian cancer and specifically in high-grade serous ovarian cancer (HGSC), the type of ovarian cancer responsible for 70-80% of deaths resulting from the most lethal gynecologic malignancy.

FNDC5 is differentially expressed in ovarian tumors from women diagnosed with HGSC

We identified FNDC5 as among the genes whose expression was most different when comparing primary HGSC tumors to normal fallopian tube tissue (2) (Chart 1). When sorting all of the transcripts measured by microarray based on change in expression between HGSC and the normal fallopian tube, in this data set, FNDC5 ranked 113 out of 40481 total transcripts (Chart 1), equating to 99.7% differential expression. Differential expression of FNDC5 in HGSC tumors was statistically significant (Chart 1; $p=2.78E-06$).

We analyzed a second microarray dataset (3) generated using normal ovarian tissues and tumors from women diagnosed with epithelial ovarian cancer (EOC) to determine whether differential expression of FNDC5 could be observed in tumors from a different group of patients, which included HGSC tumors.

1 We found that FNDC5 was differentially expressed in epithelial ovarian cancers when analyzing
2 based on disease progression (Chart 2). When sorting all of the transcripts measured by microarray based
3 on change in expression between tumors from patients early in disease progression and tumors from
4 patients late in disease progression, FNDC5 ranked 9594 out of 70523 total transcripts (Chart 2), equating
5 to 86.4% differential expression. Differential expression of FNDC5 in EOC based on disease progression
6 trended towards statistical significance (Chart 2; $p=3.91E-02$).

7 FNDC5 is expressed at significantly lower levels in HGSC when compared to the normal fallopian tubes.

8 We then extracted exact mRNA expression levels for the differentially expressed FNDC5 transcript
9 from both normal fallopian tube tissue and from epithelial ovarian tumors. FNDC5 was expressed at
10 significantly lower levels in high-grade serous ovarian cancers when compared to the normal fallopian
11 tube (Figure 1; $p=0.0055$). We calculated a mean fold change of 0.65 in FNDC5 expression when
12 comparing HGSC tumors to normal fallopian tissue, as FNDC5 was expressed at 9.39 ± 0.71 arbitrary
13 units (AU) in the fallopian tubes but at 6.14 ± 0.57 AU in tumors of the ovary.

14 Thus, we found using published and public microarray data (2, 3) that FNDC5 was among the
15 genes most differentially expressed in tumors from women with high-grade serous carcinomas, and in
16 epithelial ovarian cancer broadly. FNDC5 expression was significantly lower in HGSC tumors when
17 compared to the normal fallopian tubes.

18 Discussion

19 Transcriptional profiling of HGSC in a Cancer Genome Atlas integrated genomic analysis
20 classified HGSC into four subtypes based on gene clustering: immunoreactive, differentiated,
21 proliferative, and mesenchymal (6). We sought to describe the transcriptional landscape of epithelial
22 ovarian cancers in an unbiased fashion by using published and public microarray data from primary
23 tumors of women diagnosed with HGSC compared to transcriptome data from normal ovarian and
24 fallopian tube tissue (2, 3). In both datasets analyzed (2, 3), FNDC5 was among the genes whose
25 expression changed most significantly when comparing normal gynecological tissue to ovarian tumors,
26 from women with epithelial ovarian cancer as well as specifically from women with HGSC. FNDC5 may
27 be a molecule of relevance in understanding transformation or disease progression in human epithelial
28 ovarian cancer.

References

1. Guppy, A.E., Nathan, P.D. and Rustin, G.J., 2005. Epithelial ovarian cancer: a review of current management. *Clinical Oncology*, 17(6), pp.399-411.
2. Zhang, W., Klinebiel, D., Barger, C.J., Pandey, S., Guda, C., Miller, A., Akers, S.N., Odunsi, K. and Karpf, A.R., 2020. Global DNA hypomethylation in epithelial ovarian cancer: passive demethylation and association with genomic instability. *Cancers*, 12(3), p.764.
3. Biomarkers in older ovarian cancer patients. GSE126308.
<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE126308>. Sambavy Nadaraja. Odense University Hospital.
4. Bowtell, D.D., Böhm, S., Ahmed, A.A., Aspuria, P.J., Bast Jr, R.C., Beral, V., Berek, J.S., Birrer, M.J., Blagden, S., Bookman, M.A. and Brenton, J.D., 2015. Rethinking ovarian cancer II: reducing mortality from high-grade serous ovarian cancer. *Nature reviews Cancer*, 15(11), pp.668-679.
5. Vaughan, S., Coward, J.I., Bast, R.C., Berchuck, A., Berek, J.S., Brenton, J.D., Coukos, G., Crum, C.C., Drapkin, R., Etemadmoghadam, D. and Friedlander, M., 2011. Rethinking ovarian cancer: recommendations for improving outcomes. *Nature Reviews Cancer*, 11(10), pp.719-725.
6. Cancer Genome Atlas Research Network, 2011. Integrated genomic analyses of ovarian carcinoma. *Nature*, 474(7353), p.609.

Rank: 113
Probe ID: A_23_P420326
p-value: 2.78E-06
t: 8.3500004
B: 5.09717
Gene: FNDC5
Gene name: fibronectin type III domain containing 5

Chart 1: FNDC5 is differentially expressed in high-grade serous ovarian carcinomas (HGSC).

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Rank: 9594
Probe ID: TC01002476.hg.1
p-value: 3.91E-02
t: -2.12
B: -3.80861
Gene: FNDC5
Gene name: fibronectin type III domain containing 5

Chart 2: FNDC5 is differentially expressed in epithelial ovarian cancers when comparing tumors based on degree of disease progression - “early” versus “late”.

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Figure 1: FNDC5 is expressed at significantly lower levels in HGSC tumors when compared to normal fallopian tube.

The mRNA expression of FNDC5 in normal fallopian tube tissues (left) and in the primary tumors of patients with HGSC (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in FNDC5 expression between the normal fallopian tube and primary tumors from patients with HGSC, a *p*-value, listed above.