1 Commentary

2 COVID-19, Ebola Virus Disease and Nipah Virus Infection Reclassification as Novel
3 Acute Immune Dysrhythmia Syndrome (n-AIDS): Potential Curative Role for
4 Immunomodulators.

5 Mina T. Kelleni, MD, PhD

6 Pharmacology Department, College of Medicine, Minia University, Egypt.

7 Mobile: +201200382422

8 drthabetpharm@yahoo.com, mina.kelleni@mu.edu.eg

9 <https://orcid.org/0000-0001-6290-6025>

10 Highlights

11 COVID-19 and selected other fatal diseases are known to disrupt the immune system

12 Monocytic dysrhythmia and altered Th1/Th2 balance trigger COVID-19 mortality

13 n-AIDS is manifested by lymphopenia, causing ARDS and multi-inflammatory syndrome

14 Para COVID-19 syndrome describes potential latent immune related complications

15 Immunomodulators might be the cure of COVID-19 and other fatal diseases

16 Abstract

17 In this manuscript, a suggested reclassification of COVID-19, Ebola virus disease, Nipah
18 virus infection, SARS, and MERS to be considered as a novel acute onset immune
19 dysrhythmia syndrome (n-AIDS) due to altered monocytic, Th1/Th2 as well as cytokines

and chemokines balances is provided. n-AIDs is postulated to be the cause of the acute respiratory distress syndrome and multi-inflammatory syndrome described with COVID-19 and potential curative immunomodulators are described for the mentioned diseases as well as for other disorders caused by Th1/Th2 imbalance. Meanwhile, para COVID-19 syndrome is suggested to describe various immune-related disorders that are associated with SARS CoV-2 infection whether before or after recovery and to embrace a potential of a latent infection that might be discovered later as occurred with Ebola virus disease. Notably, our hypothesis has evolved out of our real-life practice that uses immunomodulatory drugs to manage COVID-19 safely and effectively.

Keywords: COVID-19, Ebola virus disease, Nipah virus infection, n-AIDS, Para CoVID-19 syndrome.

40 Almost three decades ago, a brilliant viewpoint has suggested that a dysregulated
41 immunological switch in favor of Th2 type responses over the Th1 type response is
42 associated with progression of HIV to AIDS and this switch was characterized by loss of
43 the protective antiviral IL-2- and IFN- γ production; Furthermore, those expert researchers
44 have suggested that thousands of seronegative, HIV-exposed (many on multiple occasions)
45 individuals have generated strong Th1 dependent IL-2 responses to HIV antigens with [1].
46 Interestingly, a similar imbalance in Th1/Th2 types with a Th2 favorable switch or failure
47 in the activation of Th1 and reduced IFN- γ production was observed in deceased SARS
48 patients or critically ill MERS patients, whereas a Th1 strong response, which is pivotal in
49 mediating virus-specific adaptive immunity, was observed in mild patients. Importantly,
50 CD4 + and CD8 + T cell cytokines were significantly diminished in COVID-19 patients as
51 compared to healthy controls and subverted T cell composition and/or homeostasis were
52 also suggested to share in COVID-19 pathogenesis and IL-6 was implied to modulate this
53 immunopathological process in patients suffering from severe or critical COVID-19[2,3].
54 Notably, SARS CoV-2 was suggested to dysregulate the antiviral immune response at an
55 early stage leading to number depletion and functional exhaustion of NK and CD8+ T cells
56 which were restored in those who survived and a recommendation to improve the immune
57 response at the early stage of SARS CoV-2 infection was concluded[4]. However, Th2,
58 Th17 cell, and Treg percentages were significantly lower in deceased COVID-19 cases
59 than recovered and healthy control and a variable immunological response was suggested
60 to predict mortality in COVID-19 patients[5], but COVID-19 high morbidity and mortality
61 were suggested to be related to low Th1 immunity[6] and spike-specific Th1 cells capable

of IL7-dependent homeostatic proliferation was shown to predict survival from severe COVID-19[7].

Notably, ACE2 was suggested to regulate the immune response in SARS and SARS CoV-2 including activation of B cells, macrophages, Th1 cells and the inhibition of Treg cells and CD8 + T cells[8] to be correlated that it has been suggested that ACE2[9] and other discovered[10] and yet to be discovered genetic polymorphisms might be reflected through different T cell virus specific and other immunological responses. Thus, while some SARS CoV-2 exposed patients would remain asymptomatic, including asymptomatic seronegative COVID-19 patients[11], others suffer from mild-moderate or severe COVID-19. Similarly, some COVID-19 patients will recover smoothly while others complain of post COVID-19 autoimmune complications hypothesized to be due to transient immunosuppression of innate and acquired immunity[12].

Moreover, we have recently suggested a new terminology for SARS CoV-2 induced dysregulated immune response; monocytic dysrhythmia[13] and an imbalanced immune-inflammatory response was previously described to drive development of COVID-19[14]. Furthermore, we suggested to name para COVID-19 syndrome [15] to embrace a potential that SARS CoV-2 might persist latent, for yet unspecified period, in some cells and tissues[16] and/or a capability to induce immune-mediated disorders such as what is currently being described of several post COVID-19 diseases affecting the nervous system [17,18] or reactivation of various types of herpes viruses which are described in critically ill COVID-19 patients [19] and we recommend further investigations to assess potential SARS CoV-2 direct latency or indirect persistent functional dysrhythmia; respectively in some immune cells such as the migrating interstitial macrophages[20,21] as it is already

well known how SARS CoV-2 possesses several adaptive differences from other coronaviruses to be noted that our knowledge about RNA viruses and their capabilities to remain latent for long duration is still evolving and it might eventually resemble the newly described latent Ebola virus [<https://www.sciencemag.org/news/2021/03/new-ebola-outbreak-likely-sparked-person-infected-5-years-ago>].

Furthermore, a fourth RNA virus induced fatal disease; Ebola virus disease might also be considered to possess a similar potential as regards to its induced dysregulation of the immune system [22], its long lasting T and B cell immunological dysfunction which was further described in Ebola survivors[23] as well as the dysregulated inflammatory and immunological immune response in both Ebola virus disease survivor and deceased cases[24]. Additionally, a fifth RNA virus with a high fatality rate; Nipah virus might be similarly reclassified as it has been shown to modulate the inflammatory and immunological response including the interferon homeostasis[25]. Moreover, Nipah virus was shown to modulate the pro-inflammatory and leucocyte attracting cytokines in a manner that determines the disease course[26] and to induce a dysregulated immune recruitment that led to acute vasculitis among other several induced immune-dysregulatory mechanisms[27].

Interestingly, we postulated that abnormal cytokine and chemokine, known and yet to be discovered, dependent lymphocyte distraction (clinically manifested by lymphopenia) into (causing ARDS) or away from the lungs (causing multiple inflammatory syndrome) might reason for COVID-19 pathogenesis and complications [28] and we would like to suggest that SARS, MERS and SARS CoV-2; the three virulent RNA corona viruses which emerged in the past two decades, which are also anticipated to be joined by other potentially

fatal similar viruses, and Ebola virus disease as well as Nipah virus infection should be considered for a novel immunopathological reclassification that acknowledges their main cause of complications and/or fatalities to be related to their peculiar immune monocytic, Th1/Th2, and potentially other immune cells dysrhythmia that though seems hyperactive, it is practically deficient/incompetent ultimately leading to an acute potentially fatal response; n-AIDS and probable latent effects as recently shown for Ebola and necessitates further investigations to be fully explored for SARS CoV-2; para COVID-19 syndrome. Furthermore, though the three potentially fatal coronaviruses share some similarities with HIV[29], and that other potential similarities between SARS CoV-2 and Nipah virus have been also described [30]; these viruses and Ebola virus also differ from HIV in several important aspects including their specific immunological targets and their main tendency for acute progressive onset and complications.

Taken together, we suggest that our suggested novel classification of acute immune-dysrhythmic syndrome might properly guide us in our quest for a cure as it is only when we know the cause, we can figure out the cure and we recommend to focus on immune-modulation as a potential effective COVID-19 therapy[6,13,31], Ebola virus disease[32] and Nipah virus infection[25] and we hypothesize that our evolved real-life immunomodulatory COVID-19 management protocol[31] that guided us to this hypothesis through its remarkable clinical efficacy against COVID-19 might be also beneficial when tested in clinical trials for early management of other RNA viruses that cause n-AIDS as well as other diseases caused by altered Th1/Th2 balance.

131 **Acknowledgments**

132 This work is dedicated to the soul of Dr. George Habib Bebawi (1938 – 2021); a great
133 Egyptian American world-renowned theologian, patristic scholar and a former director of
134 studies at the Institute for Christian Orthodox Studies, at Cambridge, England. He has
135 bravely and relentlessly fought for what he truly believed orthodox while struggling against
136 a lifelong oppression and injustice that has not ended until his departure. May his memory
137 be everlasting.

138 **Funding**

139 None

140 **Conflicts of interest/Competing interests**

141 The author declares no competing interests

142 **Availability of data and material**

143 N/A

144 **Code availability**

145 N/A

146 **Ethics approval**

147 N/A

148 **Consent to participate**

149 N/A

150 **Consent for publication**

151 N/A

152 **References**

- 153 1. Clerici M, Shearer GM. A TH1→TH2 switch is a critical step in the etiology of HIV
154 infection. *Immunology Today*. 1993 1993/01/01/;14(3):107-111.
- 155 2. Zhang Y-y, Li B-r, Ning B-t. The Comparative Immunological Characteristics of SARS-CoV,
156 MERS-CoV, and SARS-CoV-2 Coronavirus Infections [Review]. *Front Immunol*. 2020
157 2020-August-14;11(2033).
- 158 3. Rupp J, Dreo B, Gutl K, et al. T Cell Phenotyping in Individuals Hospitalized with COVID-
159 19. *J Immunol*. 2021 Feb 8.
- 160 4. Zheng M, Gao Y, Wang G, et al. Functional exhaustion of antiviral lymphocytes in COVID-
161 19 patients. *Cellular & Molecular Immunology*. 2020 2020/05/01;17(5):533-535.
- 162 5. Sami R, Fathi F, Eskandari N, et al. Characterizing the immune responses of those who
163 survived or succumbed to COVID-19: Can immunological signatures predict outcome?
164 *Cytokine*. 2021 Apr;140:155439.
- 165 6. Gupta A. Is Immuno-modulation the Key to COVID-19 Pandemic? *Indian Journal of*
166 *Orthopaedics*. 2020 2020/05/01;54(3):394-397.
- 167 7. Neidleman J, Luo X, George AF, et al. Distinctive features of SARS-CoV-2-specific T cells
168 predict recovery from severe COVID-19. *medRxiv*. 2021 Jan 28.
- 169 8. Luo J, Lu S, Yu M, et al. The potential involvement of JAK-STAT signaling pathway in the
170 COVID-19 infection assisted by ACE2. *Gene*. 2021 Feb 5;768:145325.
- 171 9. Kelleni MT. ACE2 polymorphisms interplay with the apelinergic peptide system:
172 potential tools for COVID-19 diagnosis and treatment. *OSFPREPRINTS (Preprint)*. 2021.
- 173 10. Pairo-Castineira E, Clohisey S, Klaric L, et al. Genetic mechanisms of critical illness in
174 Covid-19. *Nature*. 2020 2020/12/11.
- 175 11. Lou B, Li T-D, Zheng S-F, et al. Serology characteristics of SARS-CoV-2 infection since
176 exposure and post symptom onset. *European Respiratory Journal*. 2020:2000763.
- 177 12. Cañas CA. The triggering of post-COVID-19 autoimmunity phenomena could be
178 associated with both transient immunosuppression and an inappropriate form of
179 immune reconstitution in susceptible individuals. *Med Hypotheses*. 2020;145:110345-
180 110345.
- 181 13. Kelleni MT. Non-steroidal Anti-inflammatory Drugs/nitazoxanide/azithromycin Potential
182 Beneficial COVID-19 Effects: Preventing the Cytokine Storm via Mitigation of the
183 Interleukin-6 Amplifier and Monocytic Immunological Dysrhythmia. *OSFPREPRINTS*
184 (Preprint). 2021.
- 185 14. Blanco-Melo D, Nilsson-Payant BE, Liu W-C, et al. Imbalanced Host Response to SARS-
186 CoV-2 Drives Development of COVID-19. *Cell*. 2020;181(5):1036-1045.e9.
- 187 15. Kelleni M. SARS CoV-2 Might Exploit Cells of the Innate Immune System to Induce the
188 Novel Acute Immune Dysrhythmic Syndrome (n-AIDS) and Para COVID-19 Syndrome.
189 *OSFPREPRINTS (Preprint)*. 2021.
- 190 16. Pietsch H, Escher F, Aleshcheva G, et al. Proof of SARS-CoV-2 genomes in
191 endomyocardial biopsy with latency after acute infection. *International Journal of*
192 *Infectious Diseases*. 2021 2021/01/01/;102:70-72.

193 17. Pezzini A, Padovani A. Lifting the mask on neurological manifestations of COVID-19.
194 Nature Reviews Neurology. 2020 2020/11/01;16(11):636-644.

195 18. Taquet M, Geddes JR, Husain M, et al. 6-month neurological and psychiatric outcomes in
196 survivors of COVID-19: a retrospective cohort study using electronic health records. The
197 Lancet Psychiatry.

198 19. Simonnet A, Engelmann I, Moreau AS, et al. High incidence of Epstein–Barr virus,
199 cytomegalovirus, and human-herpes virus-6 reactivations in critically ill patients with
200 COVID-19. Infectious Diseases Now. 2021 2021/01/18/.

201 20. Tavazzi G, Pellegrini C, Maurelli M, et al. Myocardial localization of coronavirus in
202 COVID-19 cardiogenic shock [<https://doi.org/10.1002/ejhf.1828>]. European Journal of
203 Heart Failure. 2020 2020/05/01;22(5):911-915.

204 21. Liegeois M, Legrand C, Desmet CJ, et al. The interstitial macrophage: A long-neglected
205 piece in the puzzle of lung immunity. Cellular Immunology. 2018 2018/08/01;330:91-
206 96.

207 22. Falasca L, Agrati C, Petrosillo N, et al. Molecular mechanisms of Ebola virus
208 pathogenesis: focus on cell death. Cell Death Differ. 2015;22(8):1250-1259.

209 23. Wiedemann A, Foucat E, Hocini H, et al. Long-lasting severe immune dysfunction in
210 Ebola virus disease survivors. Nature Communications. 2020 2020/07/24;11(1):3730.

211 24. Colavita F, Biava M, Castilletti C, et al. Inflammatory and Humoral Immune Response
212 during Ebola Virus Infection in Survivor and Fatal Cases Occurred in Sierra Leone during
213 the 2014–2016 Outbreak in West Africa. Viruses. 2019;11(4):373.

214 25. Pelissier R, Iampietro M, Horvat B. Recent advances in the understanding of Nipah virus
215 immunopathogenesis and anti-viral approaches [version 1; peer review: 3 approved].
216 F1000Research. 2019;8(1763).

217 26. Satterfield BA, Cross RW, Fenton KA, et al. The immunomodulating V and W proteins of
218 Nipah virus determine disease course. Nature Communications. 2015
219 2015/06/24;6(1):7483.

220 27. Prescott J, de Wit E, Feldmann H, et al. The immune response to Nipah virus infection.
221 Arch Virol. 2012;157(9):1635-1641.

222 28. Kelleni MT. Early use of non-steroidal anti-inflammatory drugs in COVID-19 might
223 reverse pathogenesis, prevent complications and improve clinical outcomes. Biomed
224 Pharmacother. 2021 Jan;133:110982.

225 29. Saleemi MA, Ahmad B, Benchoula K, et al. Emergence and molecular mechanisms of
226 SARS-CoV-2 and HIV to target host cells and potential therapeutics. Infect Genet Evol.
227 2020;85:104583-104583.

228 30. Roe K. Explanation for COVID-19 infection neurological damage and reactivations.
229 Transbound Emerg Dis. 2020;67(4):1414-1415.

230 31. Kelleni MT. NSAIDs/Nitazoxanide/Azithromycin Immunomodulatory Protocol Used in
231 Adults, Children and Pregnant COVID-19 Patients: An Egyptian Prospective
232 Observational Study. OSFPREPRINTS (Preprint). 2021.

233 32. Bixler SL, Duplantier AJ, Bavari S. Discovering Drugs for the Treatment of Ebola Virus.
234 Curr Treat Options Infect Dis. 2017;9(3):299-317.