Differential expression of ZSCAN18 in cancer of the vulva.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

In these brief notes we document work using published microarray data (1, 2) to pioneer integrative transcriptome analysis comparing vulvar carcinoma to its tissue of origin, the vulva. We report the differential expression of zinc finger and SCAN domain containing 18, encoded by ZSCAN18, in cancer of the vulva. ZSCAN18 may be of pertinence to understanding transformation and disease progression in vulvar cancer (3).

Keywords: vulvar cancer, vulvar carcinoma, ZSCAN18, zinc finger and SCAN domain containing 18.

Gynecologic cancers include cervical cancer, ovarian cancer, endometrial (uterine) cancer, and vulvar cancer (4, 3). Vulvar cancer is thought to be the least prevalent of these, with 6190 cases reported in the United States in 2018 (5). We integrated separate published microarray datasets to understand, at the systems level, the transcriptional landscape of vulvar carcinoma.

Methods

We utilized microarray datasets GSE63678 (1) and GSE38228 (2) for this integrated, transcriptome-level differential gene expression analysis of human vulvar cancers in conjunction with GEO2R. GSE63678 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=7$ normal vulvar tissues and $n=6$ vulvar carcinoma tissues; analysis was performed using platform GPL571. GSE38228 was generated using Illumina HumanHT-12 V4.0 expression beadchip technology with $n=5$ normal vulvar tissues and $n=13$ vulvar SCC (squamous cell carcinoma) tissues; analysis was performed using platform GPL10558. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether ZSCAN18 expression was significantly different between normal vulva and vulvar cancer using a two-tailed t -test.

Results

We mined published microarray data (1, 2) to understand the most significant transcriptional changes between the normal vulva and cancer of the vulva.

ZSCAN18 is differentially expressed in vulvar cancer.

We found that zinc finger and SCAN domain containing 18, encoded by ZSCAN18, was among the most differentially expressed genes, transcriptome-wide, in cancer of the vulva (1). When sorting each transcript whose expression was measured across the transcriptome based on significance of change between the normal vulva and vulvar cancer tissues, ZSCAN18 ranked 39 out of 22277 total transcripts, equating to 99.8% differential expression (Chart 1) (1). Differential expression of ZSCAN18 was statistically significant (Chart 1; $p=2.65E-06$).

We validated our findings by performing this same analysis using a second published microarray dataset (2). When sorting each transcript whose expression was measured across the transcriptome based on significance of change between the normal vulva and vulvar cancer tissues, ZSCAN18 ranked 24 out of 47323 total transcripts, equating to 99.9% differential expression (Chart 2) (2). Differential expression of ZSCAN18 was again statistically significant (Chart 2; $p=3.05E-12$).

ZSCAN18 expression is higher in cancer tissue as compared to the vulva.

We obtained exact mRNA expression data from the first microarray dataset we mined to directly compare expression levels between control and cancer tissues. ZSCAN18 was expressed at significantly higher levels in cancer of the vulva as compared to normal vulva. ZSCAN18 was expressed at 9.47 ± 0.44 arbitrary units (A.U.) in normal vulva, but at 7.04 ± 0.75 A.U. in vulvar cancer tissues (Figure 1). We

1 calculated a mean fold change of 0.74 in ZSCAN18 expression when comparing vulvar carcinoma to
2 vulva.

3 **Discussion**

4 We documented here that ZSCAN18 is differentially expressed at the transcriptome level in cancer
5 of the human vulva (1, 2). ZSCAN18 is a candidate gene of interest in understanding the transcriptional
6 biology of vulvar cancer.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

References

1. Pappa, K.I., Polyzos, A., Jacob-Hirsch, J., Amariglio, N., Vlachos, G.D., Loutradis, D. and Anagnou, N.P., 2015. Profiling of discrete gynecological cancers reveals novel transcriptional modules and common features shared by other cancer types and embryonic stem cells. *PLoS One*, 10(11), p.e0142229.
2. Micci, F., Panagopoulos, I., Haugom, L., Dahlback, H.S.S., Pretorius, M.E., Davidson, B., Abeler, V.M., Tropé, C.G., Danielsen, H.E. and Heim, S., 2013. Genomic aberration patterns and expression profiles of squamous cell carcinomas of the vulva. *Genes, Chromosomes and Cancer*, 52(6), pp.551-563.
3. Canavan, T.P. and Cohen, D., 2002. Vulvar cancer. *American family physician*, 66(7), p.1269.
4. Grimes, D.A. and Economy, K.E., 1995. Primary prevention of gynecologic cancers. *American journal of obstetrics and gynecology*, 172(1), pp.227-235.
5. Weinberg, D. and Gomez-Martinez, R.A., 2019. Vulvar cancer. *Obstetrics and Gynecology Clinics*, 46(1), pp.125-135.

Rank: 39
Probe ID: 218312_s_at
p-value: 2.65E-06
t: 7.72133
B: 5.04079
Gene: ZSCAN18
Gene name: zinc finger and SCAN domain containing 18

Chart 1: Differential expression of ZSCAN18 in vulvar cancer.

Rank of differential expression, probe ID, *p*-value of global differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Rank: 24
Probe ID: ILMN_1654946
p-value: 3.05E-12
t: 16.0477584
B: 17.83728
Gene: ZSCAN18
Gene name: zinc finger and SCAN domain containing 18

Chart 2: Differential expression of ZSCAN18 in vulvar cancer.

Rank of differential expression, probe ID, *p*-value of global differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Figure 1: ZSCAN18 is expressed at significantly higher levels in vulvar cancer as compared to normal vulva.

The mRNA expression level of ZSCAN18 in the normal vulva (left) and in vulvar carcinoma (right) is graphically depicted with the result of a statistical test evaluating significance of difference in ZSCAN18 expression between the two groups, a *p*-value, listed above.