A network of olfactory receptors is differentially expressed in CD24+ CD73+ V γ 2+ $\gamma\delta$ -T cells

Shahan Mamoor, MS

Abstract

To evaluate the transcriptional consequences of differential V γ chain usage in $\gamma\delta$ -T cells, I used a publicly available dataset to compare the global gene expression profiles of immature, committed CD24+ CD73+ $\gamma\delta$ -T cells bearing either V γ 1.1+ or V γ 2+ at the $\gamma\delta$ TCR. I found that thirty-nine genes encoding olfactory receptors were among the most differentially expressed in V γ 2+ CD24+ CD73+ cells. Four vomeronasal receptors (proteins involved in pheromone detection) were also transcriptionally up-regulated in V γ 2+ CD24+ CD73+ cells compared to their V γ 1.1+ counterparts. Analysis of two further datasets revealed two other olfactory genes that were differentially expressed in $\gamma\delta$ -T cells, in these cases not based on V γ chain usage but in a stage-selective pattern. To my knowledge this is the first report documenting the differential expression of genes encoding olfactory receptors during $\gamma\delta$ -T development and based on V γ chain usage.

Keywords: $\gamma\delta$ -T cell, olfactory receptor, vomeronasal receptor, TCR V γ 2

Introduction

$\gamma\delta$ -T cells are present in mice by embryonic day 14 (16), and persist throughout adulthood. In mice and in humans, they reside in the fetus, the peripheral blood, in tissues, and at mucosal surfaces (5). $\gamma\delta$ -T cells are distinguished from $\alpha\beta$ T-cells by the composition of the T-cell receptor displayed on the cell surface. Tcr rearrangements that produce γ and δ TCR chains can generate a $\gamma\delta$ TCR, while Tcr rearrangements that produce α and β TCR chains can generate an $\alpha\beta$ TCR. Thymocytes become fully committed to the T-cell fate at stage DN2b (CD44+ CD25+), as DN1 and DN2a thymocytes maintain the capacity to differentiate into NK cells, monocytes, mast cells and dendritic cells. During stage DN3, TCR γ and TCR β rearrangements occur, but since TCR α rearrangements do not occur until the DP stage, $\gamma\delta$ -T cells have the theoretical capacity to differentiate into a $\gamma\delta$ TCR bearing lymphocyte while β -selection is still occurring (during the transition of DN4 to DP) (7).

Despite the fact that the majority of mature T cells are of the $\alpha\beta$ -lineage (3), the temporal kinetics of Tcr γ/δ rearrangement confer an advantage to the developing thymocyte in choosing the $\gamma\delta$ T-cell fate, as the instructive model of $\alpha\beta/\gamma\delta$ -T cell lineage commitment argues that once a T-cell expresses a single functional TCR ($\gamma\delta$ or $\alpha\beta$), it is thus designated a T-cell of that brand (4). Nevertheless, a study in which the TCR β transgene was overexpressed *in vivo* found T-cells that resembled $\gamma\delta$ -T cells but that expressed an $\alpha\beta$ TCR (1, 18), putting at odds the claim that the presence of an $\alpha\beta$ TCR on a particular cell would preclude it from joining the $\gamma\delta$ -T lineage. Similarly, another study utilizing a knockout of the TCR β locus demonstrated that double positive $\alpha\beta$ -T cells could still be generated despite the absence of a fully assembled $\alpha\beta$ TCR (13). Furthermore, a study that sequenced TCR repertoires revealed that $\alpha\beta$ T-cells can complete the commitment process while Tcr β rearrangement is still occurring (17). Thus, though discrete events leading to the (a) lineage specification of $\gamma\delta$ - versus $\alpha\beta$ -T cells, and (b) differentiation of $\gamma\delta$ T-cells have been characterized (1, 3, 6, 8, 9, 10, 13, 14, 15, 16), the absolute temporal boundaries of $\gamma\delta$ -T cell differentiation have not been established.

With this in mind, I performed comparative gene expression profiling analyses on existing datasets generated using RNA-sequencing or microarray of $\gamma\delta$ -T cells (2, 14, 20), aiming to gain novel insights into the transcriptional changes that accompany $\gamma\delta$ -T cell selection. Through successive comparisons using datasets from independent sources, of $\gamma\delta$ -T cells and thymocytes from multiple developmental stages and $\gamma\delta$ TCR clonotypes I aimed to identify transcripts that might otherwise have been overlooked at the systems level. In the course of these analyses, I found that 39 unique transcripts from the family of olfactory receptors were differentially up-regulated by immature committed CD24+ CD73+ $\gamma\delta$ -T cells that use the V γ 2+ $\gamma\delta$ TCR. Moreover, several receptors of the vomeronasal organ were concomitantly transcriptionally induced. To my knowledge, this is the first evidence of differential transcription of olfactory receptor genes in $\gamma\delta$ -T cells or during the $\gamma\delta$ -T cell maturation process.

Methods

Comparative Gene Expression Profiling

GEO (Gene Expression Omnibus) Dataset 75920 was utilized for this analysis. GEO Datasets 71637 and 86495 were also utilized. The GEO2R web application, when provided with a dataset, identifies 250 transcripts whose expression is most differentially regulated in a given comparison between samples. Log transformation of data was set to “Yes”, and the p-value, when adjusted, was set to “Benjamini-Hochberg”. The “Submitter supplied” category of platform annotation was used to identify gene/RNA names.

Statistics

Statistical analyses comparing the mean RNA expression value of each olfactory receptor identified in each murine cell subset ([1] CD24+ CD73+ V γ 1.1+ $\gamma\delta$ -T cells versus CD24+ CD73+ V γ 2+ $\gamma\delta$ -T cells, [2] E18 $\gamma\delta$ -T cells, wild-type and CD3 double haploinsufficient versus adult $\gamma\delta$ -T cells, wild-type and CD3 double haploinsufficient, and [3] DN3a thymocytes versus $\gamma\delta$ -T cells) were performed using GraphPad Prism Version 8.0. An unpaired t-test with Welch's correction was performed for each gene, and gaussian distribution was assumed (parametric test). P-values for each comparison are denoted on each graph.

Results

Olfactory receptors are among the most differentially expressed genes in V γ 2+ $\gamma\delta$ -T cells relative to V γ 1.1+ $\gamma\delta$ -T cells at the CD24+ CD73+ (immature committed) stage

I performed comparative gene expression profiling on exomic microarray data comparing immature committed (CD24+ CD73+) $\gamma\delta$ -T cells bearing either V γ 2 or V γ 1.1 at the $\gamma\delta$ TCR. Of the 250 most differentially expressed genes between V γ 2+ $\gamma\delta$ -T cells and V γ 1.1+ $\gamma\delta$ -T-cells at the CD24+ CD73+ stage, I identified thirty-nine transcripts of the olfactory receptor family (Table 1 and Figure 1). Each of the thirty-nine transcripts was uniformly up-regulated in the V γ 2+ subset.

Olfactory receptor Olfr157 is expressed in a stage-specific manner in $\gamma\delta$ -T cells, increasing in expression from embryonic day 18 to 6 weeks

Using a separate dataset (14), I performed comparative gene expression analysis on microarray data from 4 cell populations isolated by flow cytometry ($\gamma\delta$ -T cells from embryonic day 18 mice, both wild-type and CD3 double haploinsufficient, and $\gamma\delta$ -T cells from adult mice [6 weeks], both wild-type and CD3 double haploinsufficient). The top 250 most differentially transcribed genes after cross comparison were identified using GEO2R, and a mean RNA expression level for each transcript in each of the subsets (E18 $\gamma\delta$ -T cells and adult $\gamma\delta$ -T cells) was calculated.

Brief Report

Though two genotypes (wild-type and CD3 double haploinsufficient, [CD3 g⁺/- d⁺/-]) were included in each of the subsets, since they were grouped by developmental stage, the comparative analysis identified any transcripts that are similarly regulated across different genotypes.

Again, I identified an olfactory receptor, Olfr157, as one of the top 250 most differentially expressed genes in a global profiling of $\gamma\delta$ -T cells. Here, the analysis was between $\gamma\delta$ -T cells from day 18 embryos and $\gamma\delta$ -T cells from mice 6-weeks of age, suggesting that Olfr157 was expressed in $\gamma\delta$ -T cells in a developmental stage-selective pattern.

Transcript for the olfactory receptor Olfr1507 is present in significantly lower quantities in $\gamma\delta$ -T cells relative to DN3a thymocytes

I performed comparative gene expression profiling with a third dataset, where the authors isolated (among other populations) stage DN3a thymocytes as well as $\gamma\delta$ -T cells (20). Among the top 250 most differentially expressed genes between DN3a thymocytes and $\gamma\delta$ -T cells was Olfr1507, a receptor of the olfactory system (Figure 4). The expression of Olfr1507 was significantly decreased in $\gamma\delta$ -T cells relative to DN3a thymocytes.

Expression of several vomeronasal receptor genes is concomitantly up-regulated in in V γ 2+ $\gamma\delta$ -T cells relative to V γ 1.1+ $\gamma\delta$ -T cells at the CD24+ CD73+ immature committed stage

Finally, using the initial dataset, I identified four genes of the vomeronasal receptor family, V1rh20, V1rg7, V1ra6 and V1rh11, whose expression was differentially increased in CD24+ CD73+ V γ 2+ $\gamma\delta$ -T cells. These transcripts were also among the top 250 most differentially expressed genes between CD24+ CD73+ V γ 2+ $\gamma\delta$ -T cells and CD24+ CD73+ V γ 1.1+ $\gamma\delta$ -T cells. As these genes were not among the top 250 most differentially expressed genes between CD24+ CD73- V γ 2+ $\gamma\delta$ -T cells and CD24+ CD73- V γ 1.1+ $\gamma\delta$ -T cells (data not shown), this differential expression appeared to occur in a stage-specific fashion, with the gain of CD73+.

Discussion

In brief, I report here using three independent publicly available datasets that thirty-nine members of the olfactory receptor family are differentially transcriptionally induced in V γ 2+ CD24+ CD73+ $\gamma\delta$ -T cells. Additionally, one olfactory receptor (Olfr157) is differentially expressed in a stage-selective manner, significantly increasing in quantity from embryonic day 18 to 6 weeks of age. Intriguingly, another olfactory receptor, Olfr1507, is present in significantly less quantity in $\gamma\delta$ -T cells relative to DN3a thymocytes. All of these transcripts of the olfactory receptor family were among the 250 most differentially expressed as determined by global gene expression profiling.

In V γ 2⁺ CD24⁺ CD73⁺ $\gamma\delta$ -T cells, transcript levels of 4 receptors of the vomeronasal pheromone sensing system were also significantly increased and among the most differentially transcribed relative to V γ 1.1⁺ CD24⁺ CD73⁺ $\gamma\delta$ -T cells.

Though no literature could be identified regarding a functional relationship between the olfactory system and $\gamma\delta$ -T cells/Tcr rearrangement, a series of olfactory genes flank the 5' end of the Tcr δ locus (11).

The identification of these differentially expressed transcripts merits future investigation into the potential relationship between the olfactory system, the vomeronasal system and $\gamma\delta$ -T cell fate and/or function.

References

1. Bruno, L., Fehling, H.J. and von Boehmer, H., 1996. The $\alpha\beta$ T cell receptor can replace the $\gamma\delta$ receptor in the development of $\gamma\delta$ lineage cells. *Immunity*, 5(4), pp.343-352.
2. Buus, T.B., Geisler, C. and Lauritsen, J.P.H., 2016. The major diversification of V γ 1. 1⁺ and V γ 2⁺ thymocytes in mice occurs after commitment to the $\gamma\delta$ T-cell lineage. *European journal of immunology*, 46(10), pp.2363-2375.
3. Carding, S.R. and Egan, P.J., 2002. $\gamma\delta$ T cells: functional plasticity and heterogeneity. *Nature reviews immunology*, 2(5), p.336.
4. Ciofani, M., Knowles, G.C., Wiest, D.L., von Boehmer, H. and Zúñiga-Pflücker, J.C., 2006. Stage-specific and differential notch dependency at the $\alpha\beta$ and $\gamma\delta$ T lineage bifurcation. *Immunity*, 25(1), pp.105-116.
5. Ciofani, M. and Zúñiga-Pflücker, J.C., 2010. Determining $\gamma\delta$ versus $\alpha\beta$ T cell development. *Nature Reviews Immunology*, 10(9), p.657.
6. Dave, V.P., Cao, Z., Browne, C., Alarcon, B., Fernandez-Miguel, G., Lafaille, J., de la Hera, A., Tonegawa, S. and Kappes, D.J., 1997. CD3 δ deficiency arrests development of the $\alpha\beta$ but not the $\gamma\delta$ T cell lineage. *The EMBO Journal*, 16(6), pp.1360-1370.
7. Rothenberg, E.V., Moore, J.E. and Yui, M.A., 2008. Launching the T-cell-lineage developmental programme. *Nature Reviews Immunology*, 8(1), p.9.
8. Garbe, A.I., Krueger, A., Gounari, F., Zúñiga-Pflücker, J.C. and von Boehmer, H., 2006. Differential synergy of Notch and T cell receptor signaling determines $\alpha\beta$ versus $\gamma\delta$ lineage fate. *Journal of Experimental Medicine*, 203(6), pp.1579-1590.
9. Hayes, S.M., Li, L. and Love, P.E., 2005. TCR signal strength influences $\alpha\beta$ / $\gamma\delta$ lineage fate. *Immunity*, 22(5), pp.583-593.

10. Itohara, S., Mombaerts, P., Lafaille, J., Iacomini, J., Nelson, A., Clarke, A.R., Hooper, M.L., Farr, A. and Tonegawa, S., 1993. T cell receptor δ gene mutant mice: independent generation of $\alpha\beta$ T cells and programmed rearrangements of $\gamma\delta$ TCR genes. *Cell*, 72(3), pp.337-348.
11. Krangel, M.S., Carabana, J., Abbarategui, I., Schlimgen, R. and Hawwari, A., 2004. Enforcing order within a complex locus: current perspectives on the control of V (D) J recombination at the murine T-cell receptor α/δ locus. *Immunological reviews*, 200(1), pp. 224-232.
12. Kreslavsky, T., Gleimer, M. and von Boehmer, H., 2010. $\alpha\beta$ versus $\gamma\delta$ lineage choice at the first TCR-controlled checkpoint. *Current opinion in immunology*, 22(2), pp.185-192.
13. Livák, F., Wilson, A., MacDonald, H.R. and Schatz, D.G., 1997. $\alpha\beta$ Lineage-committed thymocytes can be rescued by the $\gamma\delta$ T cell receptor (TCR) in the absence of TCR β chain. *European journal of immunology*, 27(11), pp.2948-2958.
14. Muñoz-Ruiz, M., Ribot, J.C., Grosso, A.R., Gonçalves-Sousa, N., Pamplona, A., Pennington, D.J., Regueiro, J.R., Fernández-Malavé, E. and Silva-Santos, B., 2016. TCR signal strength controls thymic differentiation of discrete proinflammatory $\gamma\delta$ T cell subsets. *Nature immunology*, 17(6), p.721.
15. Nuñez-Cruz, S., Aguado, E., Richelme, S., Chetaille, B., Mura, A.M., Richelme, M., Pouyet, L., Jouvin-Marche, E., Xerri, L., Malissen, B. and Malissen, M., 2003. LAT regulates $\gamma\delta$ T cell homeostasis and differentiation. *Nature immunology*, 4(10), p.999.
16. Ribot, J.C., Debarros, A., Pang, D.J., Neves, J.F., Peperzak, V., Roberts, S.J., Girardi, M., Borst, J., Hayday, A.C., Pennington, D.J. and Silva-Santos, B., 2009. CD27 is a thymic determinant of the balance between interferon- γ -and interleukin 17-producing $\gamma\delta$ T cell subsets. *Nature immunology*, 10(4), p.427.
17. Sherwood, A.M., Desmarais, C., Livingston, R.J., Andriesen, J., Haussler, M., Carlson, C.S. and Robins, H., 2011. Deep sequencing of the human TCR γ and TCR β repertoires suggests that TCR β rearranges after $\alpha\beta$ and $\gamma\delta$ T cell commitment. *Science translational medicine*, 3(90), pp.90ra61-90ra61.
18. Terrence, K., Pavlovich, C.P., Matechak, E.O. and Fowlkes, B.J., 2000. Premature expression of T cell receptor (TCR) $\alpha\beta$ suppresses TCR $\gamma\delta$ gene rearrangement but permits development of $\gamma\delta$ lineage T cells. *Journal of Experimental Medicine*, 192(4), pp.537-548.
19. Ueda-Hayakawa, I., Mahlios, J. and Zhuang, Y., 2009. Id3 restricts the developmental potential of $\gamma\delta$ lineage during thymopoiesis. *The Journal of Immunology*, 182(9), pp.5306-5316.
20. Yang, K., Blanco, D.B., Chen, X., Dash, P., Neale, G., Rosencrance, C., Easton, J., Chen, W., Cheng, C., Dhungana, Y. and Anil, K.C., 2018. Metabolic signaling directs the reciprocal lineage decisions of $\alpha\beta$ and $\gamma\delta$ T cells. *Science immunology*, 3(25), p.eaas9818.

Brief Report

Table 1: Rank-ordered list of each olfactory gene identified in differential gene expression analysis of CD24+ CD73+ V γ 1.1+ $\gamma\delta$ -T cells ($n=3$) and CD24+ CD73+ V γ 2+ $\gamma\delta$ -T cells ($n=4$). .

Table 1

Rank (out of 250)	Gene	Rank (out of 250)	Gene
3	Olfr889	43	Olfr1099
5	Olfr1459	50	Olfr1188
8	Olfr805	51	Olfr525
9	Olfr27	52	Olfr2
12	Olfr716	54	Olfr130
13	Olfr218	55	Olfr808
16	Olfr51	57	Olfr1179
22	Olfr1448	62	Olfr464
26	Olfr1023	77	Olfr1408
28	Olfr311	88	Olfr589
30	Olfr1312	95	Olfr910
31	Olfr1325	103	Olfr107
32	Olfr49	111	Olfr187
33	Olfr1365	120	Olfr53
35	Olfr1212	132	Olfr417
36	Olfr1305	144	Olfr695
38	Olfr829	151	Olfr926
39	Olfr568	173	Olfr645
41	Olfr199	178	Olfr807
43	Olfr1099	201	Olfr398

Figure 1: mRNA expression level of each olfactory gene identified in differential gene expression analysis of CD24⁺ CD73⁺ V γ 1.1⁺ $\gamma\delta$ -T cells ($n=3$) and CD24⁺ CD73⁺ V γ 2⁺ $\gamma\delta$ -T cells ($n=4$). Each dot represents one population of cells, and the error bars represent the standard deviation.

Figure 1: (continued)

Figure 2: Summary of mRNA expression levels of each olfactory gene identified in differential gene expression analysis between both cell subsets (CD24+ CD73+ V γ 1.1+ $\gamma\delta$ -T cells and CD24+ CD73+ V γ 2+ $\gamma\delta$ -T cells). Each point represents one population of cells, and the black line represents the mean expression level of each gene in 3 or 4 replicate populations.

Figure 3: mRNA expression level of olfactory receptor 157, identified in differential gene expression analysis of $\gamma\delta$ -T cells from day 18 embryos ($n=2$) and 6 week-old adult ($n=2$) mice ($n=2$), each including one population of cells from wild-type and CD3 double haploinsufficient mice. Each dot represents one population of cells, and the error bars representation the standard deviation.

Figure 4: mRNA expression level of olfactory receptor 1507, identified in differential gene expression analysis of $\gamma\delta$ -T cells ($n=3$) and DN3a thymocytes ($n=3$) from mice. Each dot represents a replicate population of cells from each cell type, and the error bars represent the standard deviation.

Figure 5: mRNA expression level of vomeronasal receptor genes identified in differential gene expression analysis of CD24⁺ CD73⁺ V γ 1.1⁺ $\gamma\delta$ -T cells ($n=3$) and CD24⁺ CD73⁺ V γ 2⁺ $\gamma\delta$ -T cells ($n=4$). Each dot represents one population of cells, and the error bars represent the standard deviation.