1 **Mechanistic Inferences from Clinical Reports of SARS-CoV-2**

2
3 Meagan M Jenkins^{1*}, Tyler R McCaw^{1*}, Paul A Goepfert²

4
5 ¹Department of Medicine, Division of Clinical Immunology & Rheumatology, University of
6 Alabama at Birmingham, Birmingham, AL, USA

7 ²Department of Medicine, Division of Infectious Diseases, University of Alabama at
8 Birmingham, Birmingham, AL, USA

9 *Authors contributed equally to this work

10
11 **Corresponding Author:**

12 Tyler R McCaw, PhD
13 Medical Scientist Training Program
14 University of Alabama at Birmingham
15 121 Shelby Building
16 1825 University Blvd
17 Birmingham, AL 35233
18 Email: trmccaw@uab.edu
19

Abstract

SARS-CoV-2 was identified as the causative pathogen in an outbreak of viral pneumonia cases originating in Wuhan, China, with an ensuing rapid global spread that led it to be declared a pandemic by the WHO on March 11, 2020. Given the threat to public health posed by sequelae of SARS-CoV-2 infection, the literature surrounding patient presentation in severe and non-severe cases, transmission rates and routes, management strategies, and initial clinical trial results have become available at an unprecedented pace. In this review we collate current clinical and immunologic reports, comparing these to reports of previous coronaviruses to identify mechanisms driving progression to severe disease in some patients. In brief, we propose a model wherein dysregulated type I interferon signaling leads to aberrant recruitment and accumulation of innate immune lineages in the lung, impairing establishment of productive adaptive responses, and permitting a pathologic pro-inflammatory state. Finally, we extend these findings to suggest possible treatment options that may merit investigation in randomized clinical trials.

Keywords: SARS-CoV-2, COVID-19, COVID-19 pneumonia, immune response, COVID-19 mechanisms

Introduction

SARS-CoV-2 was determined to be the causative pathogen of the disease COVID-19 resulting in an outbreak of viral pneumonia cases originating in Wuhan, China in 2019. The rapid spread of the virus on a global scale led to WHO designation as a pandemic on March 11, 2020. At the time of this review, there are over 3 million confirmed global cases of COVID-19 and at least 210,000 attributable deaths, with over 987,000 cases and 56,000 attributable deaths in the United States [1]. Relative to annual influenza infections, COVID-19 appears to have significantly higher rates of patient morbidity and mortality, though they remain lower than what was seen for SARS-CoV and MERS. Unfortunately, no specific therapy is available and the management of infected patients has been variable among institutions and nations. Given the rapid pace at which the global situation and our understanding of the virus is progressing, this review intends to collate currently available published case series (peer-reviewed and pre-print) and integrate these clinical findings with immunologic insights from COVID-19, SARS-CoV, and MERS studies.

Clinical characteristics among patients with COVID-19 infection

Emerging data from Wuhan, China suggests that while some patients are hospitalized with more severe COVID-19 infections (17%), most cases are not life-threatening [2]. Preliminary estimates suggest the time from infection to symptom onset to be around 5 days, and a mean duration of 11-12.5 days from the time of infection to hospitalization [3]. Despite initial controversy, viral shedding and transmission from person-to-person does occur during an asymptomatic period [4], likely contributing to the rapid global

spread. Indeed, most of the viral shedding occurs before symptom onset [5]. Of note, viral shedding has been found to align with cough symptoms, in nasopharyngeal samples and sputum. Shedding can also occur through stool although infectious viral transmission has not been reported through this route [6]. A notable limitation to most studies are the relatively small sample sizes and the relative exclusion of infected patients with only mild symptoms who do not seek medical attention. The severity of patient illness has repeatedly been found to correlate with symptomology, clinical parameters, and outcome; therefore, most published reports stratify patient parameters by "non-severe" versus "severe" disease (studies stratifying by "no ARDS" versus "ARDS", or "ICU care" versus "No ICU care" were conformed to this strategy for ease of reporting). Published case series indicate the most common presenting symptoms to include fever in the majority and cough with fatigue, sputum production, myalgias/arthralgias, and shortness of breath less commonly present but enriched in patients with severe disease [2,7–10]. Although distinct viral strains may predominate in different geographical regions, the clinical presentations from these Chinese studies were mirrored by laboratory-confirmed COVID-19 infected patients treated at an academic health system in New York, New York, USA [11,12], though respective frequencies of fever (61%) and cough (56%) were notably lower in an Italian case series [13] (**Table 1**).

Following hospital admission, clinical parameters tend to correlate with disease severity. The acute phase protein C-reactive protein (CRP) was significantly elevated in severe versus non-severe cases. Elevations in lactic acid dehydrogenase (LDH), aspartate aminotransferase (AST), and alanine aminotransferase (ALT) were also more frequent

and more significant in severe cases, suggesting a greater extent of systemic tissue damage and corroborating a more inflammatory environment in this cohort [2,7,9,14,15]. Immunologic perturbations are increasingly recognized as a defining feature of severe infection with COVID-19. Serum cytokine derangements include a nearly invariable elevation in interleukin 6 (IL-6). IL-8 and type I interferon are also frequently elevated in the blood of patients with severe COVID-19 pneumonia [16,17]. Although increased circulating TNF α has been reported in more severe cases, differences in concentration appear to be modest and inconsistent (**Table 1**).

Derangements in predominantly pro-inflammatory cytokines, or “cytokine storm” suggest propagation of a systemic inflammatory state in hospitalized patients that develop severe disease. Specifically, elevated ferritin and IL-6 were shown to correlate with mortality in Wuhan, China [9]. The neutrophil-to-lymphocyte ratio (NLR) is a surrogate of systemic inflammation [18] and can serve as a predictor of poor prognosis in critically-ill septic [19,20] and cancer patients [21]. Extending its utility to COVID-19 patients, NLR is significantly elevated in those with severe disease [8,15,22], serving as an independent risk factor and prognosticator of patients at high risk for progression to severe pneumonia [10]. Specifically, circulating neutrophil counts are significantly elevated while nearly all adaptive populations are negatively impacted by severe disease (**Table 1**).

Lessons from previous coronaviruses

With our understanding still in its infancy, mechanisms contributing to severe COVID-19 pneumonia are yet to be determined; however, lessons can be gleaned from the previous

coronavirus epidemics (SARS-CoV and MERS-CoV). Sequencing of the SARS-CoV-2 viral genome revealed a concordance rate of approximately 80% with SARS-CoV, its closest relative aside from bat CoVs (the likely source of COVID-19) [23,24]. Both coronaviruses also bind Angiotensin Converting Enzyme 2 (ACE2), a surface enzyme expressed predominantly on type 2 epithelial cells of the lower respiratory tract, to gain cellular entry, though SARS-CoV-2 binds with 10-20x greater affinity [25]. Clinically, patient presentation during the SARS-CoV epidemic in 2002-2003 bore striking similarities to that being reported during the current COVID-19 pandemic. Common symptoms of SARS-CoV included fever, chills, myalgia, and dry cough [26]. LDH and CK were frequently elevated in the context of neutrophilia and lymphopenia, again reflecting a systemic inflammatory state [26–29]. Here, a dysregulated immune response frequently resulted in ARDS, a pathologic outcome similar to COVID-19 [26], and similar findings on autopsy [30,31]. Given the substantial biochemical and clinical similarities, it is logical to assume similar mechanisms dictate the immune response to these pathogens. Therefore, lessons learned in the study of SARS-CoV can be cautiously applied to our evolving understanding of COVID-19.

Putative mechanism

SARS-CoV and SARS-CoV-2 enter cells in the respiratory tract via ACE2 and release damage associated molecular patterns (DAMPs; e.g. ATP, HMGB1, nucleic acids, etc.) as well as viral particle-derived pathogen associated molecular patterns (PAMPs) into the extracellular environment. Binding of these molecules to cognate pattern recognition receptors (PRRs) stimulates an innate immune response. In this manner, lung dendritic

cells recognize an infection, mature, and traffic to the draining lymph node wherein antigen is presented to T cells [32]. Stimulation of adaptive immunity then leads to viral clearance through cellular and humoral mechanisms--the likely scenario in asymptomatic patients, or with only mild disease. Progression to severe disease, however, is likely driven by dysregulation of this process.

Adaptive dysregulation

Levels of CD4 and CD8 T cells negatively correlate with disease severity in COVID-19 patients and are similarly decreased in SARS-CoV patients [27,29]. Demonstrating their central role in viral clearance, adoptive transfer of virus-specific CD4 or CD8 T cells significantly improved mortality and expedited viral clearance in a lethal challenge model of SARS-CoV. Moreover, vaccination with peptide-coated DCs one week prior to infection was able to elicit a protective CD8 T cell response [33]. In a different approach, Chen et al. depleted T cell subsets before infection and found CD4, but not CD8, T cells to be critical for efficient mouse clearance of SARS-CoV infection. In this same study, the administration of neutralizing antibodies following CD4 T cell depletion promoted viral clearance, suggesting a requirement for effective B cell help and production of neutralizing antibodies for viral control [34]. In line with these findings, antibodies to type A blood antigens appear to be cross-reactive and somewhat protective, as patients with type B and O blood are less frequently infected with SARS-CoV and SARS-CoV-2 [35,36]. However, declining numbers of circulating lymphocytes in severe disease seemingly suggests impairment of these responses.

In COVID-19-mediated lymphopenia, B cells, activated CD4 T cells, memory CD4 T cells, and CD8 T cells are reduced. One proposed explanation is that SARS-CoV-2 might directly infect T cells and initiate cell death by viral lysis [31]. This outcome seems unlikely, as Banerjee et al. found viral-like particles in CD4 T cells but demonstrated an absence of viral replication in healthy donor PBMCs of any lineage [37]. Furthermore, single-cell RNA-sequencing of PBMCs from hospitalized COVID-19 patients failed to find SARS-CoV-2 viral reads in any samples [16]. Although lymphopenia in the circulation could be driven by massive recruitment of these cells into the lungs, autopsy of patients having succumbed to severe COVID-19 pneumonia showed a paucity of infiltrating lymphocytes [31], rendering this an unlikely scenario as well. The systemic inflammatory state imposed by severe COVID-19 disease, much like sepsis, may then be the impetus behind observed lymphopenia and elevated NLR [38]. In sepsis, circulating lymphocytes display signs of early apoptosis, Annexin V surface expression and lymphocyte shrinkage [39], implicating loss of these populations through programmed cell death [40]. Thus, the systemic inflammatory state during severe COVID-19 pneumonia and/or viral sepsis likely induces lymphocyte apoptosis and dysregulated adaptive responses.

A recent report from China has found a positive correlation between abundance of SARS-CoV-2 Nucleoprotein (NP) neutralizing antibodies and disease severity, noting that earlier, stronger responders for NP specific anti-IgG and anti-IgM associate with increased disease severity. Conversely, patients with fewer circulating neutralizing antibodies were found to have a decreased viral load [6]. In agreement with this, Wu et al. reported about 30% of non-severe patients generated very low neutralizing antibody

titers against the spike (S) protein. It was also found that patients who were older with lower lymphocyte counts and increased CRP had increased neutralizing antibody titer, however, none of these patients had severe disease [41]. Although these are small observational studies, results are consistent with reports from MERS infection, where patients having succumbed to disease had robust neutralizing antibody responses during infection [42]. Similarly, in SARS-CoV infection, patients with severe illness had higher antibody titers at earlier stages during infection [43]. A macaque model of SARS-CoV proposed one explanation for an ostensible role of neutralizing antibodies in disease progression. Macaques vaccinated with S protein and challenged with SARS-CoV developed diffuse alveolar damage compared to only mild pathology in controls. Disease severity was a consequence of increased inflammatory monocyte-macrophages in the lungs and was mediated in part by Fc γ receptor signaling, as the combination of blocking Fc γ receptors and giving anti-S IgG decreased macrophage production of IL-8 and MCP-1 *in vitro* [44]. A second explanation is that viral S protein and N protein activate complement directly and/or in complex with neutralizing antibodies, contributing to the widespread complement activation that culminates in characteristic tissue damage and diffuse microthrombi [45,46]. Activated complement protein C5a is a potent anaphylatoxin that enhances Fc γ RIII signaling upon receptor binding (C5aR) to promote alveolar macrophage activation [47], potentially serving as a link between these two hypotheses. Indeed, this scenario accounts for observed elevations in D-dimer, reduced platelets, and diffuse tissue thrombi found on autopsy [48].

Antibody-mediated enhancement of viral disease—an increase in viral uptake, replication, and/or pathogenicity due to the binding of virions to antibodies—cannot be excluded. This mechanism has been demonstrated in various other respiratory infections, including influenza virus, respiratory syncytial virus, and is controversial in SARS-CoV and MERS [49].

It is important to consider, however, that these clinical findings are subject to a temporal sampling bias in that patients do not typically seek medical attention until symptoms are relatively progressed. Therefore, it is difficult to discern whether antibody production contributed to a dysregulated immune response and disease severity, or whether disease severity and an altered cytokine milieu drove the production of higher antibody titers. In the latter case, neutralizing antibodies could still be protective if present early after infection and would help explain the potential efficacy of passive immunization. Indeed, Hoffmann et al. found SARS-CoV neutralizing antibodies directed against the viral S protein to be cross-protective, inhibiting both SARS-CoV and SARS-CoV-2 host cell entry *in vitro* [50].

Innate Dysregulation

Frequent and significant elevations in circulating IL-6 and IL-8, which contribute to neutrophil activation and recruitment, suggest a potential pathologic role for neutrophils in patients developing severe COVID-19 pneumonia. Using a rat model of SARS-CoV, Haick et al. showed a critical contribution of neutrophils to the initial anti-viral immune response, elaborating IL-1, TNF α , CXCL1, CXCL3, CCL2, and IP10 to orchestrate the

recruitment of a productive cellular response to the lungs. Expectedly, depletion of neutrophils prior to infection impaired viral clearance. However, subsequent repopulation and influx of neutrophils to the lungs in late-stage disease contributed to significant morbidity and mortality without improvement in viral clearance [51]. Bulk RNA-sequencing and pathway analysis of PBMCs from COVID-19 patients with mild versus severe symptoms revealed upregulation of neutrophil chemotaxis, neutrophil activation, and type I IFN signaling pathways, consistent with these mouse studies. Virus-host interactome analysis then indicated that non-structural proteins 9 and 10 impaired the function of NKRF (a competitive inhibitor of NF κ B) to induce IL-6 and IL-8 expression in lung epithelia, which the authors postulate could incite tissue destruction and eventual ARDS [52].

A role for type I IFN signaling

What is driving this perturbed immune response that causes significantly increased rates of mortality in the elderly and those with chronic disease? Accumulating evidence suggests type I interferon signaling, or dysregulation thereof, dictates establishment of a productive immune response to then preempt progressive viral replication, development of severe pneumonia, and ARDS. Channappanavar et al. showed that delayed type I interferon signaling promotes lethal respiratory infection with SARS-CoV in mice, while supplementation with IFN β prior to peak viral replication abrogated mortality with no significant difference in pulmonary viral load. Knockout of the interferon alpha receptor led to reduced expression of CCL2 (among other chemokines), which functions to recruit inflammatory monocyte-macrophages via CCR2. Antibody-mediated depletion of this

population led to reduced levels of $\text{TNF}\alpha$, $\text{IL-1}\beta$, IL-6 , and iNOS and ameliorated disease severity, directly linking pulmonary macrophages to morbidity and mortality in SARS-CoV [53]. Extending these findings, the late administration of type I interferon led to recruitment of pathogenic neutrophils and macrophages to the lung and development of fatal pneumonia in a mouse model of MERS-CoV [54]. Single-cell RNA-sequencing of PBMCs from four COVID-19 patients at discrete stages of disease severity (pre-ICU, ICU, and post-ICU samples) also identified a relationship between dysregulated type I interferon signaling in PBMCs and clinical status. This correlated with drastically increased circulating inflammatory monocytes and fewer T and B cells while patients were in the intensive care unit [16]. In addition, an autopsy of two patients having succumbed to COVID-19 infection showed significant aggregation and activation of infiltrating macrophages--homogeneously expressing IL-6 , IL-10 , and $\text{TNF}\alpha$ --amidst extensive tissue destruction, corroborating a direct role for these cells in pulmonary pathology [31].

Failure to clear viral infection early might then lead to prolonged type I signaling that can instead impair adaptive immune functions [55]. In support of this notion, flow cytometry on PBMCs from COVID-19 patients with mild versus severe disease found a significant decline in the frequency of multifunctional CD4 T cells in the severe cohort alongside the increased expression of activation and exhaustion surface proteins on CD8 T cells in the total T cell pool. The authors compare these phenotypic changes to an exhaustion-like state [56]. Indeed, prolonged type I interferon signaling is known to reinforce CD8 T cell exhaustion and impair viral control [57]. The extent to which the development of functional T cell exhaustion (that is, loss of cytokine production and cytotoxicity) contributes to

disease severity on the timescale of COVID-19 infection and resolution remains to be determined.

Collectively, these studies suggest a model wherein a delayed initial immune response to infection, including type I interferon signaling, permits high levels of viral replication and persistence. Neutrophil recruitment to the lungs and subsequent activation then elaborates inflammatory cytokines (IL-6 and IL-8) and chemokines (notably CCL2), driving substantial monocyte recruitment and activation. It is likely a combination of these innate populations that mediate eventual tissue destruction manifested by severe disease. The systemic inflammatory state also hinders adaptive immunity, inducing lymphocyte apoptosis, and reducing the production of canonical T cell cytokines as well as neutralizing antibodies, impairing viral clearance. These factors culminate in the cytokine storm with local and systemic tissue destruction observed in patients with severe COVID-19 pneumonia. Delayed and/or blunted immune responses characteristic in the elderly [58] and an already dysregulated inflammatory state present in common comorbidities (e.g. insulin resistance, hypertension) [59,60] compromise prompt responses to infection, potentially driving the significantly increased morbidity and mortality observed in these patient populations (**Figure 1**).

Immunological Memory to COVID-19

Whether or not protective memory responses are established following infection with COVID-19 remains uncertain. Convalescent plasma transfer from recovered to currently infected patients has shown promise in reducing clinical severity and expediting clearance

of disease [61,62], implying that antibodies to at least some viral epitopes can confer protection. Because the kinetics of B cell contraction in COVID-19 are unknown, it could be that memory cells persist for some time following infection resolution, or that antibodies produced by antibody secreting cells prior to contraction are predominantly responsible for observed benefits. Importantly, if activation induced cell death is occurring in the T lymphocytes, which has been shown in CD44-positive memory or antigen experienced cells, this mechanism could also impede memory cell formation, either by depleting memory B or T cells directly or by interfering with T-B cell help [63].

It appears that memory B cell responses in SARS-CoV are relatively short lived. Patients having recovered from SARS-CoV maintained neutralizing antibodies for 16 months after infection, tending to peak 4 months after infection and decreasing thereafter [64]. After six years, anti-SARS antibodies were undetectable in 91% of patients and no SARS specific memory B cells were found. Memory T cells were, however, detectable 6 years after infection, but their ability to confer protective immunity remains uncertain [65]. If SARS-CoV-2 proves to behave similarly, it will be important to determine the cause of this temporary persistence and approaches to address it.

Implications for treatment strategies

As of April 27th, the NIH shows almost 950 clinical trials registered in association with COVID-19 (search term “COVID-19” in clinicaltrials.gov). In addition, new papers are becoming available on preprint servers daily, giving insight into new therapy options and preliminary results on treatment options. Of note, the SARS-CoV-2 viral and protease

structure, which was found to be highly conserved throughout CoVs, has been identified, and specific protease inhibitors with antiviral effects *in vitro* are currently under investigation [66,67]. The development of specific anti-viral small molecule inhibitors along with a vaccine capable of generating protective immunity is undoubtedly critical to managing the COVID-19 pandemic.

Until such targeted therapies are available, potential points of clinical intervention for patients at high risk of progression can be surmised from the available literature. Administration of type I interferon quickly after infection may be sufficient to expedite a productive immune response. Unfortunately, this is not likely to be a clinically viable approach given the brief window of opportunity between infection and intervention, well before the onset of symptoms. Interrupting early stages of the pathologic response might curtail subsequent immune derangements and facilitate viral clearance. To this end, Li et al. proposed to inhibit neutrophil production of IL-6 and IL-8 using small molecule JAK inhibitors or IL-6 neutralizing antibodies [52]. Because anti-IL-6 has proven efficacious in the alleviation of chimeric antigen receptor T cell therapy cytokine release syndrome [68], it is possibly a useful adjuvant to any treatment strategy for severe COVID-19 pneumonia. Alternatively, blocking CCL2-mediated recruitment of monocytes to the lung with neutralizing antibodies may be efficacious over a longer period and more closely correspond with the development of symptoms (given the role of macrophages in lung pathology). Of course, a patient's clinical picture should be evaluated when considering administration of broadly immunosuppressive agents. Giving these during a critical time in the initial stages of response, which seem to determine whether severe disease will

occur, might be detrimental to patient outcome. In these initial stages blocking monocyte recruitment, for example, may prove more advantageous.

For patients that do not come to clinical attention until pathologic inflammation has been established, or for whom clinical progression is imminent, management should focus on targeted immunosuppression and conferral of passive immunity. Early management of such patients has included broad-spectrum anti-virals, prophylactic antibiotics, and rather indiscriminate immunosuppression with corticosteroids [69]. More selective immune suppression could be achieved by specifically targeting neutrophils or macrophages. The depletion of these populations responsible for lung pathology might help to restore balance and promote an effective anti-viral immune response. Additionally, blocking type I interferons with neutralizing antibodies in advanced disease may delimit lung pathology [70].

Once pulmonary pathology becomes sufficiently advanced, however, the ability to stimulate a productive immune response is compromised and passive immunization with virus-neutralizing antibodies might be of greatest benefit. This approach recently demonstrated favorable outcomes in two of three patients, while the failed case seems to have been an error in cross-matching [61]. A second study with 10 patients reported similarly auspicious outcomes [62]. According to the FDA's Recommendations for investigational COVID-19 convalescent plasma, plasma can be used in life-threatening COVID-19 cases in the United States, but trial results are still pending [71]. While seemingly promising, plasma transfer will likely be unable to reverse lung damage from

immune dysregulation; therefore, the goal must remain improvement of screening measures and early identification of patients at high risk of severe pneumonia. Furthermore, these and other potentially beneficial therapies need to be studied in randomized control studies in sufficient numbers of patients to accurately assess their utility.

Concluding Remarks

Our understanding of biologic parameters governing patient presentation following infection with SARS-CoV-2 is expanding at an extraordinary pace. Here we have integrated clinical data from mild and severe COVID-19 patients with mechanistic findings from similar coronaviruses to propose a model wherein an aberrant initial innate immune response cascades into dysregulation of adaptive immunity, preventing efficient viral clearance. Exploring this possible mechanism for severe COVID-19 pathogenesis identified several treatment options meriting continued investigation.

Funding

This work was supported by funds from the UAB School of Medicine Dean's Office.

Acknowledgments

The authors would like to thank Dr. Jeremy Walker for his clinical input reviewing this manuscript. The authors also thank the UAB School of Medicine Dean's Office for supporting this work.

384 **Conflict of Interest**

385 The authors declare no potential conflicts of interest.

386

387

References

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis.* **2020**; 3099(20):19–20.
2. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* Lancet Publishing Group; **2020**; 395(10223):497–506.
3. Li X, Geng M, Peng Y, Meng L, Lu S. Molecular immune pathogenesis and diagnosis of COVID-19. *J. Pharm. Anal.* Xi'an Jiaotong University; 2020.
4. Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-NCOV infection from an asymptomatic contact in Germany. *N Engl J Med.* **2020**; 382(10):970–971.
5. He X, Lau EH, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *medRxiv [Internet].* **2020**; :2020.03.15.20036707. Available from: <https://www.medrxiv.org/content/10.1101/2020.03.15.20036707v2>
6. Tan W, Lu Y, Zhang J, et al. Viral Kinetics and Antibody Responses in Patients with COVID-19. *medRxiv [Internet].* **2020**; :2020.03.24.20042382. Available from: <http://medrxiv.org/content/early/2020/03/26/2020.03.24.20042382.abstract>
7. Guan W, Ni Z, Hu Y, et al. Clinical characteristics of 2019 novel coronavirus infection in China. *N Engl J Med.* **2020**; .
8. Qin C, Zhou L, Hu Z, et al. Dysregulation of Immune Response in Patients with COVID-19 in Wuhan, China. *SSRN Electron J [Internet].* **2020**; . Available from: <https://academic.oup.com/cid/advance-article-abstract/doi/10.1093/cid/ciaa248/5803306>

- 411 9. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult
412 inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*.
413 Lancet Publishing Group; **2020**; .
- 414 10. Liu J, Liu Y, Xiang P, et al. Neutrophil-to-Lymphocyte Ratio Predicts Severe
415 Illness Patients with 2019 Novel Coronavirus in the Early Stage. *medRxiv*
416 [Internet]. **2020**; 807:2020.02.10.20021584. Available from:
417 <https://doi.org/10.1101/2020.02.10.20021584>
- 418 11. Petrilli CM, Jones SA, Yang J, et al. Title: Factors associated with hospitalization
419 and critical illness among 4,103 patients with. *medRxiv* [Internet]. **2020**; .
420 Available from: <https://doi.org/10.1101/2020.04.08.20057794>
- 421 12. Goyal P, Choi J, Pinheiro L, Schenck E, Al. E. Clinical Characteristics of Covid-19
422 in New York City. *N Engl J Med*. **2020**; .
- 423 13. Caruso D, Zerunian M, Polici M, et al. Chest CT Features of COVID-19 in Rome,
424 Italy. *Radiology*. **2020**; :201237.
- 425 14. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to
426 COVID-19 based on an analysis of data of 150 patients from Wuhan, China.
427 *Intensive Care Med*. **2020**; .
- 428 15. Wu C, Chen X, Cai Y, et al. Risk Factors Associated with Acute Respiratory
429 Distress Syndrome and Death in Patients with Coronavirus Disease 2019
430 Pneumonia in Wuhan, China. *JAMA Intern Med*. **2020**; .
- 431 16. Wei L, Ming S, Zou B, et al. Viral Invasion and Type I Interferon Response
432 Characterize the Immunophenotypes during COVID-19 Infection. *SSRN Electron*
433 *J*. **2020**; .

- 434 17. Wan S, Yi Q, Fan S, et al. Characteristics of lymphocyte subsets and cytokines in
435 peripheral blood of 123 hospitalized patients with 2019 novel coronavirus
436 pneumonia (NCP). medRxiv. **2020**; .
- 437 18. Zahorec R. Ratio of neutrophil to lymphocyte counts — rapid and simple
438 parameter of systemic inflammation and stress in critically ill. Bratisl Lek List.
439 **2001**; 102(1):5–14.
- 440 19. Cataudella E, Giraffa CM, Marca S Di, et al. Neutrophil-To-Lymphocyte Ratio: An
441 Emerging Marker Predicting Prognosis in Elderly Adults with Community-Acquired
442 Pneumonia. J Am Geriatr Soc. **2017**; 65(8):1796–1801.
- 443 20. Hwang SY, Shin TG, Jo IJ, et al. Neutrophil-to-lymphocyte ratio as a prognostic
444 marker in critically-ill septic patients. Am J Emerg Med [Internet]. Elsevier; **2017**;
445 35(2):234–239. Available from: <https://doi.org/10.1016/j.ajem.2016.10.055>
- 446 21. Templeton AJ, McNamara MG, Šeruga B, et al. Prognostic role of neutrophil-to-
447 lymphocyte ratio in solid tumors: A systematic review and meta-analysis. J Natl
448 Cancer Inst. **2014**; 106(6).
- 449 22. Zhang B, Zhou X, Zhu C, et al. Immune phenotyping based on neutrophil-to-
450 lymphocyte ratio and IgG predicts disease severity and outcome for patients with
451 COVID-19. medRxiv [Internet]. **2020**; :2020.03.12.20035048. Available from:
452 <https://doi.org/10.1101/2020.03.12.20035048>
- 453 23. Zhou P, Yang X-L, Wang X-G, et al. Discovery of a novel coronavirus associated
454 with the recent pneumonia outbreak in humans and its potential bat origin. Nature.
455 **2020**; .
- 456 24. Boni MF, Lemey P, Jiang X, et al. Evolutionary origins of the SARS-CoV-2

sarbecovirus lineage responsible for the COVID-19 pandemic. bioRxiv [Internet].
2020; . Available from: <https://doi.org/10.1101/2020.03.30.015008>

25. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike
in the prefusion conformation. Science (80-) [Internet]. **2020**; 367(6483):1260–
1263. Available from: <http://science.sciencemag.org/>

26. Lee HKK, Tso EYK, Chau TN, Tsang OTY. Asymptomatic Severe Acute
Respiratory Syndrome-associated Coronavirus Infection. Emerg Infect Dis
[Internet]. **2003**; . Available from: www.cdc.gov/eid

27. He Z, Zhao C, Dong Q, et al. Effects of severe acute respiratory syndrome
(SARS) coronavirus infection on peripheral blood lymphocytes and their subsets.
Int J Infect Dis. **2005**; 9(6):323–330.

28. Booth CM, Matukas LM, Tomlinson GA, et al. Clinical Features and Short-term
Outcomes of 144 Patients With SARS in the Greater Toronto Area. JAMA
[Internet]. **2003**; 289(21):2801–2809. Available from:
<https://doi.org/10.1001/jama.289.21.JOC30885>

29. Li T, Qiu Z, Zhang L, et al. Significant Changes of Peripheral T Lymphocyte
Subsets in Patients with Severe Acute Respiratory Syndrome. J Infect Dis
[Internet]. **2004**; 189(4):648–651. Available from:
<https://academic.oup.com/jid/article-abstract/189/4/648/839067>

30. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with
acute respiratory distress syndrome. Lancet Respir Med. Lancet Publishing
Group; **2020**; 8(4):420–422.

31. Wang C, Xie J, Zhao L, et al. Alveolar macrophage activation and cytokine storm

480 in the pathogenesis of severe COVID-19.pdf. Res Sq. **2020**; .

481 32. Lau YL, Peiris JSM, Law HKW. Role of dendritic cells in SARS coronavirus
482 infection. Hong Kong Med J. **2012**; 18(4):S28–S30.

483 33. Zhao J, Zhao J, Perlman S. T Cell Responses Are Required for Protection from
484 Clinical Disease and for Virus Clearance in Severe Acute Respiratory Syndrome
485 Coronavirus-Infected Mice. J Virol. **2010**; 84(18):9318–9325.

486 34. Chen J, Lau YF, Lamirande EW, et al. Cellular Immune Responses to Severe
487 Acute Respiratory Syndrome Coronavirus (SARS-CoV) Infection in Senescent
488 BALB/c Mice: CD4+ T Cells Are Important in Control of SARS-CoV Infection. J
489 Virol. American Society for Microbiology; **2010**; 84(3):1289–1301.

490 35. Cheng Y, Cheng G, Chui CH, et al. ABO blood group and susceptibility to severe
491 acute respiratory syndrome [5]. J Am Med Assoc. **2005**; 293(12):1450–1451.

492 36. Zhao J, Yang Y, Huang H, et al. Relationship between the ABO Blood Group and
493 the COVID-19 Susceptibility. medRxiv [Internet]. **2020**; :2020.03.11.20031096.
494 Available from: <https://doi.org/10.1101/2020.03.11.20031096>

495 37. Banerjee A, Nasir JA, Budylowski P, et al. Isolation, sequence, infectivity and
496 replication kinetics of SARS-CoV-2 1. bioRxiv [Internet]. **2020**; . Available from:
497 <https://doi.org/10.1101/2020.04.11.037382>

498 38. Wesche DE, Lomas-Neira JL, Perl M, Chung C-S, Ayala A. Leukocyte apoptosis
499 and its significance in sepsis and shock. J Leukoc Biol. **2005**; 78(2):325–337.

500 39. Schroeder S, Lindemann C, Decker D, et al. Increased susceptibility to apoptosis
501 in circulating lymphocytes of critically ill patients. Langenbeck's Arch Surg. **2001**;
502 386(1):42–46.

40. Roth G, Moser B, Krenn C, et al. Susceptibility to programmed cell death in T-lymphocytes from septic patients: A mechanism for lymphopenia and Th2 predominance. *Biochem Biophys Res Commun.* **2003**; 308(4):840–846.
41. Wu F, Wang A, Liu M, et al. Neutralizing antibody responses to SARS-CoV-2 in a COVID-19 recovered patient cohort and their implications. *medRxiv [Internet].* **2020**; :2020.03.30.20047365. Available from: <https://www.medrxiv.org/content/10.1101/2020.03.30.20047365v1>
42. Al-Abdely HM, Midgley CM, Alkhamis AM, et al. Middle east respiratory syndrome coronavirus infection dynamics and antibody responses among clinically diverse patients, Saudi Arabia. *Emerg Infect Dis.* **2019**; 25(4):753–766.
43. Ho MS, Chen WJ, Chen HY, et al. Neutralizing antibody response and SARS severity. *Emerg Infect Dis.* **2005**; 11(11):1730–1737.
44. Liu L, Wei Q, Lin Q, et al. Anti-spike IgG causes severe acute lung injury by skewing macrophage responses during acute SARS-CoV infection. *JCI insight.* **2019**; 4(4):1–19.
45. Gao T, Hu M, Zhang X, et al. Highly pathogenic coronavirus N protein aggravates lung injury by MASP-2-mediated complement over-activation. *medRxiv.* **2020**; :2020.03.29.20041962.
46. Magro C, Mulvey JJ, Berlin D, et al. Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID- 19 . The COVID-19 resource centre is hosted on Elsevier Connect , the company ’ s public news and information website . Elsevier hereby grants permission to make all its COVID-19-related research that is

available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories , such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source . These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active . **2020**; (January).

47. Shushakova N, Skokowa J, Schulman J, et al. C5a anaphylatoxin is a major regulator of activating versus inhibitory FcγRs in immune complex-induced lung disease. *J Clin Invest.* **2002**; 110(12):1823–1830.
48. Fox SE, Akmatbekov A, Harbert JL, Li G, Brown JQ. Pulmonary and Cardiac Pathology in Covid-19 : The First Autopsy Series from New Orleans 1) Department of Pathology , LSU Health Sciences Center , New Orleans 2) Pathology and Laboratory Medicine Service , Southeast Louisiana Veterans Healthcare System 3. *medRxiv.* **2020**; .
49. Smatti MK, Thani AA Al, Yassine HM. Viral-induced enhanced disease illness. *Front Microbiol.* **2018**; 9(DEC).
50. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell.* **2020**; :271–280.
51. Haick AK, Rzepka JP, Brandon E, Balemba OB, Miura TA. Neutrophils are needed for an effective immune response against pulmonary rat coronavirus infection, but also contribute to pathology. *J Gen Virol.* **2014**; 95(PART3):578–590.

52. Li J, Guo M, Tian X, et al. Virus-host interactome and proteomic survey of PMBCs from COVID-19 patients reveal potential virulence factors influencing SARS-CoV-2 pathogenesis. *bioRxiv*. **2020**; :2020.03.31.019216.
53. Channappanavar R, Fehr AR, Vijay R, et al. Dysregulated Type I Interferon and Inflammatory Monocyte-Macrophage Responses Cause Lethal Pneumonia in SARS-CoV-Infected Mice. *Cell Host Microbe*. Cell Press; **2016**; 19(2):181–193.
54. Channappanavar R, Fehr AR, Zheng J, et al. IFN-I response timing relative to virus replication determines MERS coronavirus infection outcomes. *J Clin Invest*. **2019**; 129(9):3625–3639.
55. Ivashkiv LB, Donlin LT. Regulation of type i interferon responses. *Nat Rev Immunol*. Nature Publishing Group; **2014**; 14(1):36–49.
56. Zheng HY, Zhang M, Yang CX, et al. Elevated exhaustion levels and reduced functional diversity of T cells in peripheral blood may predict severe progression in COVID-19 patients. *Cell Mol Immunol*. **2020**; (March).
57. Wu T, Ji Y, Ashley Moseman E, et al. The TCF1-Bcl6 axis counteracts type I interferon to repress exhaustion and maintain T cell stemness. *Sci Immunol*. **2016**; 1(6):1–27.
58. Brubaker AL, Palmer JL, Kovacs EJ. Age-related dysregulation of inflammation and innate immunity: Lessons learned from rodent models. *Aging Dis*. **2011**; 2(5):346–360.
59. Smith AG, Sheridan PA, Harp JB, Beck MA. Diet-Induced Obese Mice Have Increased Mortality and Altered Immune Responses When Infected with Influenza Virus. *J Nutr*. **2007**; 137(5):1236–1243.

- 572 60. Galkina E, Ley K. Immune and Inflammatory Mechanisms of Atherosclerosis.
573 Annu Rev Immunol. **2009**; 27(1):165–197.
- 574 61. Pei S, Yuan X, Zhang Z, et al. Convalescent Plasma to Treat COVID-19: Chinese
575 Strategy and Experiences. medRxiv [Internet]. **2020**; . Available from:
576 <https://doi.org/10.1101/2020.04.07.20056440>
- 577 62. Duan K, Liu B, Li C, et al. The feasibility of convalescent plasma therapy in severe
578 COVID-19 patients: a pilot study. medRxiv [Internet]. **2020**;
579 (8):2020.03.16.20036145. Available from:
580 <https://doi.org/10.1101/2020.03.16.20036145>
- 581 63. Zarozinski CC, McNally JM, Lohman BL, Daniels KA, Welsh RM. Bystander
582 Sensitization to Activation-Induced Cell Death as a Mechanism of Virus-Induced
583 Immune Suppression. J Virol. **2000**; 74(8):3650–3658.
- 584 64. Cao WC, Liu W, Zhang PH, Zhang F, Richardus JH. Disappearance of antibodies
585 to SARS-associated coronavirus after recovery [18]. N Engl J Med [Internet].
586 **2007**; 357(11):1162–1163. Available from: <http://authors.nejm.org>
- 587 65. Tang F, Quan Y, Xin Z-T, et al. Lack of Peripheral Memory B Cell Responses in
588 Recovered Patients with Severe Acute Respiratory Syndrome: A Six-Year Follow-
589 Up Study. J Immunol. The American Association of Immunologists; **2011**;
590 186(12):7264–7268.
- 591 66. Bestle, Dorothea et. al. TMPRSS2 and furin are both essential for proteolytic
592 activation and spread of SARS-CoV-2 in human airway epithelial cells and
593 provide promising drug targets. Available from:
594 <https://doi.org/10.1101/2020.04.15.042085>

67. Jin Z, Du X, Xu Y, et al. Structure of Mpro from COVID-19 virus and discovery of its inhibitors. *bioRxiv* [Internet]. **2020**; :2020.02.26.964882. Available from: <http://www.nature.com/articles/s41586-020-2223-y>
68. Le RQ, Li L, Yuan W, et al. FDA Approval Summary: Tocilizumab for Treatment of Chimeric Antigen Receptor T Cell-Induced Severe or Life-Threatening Cytokine Release Syndrome. *Oncologist*. **2018**; 23(8):943–947.
69. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. **2020**; 395(10223):497–506.
70. Sallard E, Lescure F-X, Yazdanpanah Y, et al. Type 1 interferons as a potential treatment against COVID-19. *Antiviral Res* [Internet]. **2020**; (January):104791. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0166354220302059>
71. FDA. Recommendations for Investigational COVID-19 Convalescent Plasma [Internet]. 2020. Available from: <https://www.fda.gov/vaccines-blood-biologics/investigational-new-drug-ind-or-device-exemption-ide-process-cber/recommendations-investigational-covid-19-convalescent-plasma#Pathways> for

Table 1: Compiled clinical data from multiple studies of severe and non-severe COVID-19 patients. Data is compiled based on information provided in respective studies. Data is presented as mean (range) when applicable, only mean is presented if data is from a single study. If data conforming to the table format could not be extracted from a report then that data was excluded from the table. Abbreviations: shortness of breath (SOB), body mass index (BMI), chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), interleukin (IL), tumor necrosis factor (TNF), interferon type I (IFN), C-reactive protein (CRP), lactate dehydrogenase (LDH), aspartate transaminase (AST), alanine transaminase (ALT), neutrophil-to-lymphocyte ratio (NLR), regulatory T cell (Treg).

Symptoms (%)	Non-severe	Severe	Reference no.
Fever	91.94 (83-97.7)	94.29 (87-100)	2, 7, 8, 9, 10, 14, 15
Cough	66.27 (33.7-82)	71.36 (33.6-85)	2, 7, 8, 9, 10, 14, 15
Sputum	31.61 (33.6-85)	42.59 (26-64.7)	2, 7, 8, 9, 10, 14, 15
Fatigue	35.54 (21-52.3)	42.57 (22-70.6)	2, 7, 8, 9, 10, 14, 15
SOB	30.18 (9.1-62)	52.02 (17.6-87)	7, 8, 10, 14, 15
Myalgia/arthralgia	16.23 (14.4-19.3)	18.47 (15-23.1)	7, 8, 9
Diarrhea	18.95 (11.4-26.5)	16.6 (5.9-27.3)	8, 10
Comorbidity (%)	Non-severe	Severe	Reference no.
Male	59.10 (47.7-68)	71 (58.8-85)	2, 10, 11, 12, 14, 15
Obesity (BMI 30-40)	32.7 (31.9-33.5)	37.85 (32.3-43.4)	11, 12
Chronic conditions	70.70	73.40	11
Cardiovascular conditions	9.57 (0-42)	18.23 (5.9-47.1)	2, 8, 10, 11, 14, 15,
Diabetes	15.82 (4.5-22.9)	19.41 (8-27.7)	2, 8, 10, 11, 12, 14, 15
Asthma/COPD	5.5 (0-12.2)	10.56 (3-17.6)	2, 10, 11, 12, 14
Hypertension	22.62 (13.6-48.3)	35.2 (15-53.8)	2, 8, 10, 12, 14, 15
CKD	2.40	2.10	8
Cytokines (pg/mL)	Non-severe	Severe	Reference no.
IL-1b	5.00	5.00	8
IL-6	9.22 (6.3-13.41)	18.55 (7.39-37.7)	8, 9, 14, 15, 17
IL-8	13.70	18.40	8
IL-10	3.73 (2.4-5)	5.6 (4.59-6.6)	8, 17
TNFa	6.24 (4.1-8.4)	5.82 (2.9-8.7)	8, 17
Labs	Non-severe	Severe	Reference no.
CRP (mg/L)	22.6 (10.3-34.1)	77.7 (23.5-126.6)	10, 14, 15
LDH (U/L)	263.83 (253.5-281)	439 (396-521)	2, 9, 15
AST (U/L)	32 (30-34)	41 (38-44)	2, 15
ALT (U/L)	27 (27-27)	41.33 (35-49)	2, 9, 15
D-dimer (µg/mL)	0.56 (0.52-0.6)	3.18 (1.16-5.2)	9, 15
NLR	2.74 (2.2-3.2)	6.53 (3.6-10.5)	8, 10, 15
Leukocytes (*10 ⁹ /L)	4.8 (4.3-5.2)	6 (3.7-9.8)	7, 9, 10
Lymphocytes (*10 ⁹ /L)	1.05 (1-1.1)	0.68 (0.4-0.9)	2, 7, 9, 10
Platelets (*10 ⁹ /L)	177.13 (149-220)	163 (137.5-196)	2, 7, 9, 10
Neutrophils (*10 ⁹ /L)	2.89 (2.4-3.2)	4.71 (2.8-7.04)	8, 10, 15
Monocytes (*10 ⁹ /L)	0.37 (0.34-0.4)	0.35 (0.29-0.4)	8, 15
B cells (*10 ⁶ /L)	181.05 (166-196.1)	147.15 (125.3-169)	8, 17
Total T cells (*10 ⁶ /L)	648.4 (633-663.8)	454.05 (446.5-461.6)	8, 15
CD4 T cells (*10 ⁶ /L)	414.27 (371-451.3)	260.77 (234-285.1)	8, 15, 17
Naive CD4 T cells (*10 ⁶ /L)	35.00	44.50	8
Memory CD4 T cells (*10 ⁶ /L)	65.00	55.50	8
CD8 T cells (*10 ⁶ /L)	243.83 (201.9-288.6)	163.73 (154.7-179)	8, 15, 17
Treg (*10 ⁶ /L)	4.50	3.70	8

Figure 1: Proposed model of COVID-19 immune dysregulation leading to severe disease. Schematic of the proposed mechanism responsible for severe COVID-19 disease in some patients. Delayed type I IFN production by dendritic cells (DCs), or other IFN-producing populations, instigates a defective response that cascades into other responding immune lineages. This delay allows increased recruitment of neutrophils, secreting inflammatory cytokines and chemokines, then driving the accumulation of inflammatory monocyte-macrophages in the lungs, particularly the alveoli. Skewing of the innate response creates a potent inflammatory cytokine environment (including IL-6, IL-8, TNF α , reactive oxygen species (ROS), etc.) acting to propagate pulmonary pathology. Additionally, the systemic inflammatory state leads to lymphopenia and stymies establishment of a productive adaptive immune response.