TLR10 is a differentially expressed gene in lymph node metastatic human breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³; the lymph nodes reside between the breast and the brain. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with lymph node metastasis in humans with breast cancer. We found that toll-like receptor 10, encoded by TLR10, was among the genes whose expression was most different in the lymph node metastases of patients with metastatic, or “regional” breast cancer as compared to primary tumors of the breast. TLR10 mRNA was present at increased quantities in lymph node metastatic tissues as compared to primary tumors of the breast. Importantly, expression of TLR10 in primary tumors was significantly correlated with patient recurrence-free survival. Modulation of TLR10 expression may be relevant to the biology by which tumor cells metastasize from the breast to the lymph nodes in humans with metastatic breast cancer.

Keywords: breast cancer, metastasis, lymph node metastases, central nervous system metastases, toll-like receptor 10, TLR10, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. The lymph nodes reside between the breast and the brain. We performed a global comparative analysis of primary and metastatic tumors in patients with lymph node metastatic breast cancer^{4,5}. We discovered differential and increased expression of the gene encoding toll-like receptor 10, TLR10, in lymph node metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE42568⁵ for this global differential gene expression analysis of lymph node metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=71$ primary breast tumors and $n=7$ lymph node metastases from patients with breast cancer; analysis was performed using platform GPL887. GSE42568 was generated using Affymetrix Human Genome U133 Plus 2.0 array technology with $n=17$ normal breast tissue biopsies and $n=104$ primary breast tumor biopsies from patients with breast cancer; analysis was performed using platform GPL570. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether TLR10 gene expression was significantly different between primary tumors of the breast and lymph node metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of TLR10 mRNA expression levels with recurrence-free survival in $n=1764$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic and primary tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of lymph node metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

TLR10 is differentially expressed in the lymph node metastases of patients with lymph node metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified toll-like receptor 10, encoded by TLR10, as a differentially expressed gene in the breast metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes expressed in lymph node

metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, TLR10 ranked 117 out of 18034 total transcripts (Chart 1), equating to 99.4% differential expression. Differential expression of TLR10 in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=1.07E-03$).

To validate transcriptome-wide differential expression of TLR10 in human breast cancer, we queried a second microarray dataset⁵, here comparing normal breast tissues and primary tumors of the breast. Again, we identified TLR10 as a differentially expressed gene in the lymph node metastatic tissues of patients with breast cancer (Chart 2). When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, TLR10 ranked 19617 out of 54675 total transcripts (Chart 2), equating to 64.1% differential expression. Differential expression of TLR10 in the primary tumors of patients with breast cancer was statistically significant (Chart 2; $p=3.48E-02$). Thus, differential expression of TLR10, transcriptome-wide, in the tumor tissues of women with breast cancer was conserved across two independent microarray datasets, both in primary tumors and in lymph node metastases.

TLR10 is expressed at higher levels in the lymph node metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for TLR10, in primary tumors of the breast and in lymph node metastasis of patients with lymph node metastatic breast cancer to determine direction and statistical significance of change in TLR10 expression in lymph node metastatic tissues. TLR10 was expressed at higher levels in the lymph node metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 1; $p=0.001055$).

TLR10 expression is significantly correlated with survival outcomes in human breast cancer.

We performed Kaplan-Meier survival analysis⁸ in 1764 breast cancer patients in total, to evaluate whether TLR10 tumor expression was correlated with survival outcomes in breast cancer. We observed a statistically significant correlation between primary tumor expression of TLR10 and recurrence-free survival (RFS) in patients with breast cancer (Figure 2). Patients whose primary tumors expressed low levels of TLR10 possessed median RFS of 30 months, while patients whose tumors expressed high levels of TLR10 possessed median RFS of 47 months. This difference in RFS based on TLR10 tumor expression in patients with breast cancer was statistically significant (Figure 2, Chart 3; logrank p -value: $2.4e-6$; hazard ratio: 0.69 (0.59-0.8))

Thus, by mining published microarray data^{4,5} in an unbiased fashion, we identified toll-like receptor 10 encoded by TLR10, as among the genes whose expression was most different, transcriptome-wide, in the lymph node metastases and primary tumors of patients with breast cancer; we observed significantly increased expression of TLR10 in lymph node metastases as compared to primary tumors of the breast. Further, we found a significant correlation between TLR10 expression and patient survival outcomes, as recurrence-free survival was significantly

greater in patients whose primary tumors expressed higher levels of TLR10 as compared to patients whose primary tumors expressed lower levels of TLR10.

Discussion

We provided evidence here that toll-like receptor 10, encoded by TLR10, is among the genes whose expression is most different in the lymph node metastases of patients with lymph node metastatic breast cancer, that TLR10 mRNA is present at significantly increased quantities in lymph node metastatic tissues as compared to primary tumors of the breast, that TLR10 is also differentially expressed in primary tumors of patients with breast cancer and that primary tumor TLR10 expression is significantly correlated with patient survival outcomes in human breast cancer. The ligand for TLR10 is unknown⁸. Evaluation of the effects of genetic depletion of TLR10 in mouse models of metastatic breast cancer on metastasis to the lymph nodes and to the central nervous system is merited. Modulation of TLR10 expression may be relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier and colonize the brain. These data establish that a pattern recognition receptor with unknown ligand specificity is among the genes whose expression changes most significantly transcriptome-wide between primary tumors of the breast and metastases to the lymph nodes in humans with breast cancer.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalín, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Clarke, C., Madden, S.F., Doolan, P., Aherne, S.T., Joyce, H., O'driscoll, L., Gallagher, W.M., Hennessy, B.T., Moriarty, M., Crown, J. and Kennedy, S., 2013. Correlating transcriptional networks to breast cancer survival: a large-scale coexpression analysis. *Carcinogenesis*, 34(10), pp.2300-2308.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.
8. Oosting, M., Cheng, S.C., Bolscher, J.M., Vestering-Stenger, R., Plantinga, T.S., Verschuere, I.C., Arts, P., Garritsen, A., van Eenennaam, H., Sturm, P. and Kullberg, B.J., 2014. Human TLR10 is an anti-inflammatory pattern-recognition receptor. *Proceedings of the National Academy of Sciences*, 111(42), pp.E4478-E4484.

Rank: 117
Probe ID: 10635
p-value: 1.07E-03
t: -3.3963839
B: -0.8697
Gene: TLR10
Gene name: toll-like receptor 10

Chart 1: TLR10 is differentially expressed in lymph node metastatic breast cancer when comparing lymph node metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 19617
probe ID: 223751_x_at
p-value: 3.48E-02
t: 2.13
B: -4.9568471
Gene: TLR10
Gene name: toll-like receptor 10

Chart 2: TLR10 is differentially expressed in human breast cancer when comparing primary tumors of the breast to normal breast tissues.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: TLR10 is expressed at significantly higher levels in the lymph node metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of TLR10 in primary tumors of the breast (left) and in lymph node metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in TLR10 expression between primary tumors of the breast and lymph node metastases is $p=0.001055$.

Figure 2: Significant correlation between TLR10 primary tumor expression and recurrence-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=1764$ total patients stratified into two groups, based on low or high expression of TLR10 in patient primary tumors. The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on TLR10 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low TLR10 expression: 30 months

High TLR10 expression: 47 months

Chart 3: Median recurrence-free survival is inferior in patients with low primary tumor expression of TLR10.

The median RFS (recurrence-free survival) of patients with low primary tumor expression of TLR10 and high primary tumor expression of TLR10 is listed in this chart, for $n=1764$ breast cancer patients.