Stabilin 2, encoded by STAB2 is differentially expressed in metastatic breast cancer, both in metastases to the brain and to the lymph nodes.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes to discover genes associated with brain metastasis in patients with metastatic breast cancer. We found that stabilin 2, encoded by STAB2, was among the genes whose expression was most different in the brain and lymph node metastases of patients with metastatic breast cancer as compared to normal breast tissues. STAB2 mRNA was present at significantly reduced quantities in brain metastatic tissues as compared to normal breast tissues. Down-regulation of STAB2 may be an important event for metastasis of primary tumor-derived cancer cells to the brain in humans with metastatic breast cancer, and irrespective of its importance, appears to be a transcriptional event generally associated with metastasis to the brain. We have recently described transcriptome-wide differential expression of Factor XII (Hageman Factor) and the coagulation factor II thrombin receptor-like 2 (F2RL2) in brain metastatic breast cancer^{6,7}; thus, STAB2⁸, a von Willebrand Factor quantitative trait locus⁹ and therefore a genetic modifier of a major factor involved in clotting of the blood is among at least three components of the hematopoietic clotting cascade whose transcriptional perturbation comprises the transcriptional fingerprint of brain metastases in patients with metastatic breast cancer.

Keywords: breast cancer, metastasis, brain metastases, central nervous system metastases, STAB2, stabilin 2, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in trastuzumab-treated patients with breast cancer². This alarmingly high frequency of CNS metastasis events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with brain metastatic breast cancer^{4,5}. We discovered significant differential and decreased expression of the stabilin 2, encoded by STAB2, in brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE52604⁴ and GSE124648⁵ and for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE52604 was generated using Whole Human Genome Microarray 4x44K G4112F technology with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer, analysis performed using platform GPL6480. GSE124648, a microarray dataset, was generated using Affymetrix Human Genome U133A Array technology with $n=10$ normal breast tissues and $n=44$ lymph node metastases from patients with breast cancer; platform GPL96 was used for this analysis. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether STAB2 gene expression was significantly between normal breast tissues and brain metastases in humans with breast cancer using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

Results

We performed global comparative transcriptome analysis of metastatic tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of brain and lymph node metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

STAB2 is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Identification of differentially expressed genes across datasets in cancer can be difficult and more challenging than when doing so in benign organ tissue and sorted cell populations due to higher levels of heterogeneity in tumors, and differing methods of tumor sampling between laboratories. Thus, we asked how gene expression was most significantly different between brain metastases from patients with breast cancer and normal breast tissue as opposed to primary tumors of the breast, by comparing global gene expression profiles of 10 normal breast tissues to 35 brain metastases⁴. Using this strategy, we identified stabilin 2, STAB2, as among the genes whose expression was most significantly different in the brain metastases of patients with brain metastatic breast cancer⁴ (Table 1). When sorting each of the genes expressed in brain metastases based on significance of difference in expression between brain metastases and normal breast tissues, STAB2 ranked 10 out of 41093 total transcripts (Table 1), equating to 99.98% differential expression. Differential expression of STAB2 in brain metastases from patients with brain metastatic breast cancer in this dataset was statistically significant (Table 1; $p=1.53E-21$).

1 Next, we queried a second microarray dataset⁵ generated using lymph node metastases and
2 normal breast tissues from patients with breast cancer to determine whether we could validate differential
3 expression of STAB2 in metastatic tissues of humans with breast cancer, and whether differential
4 expression of STAB2 was specific to brain metastases or could be observed in metastases to other sites in
5 patients with metastatic breast cancer. When sorting each of the genes expressed in lymph node
6 metastases based on significance of difference as compared to normal breast tissues in patients with breast
7 cancer, STAB2 ranked 7548 out of 22283 total transcripts (Table 2), equating to 66.1% differential
8 expression. Differential expression of STAB2 in the lymph node metastases of patients with metastatic
9 breast cancer was statistically significant (Table 2; $p=3.65E-02$). Thus, differential expression of STAB2
10 was conserved in the brain and lymph node metastases of patients with metastatic breast cancer in two
11 independent microarray datasets^{4,5}.

12 **STAB2 is expressed at significantly lower levels in the brain metastases of patients with brain** 13 **metastatic breast cancer.**

14 We obtained exact mRNA expression levels for STAB2, in normal breast tissues and in brain
15 metastasis of patients with brain metastatic breast cancer to determine direction and statistical
16 significance of change in STAB2 expression in brain metastatic tissues. STAB2 was expressed at lower
17 levels in the brain metastases of patients with breast cancer as compared to normal breast tissues, and this
18 difference was statistically significant (Figure 1; $p<0.0001$).

19 Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified
20 STAB2 as among the genes whose expression was most different in the brain metastases of patients with
21 breast cancer both when compared to the normal breast; STAB2 expression, as compared to the benign
22 breast, was significantly lower levels in brain metastases.

23 **Discussion**

24 We provided evidence here that stabilin 2, encoded by STAB2, is among the genes whose
25 expression is most different in the brain metastases of patients with brain metastatic brain cancer, and that
26 STAB2 is expressed at significantly lower levels in brain metastases as compared normal breast tissues.
27 Independent evaluation of the effects of ectopic expression of STAB2 in mouse models of metastatic
28 breast cancer on metastasis to the central nervous system are merited.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Salhia, B., Kiefer, J., Ross, J.T., Metapally, R., Martinez, R.A., Johnson, K.N., DiPerna, D.M., Paquette, K.M., Jung, S., Nasser, S. and Wallstrom, G., 2014. Integrated genomic and epigenomic analysis of breast cancer brain metastasis. *PloS one*, 9(1), p.e85448.
5. Sinn, B.V., Fu, C., Lau, R., Litton, J., Tsai, T.H., Murthy, R., Tam, A., Andreopoulou, E., Gong, Y., Murthy, R. and Gould, R., 2019. SET ER/PR: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ breast cancer*, 5(1), pp.1-8.
6. Mamoor, S., 2020. Factor XII (Hageman factor) is over-expressed in brain metastatic breast cancer.
7. Mamoor, S., 2020. Coagulation factor II thrombin receptor like 2 (F2RL2) is differentially expressed in brain metastatic breast cancer.
8. Swystun, L.L., Lai, J.D., Notley, C., Georgescu, I., Paine, A.S., Mewburn, J., Nesbitt, K., Schledzewski, K., Géraud, C., Kzhyshkowska, J. and Goerdts, S., 2018. The endothelial cell receptor stabilin-2 regulates VWF-FVIII complex half-life and immunogenicity. *The Journal of clinical investigation*, 128(9), pp.4057-4073.
9. Smith, N.L., Chen, M.H., Dehghan, A., Strachan, D.P., Basu, S., Soranzo, N., Hayward, C., Rudan, I., Sabater-Lleal, M., Bis, J.C. and de Maat, M., 2010. Novel associations of multiple genetic loci with plasma levels of factor VII, factor VIII, and von Willebrand factor: The CHARGE Consortium. *Circulation*, 121(12), p.1382.
10. Park, S.Y., Jung, M.Y., Kim, H.J., Lee, S.J., Kim, S.Y., Lee, B.H., Kwon, T.H., Park, R.W. and Kim, I.S., 2008. Rapid cell corpse clearance by stabilin-2, a membrane phosphatidylserine receptor. *Cell Death & Differentiation*, 15(1), pp.192-201.
11. Politz, O., Gratchev, A., McCourt, P.A., Schledzewski, K., Guillot, P., Johansson, S., Svineng, G., Franke, P., Kannicht, C., Kzhyshkowska, J. and Longati, P., 2002. Stabilin-1 and-2 constitute a novel family of fasciclin-like hyaluronan receptor homologues. *Biochemical Journal*, 362(1), pp.155-164.
12. Park, S.Y., Yun, Y., Lim, J.S., Kim, M.J., Kim, S.Y., Kim, J.E. and Kim, I.S., 2016. Stabilin-2 modulates the efficiency of myoblast fusion during myogenic differentiation and muscle regeneration. *Nature communications*, 7(1), pp.1-15.

Rank	ID	p-value	t	B	Gene	Gene name
10	A_23_P162607	1.53E-21	-16.8573414	38.135946	STAB2	stabilin 2

Table 1: STAB2 is differentially expressed in brain metastases in patients with metastatic breast cancer when compared to normal breast tissues.

Rank of differential expression, probe ID, *p*-value with respect to differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
7548	220114_s_at	3.65E-02	2.14283	-4.743419	STAB2	stabilin 2

Table 2: STAB2 is differentially expressed in lymph node metastases of patients with metastatic breast cancer when compared to normal breast tissue.

Rank of differential expression, probe ID, *p*-value with respect to differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: STAB2 is expressed at significantly lower levels in the brain metastases of patients with brain metastatic breast cancer as compared to normal breast tissues.

The mRNA expression level of STAB2 in normal breast tissues (left) and in the brain metastases of brain metastatic breast cancer patients (right) is graphically represented here with mean STAB2 levels marked and the result of a statistical test evaluating difference in mRNA expression level between brain metastases and normal tissues of the breast, a *p*-value, listed above.