Myelin transcription factor 1 (MYT1) is differentially expressed in the tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab (Herceptin) is a monoclonal antibody targeting the extracellular domain of the human epidermal growth factor receptor 2 (HER2)¹ utilized for the treatment of adjuvant and metastatic breast cancer² in the United States and worldwide. We mined published microarray and gene expression data^{3,4} to discover in an unbiased manner the most significant transcriptional changes associated with trastuzumab treatment. We identified myelin transcription factor 1, encoded by MYT1 among the genes most differentially expressed in the primary tumors of patients with breast cancer treated with trastuzumab. The primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of MYT1 messenger RNA than did patients not treated with trastuzumab, and a single administration of trastuzumab was sufficient to result in differential expression of MYT1 in primary tumors of the breast, demonstrating increased expression of a zinc finger DNA-binding protein and neuronal-specific members of the LSD1 complex with the capacity to interact with Sin3B and with functions in neurogenesis⁵⁻⁹ as a direct transcriptional consequence of treatment with trastuzumab.

Keywords: trastuzumab, breast cancer, MYT1, systems biology of breast cancer, targeted therapeutics in breast cancer.

1 The mechanism of action ascribed to trastuzumab function is thought to be binding of the F_{ab}
2 domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2
3 (HER2)¹, but the transcriptional consequences resulting from signaling downstream of trastuzumab
4 binding to HER2 is less well understood. Through unbiased, systems-level comparative transcriptome
analysis of published microarray and gene expression data^{3,4}, we report here that MYT1 is a differentially
expressed gene in the tumors of trastuzumab-treated patients: patients treated with trastuzumab expressed
significantly higher levels of MYT1 than did those of patients not treated with trastuzumab.

5 **Methods**

6 We used datasets GSE130788³, and GSE114082⁴ for this differential gene expression analysis of
7 the transcriptional response to trastuzumab treatment in patients with breast cancer. GSE130788 was
8 generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with
9 *n*=11 primary breast tumors from patients treated with trastuzumab and *n*=36 primary tumors from
10 patients treated with trastuzumab, analysis performed using platform GPL6480. GSE114082 was
11 generated using Illumina HumanHT-12 WG-DASL V4.0 R2 expression beadchip technology with *n*=17
12 primary breast tumors from patients before trastuzumab treatment and *n*=17 primary tumors from patients
13 after trastuzumab treatment, analysis performed using platform GPL14951. The Benjamini and Hochberg
14 method of *p*-value adjustment was used for ranking of differential expression but raw *p*-values were used
15 to assess statistical significance of global or multiplexed differential expression. Log-transformation of
16 data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical
17 test was performed to evaluate whether MYT1 gene expression was significantly between primary breast
18 tumors of patients receiving trastuzumab using a two-tailed, unpaired t-test with Welch's correction. We
19 used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version
8.4.0)(455).

16 **Results**

17 We performed unbiased comparative transcriptome analysis using published microarray and gene
18 expression data^{3,4} to identify the most significant transcriptional features of the primary breast tumors of
patients treated with trastuzumab (Herceptin).

19 **MYT1 is differentially expressed in the tumors of patients treated with trastuzumab.**

20 We identified myelin transcription factor 1, encoded by MYT1 as among the genes whose
21 expression changed most significantly in the tumors of breast cancer patients treated with trastuzumab in
22 a multiplexed gene expression dataset³ (Table 1). When sorting each of the genes expressed in the
23 primary tumors of breast cancer patients treated with trastuzumab based on significance of change as
24 compared to that of patients not treated with trastuzumab, MYT1 ranked 88 out of 41000 total transcripts
(Table 1), equating to 99.8% differential expression. MYT1 differential expression in the tumors of
patients treated with trastuzumab trended towards statistical significance (Table 1; *p*=8.23E-59).

25 We turned to a second microarray dataset⁴ to validate our findings of MYT1 differential
26 expression in the patients treated with trastuzumab. The tumors compared in this dataset were sampled
27 from the same patient before and after a single, brief exposure to trastuzumab, not comparing tumors from
28 different patients treated with a course of trastuzumab or not. Thus, while differentially expressed genes
when comparing tumors based on trastuzumab treatment can result from a variety of sources, including
gene expression changes resulting from tumor evolution over the course of treatment, changes identified
in this dataset, occurring after only a single treatment of trastuzumab, are more likely to be a direct

transcriptional result of trastuzumab binding events. We found again that MYT1 was among the genes most differentially expressed in the tumors of patients administered a single treatment of trastuzumab (Table 2). When sorting each of the genes expressed in the tumors of patients administered a single treatment of trastuzumab, MYT1 ranked 5775 out of 29377 total transcripts (Table 2), equating to 80.3% differential expression. Differential expression of MYT1 in the primary tumors of breast cancer patients following treatment with trastuzumab was statistically significant (Table 2; $p=1.01E-02$).

MYT1 is expressed at significantly higher levels in the tumors of breast cancer patients treated with trastuzumab than in the tumors of those not treated with trastuzumab.

We obtained exact mRNA expression levels for MYT1 from tumors of patients treated with trastuzumab and from the tumors of patients not treated with trastuzumab to directly evaluate direction and statistical significance of change in expression of MYT1 based on history of trastuzumab treatment. MYT1 was expressed at higher levels in the tumors of patients treated with trastuzumab as compared to those of patients not treated with trastuzumab, and this difference was deemed statistically significant (Figure 1; $p<0.0001$).

Thus, we found using published multiplexed gene expression and microarray data^{3,4} that MYT1 was among the genes most differentially expressed in the tumors of patients treated with trastuzumab, that MYT1 was expressed at higher levels in the tumors of patients treated with trastuzumab as compared to those not treated with trastuzumab, and that this transcriptional change was likely a direct result of trastuzumab treatment as opposed to a gene expression change associated with clonal selection and evolution of the tumor across the course of treatment.

Discussion

We provided evidence here that MYT1 is a differentially expressed gene in the tumors of patients treated with trastuzumab, and expressed at significantly higher levels in the primary tumors of breast cancer patients treated with trastuzumab as compared to patients not treated with trastuzumab. Thus, trastuzumab administration in patients with breast cancer results in increased primary tumor expression of a zinc finger DNA-binding protein and subunit of a neuronal LSD1 histone demethylase complex, with the ability to interact with Sin3B and with functions in oligodendrocyte fate choice⁵⁻⁹.

References

1. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
2. Herceptin (Trastuzumab). Indications. <https://www.herceptin.com/hcp/indications.html>.
3. GSE130788. A Phase II Randomized Trial of Neoadjuvant Trastuzumab or Lapatinib or the Combination of Trastuzumab and Lapatinib, Followed by Docetaxel and Carboplatin with Trastuzumab and/or Lapatinib in Patients with HER2+ Breast Cancer. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788>. Hsiaowang Chen. UCLA, Department of Medicine. Santa Monica, CA.
4. Triulzi, T., Regondi, V., De Cecco, L., Cappelletti, M.R., Di Modica, M., Paolini, B., Lollini, P.L., Di Cosimo, S., Sfondrini, L., Generali, D. and Tagliabue, E., 2018. Early immune modulation by single-agent trastuzumab as a marker of trastuzumab benefit. *British journal of cancer*, 119(12), pp. 1487-1494.
5. Bellefroid, E.J., Bourguignon, C., Hollemann, T., Ma, Q., Anderson, D.J., Kintner, C. and Pieler, T., 1996. X-MyT1, a Xenopus C2HC-type zinc finger protein with a regulatory function in neuronal differentiation. *Cell*, 87(7), pp.1191-1202.
6. Vasconcelos, F.F., Sessa, A., Laranjeira, C., Raposo, A.A., Teixeira, V., Hagey, D.W., Tomaz, D.M., Muhr, J., Broccoli, V. and Castro, D.S., 2016. MyT1 counteracts the neural progenitor program to promote vertebrate neurogenesis. *Cell reports*, 17(2), pp.469-483.
7. Romm, E., Nielsen, J.A., Kim, J.G. and Hudson, L.D., 2005. Myt1 family recruits histone deacetylase to regulate neural transcription. *Journal of neurochemistry*, 93(6), pp.1444-1453.
8. Yokoyama, A., Igarashi, K., Sato, T., Takagi, K., Otsuka, M., Shishido, Y., Baba, T., Ito, R., Kanno, J., Ohkawa, Y. and Morohashi, K.I., 2014. Identification of myelin transcription factor 1 (MyT1) as a subunit of the neural cell type-specific lysine-specific demethylase 1 (LSD1) complex. *Journal of Biological Chemistry*, 289(26), pp.18152-18162.
9. Nielsen, J.A., Berndt, J.A., Hudson, L.D. and Armstrong, R.C., 2004. Myelin transcription factor 1 (Myt1) modulates the proliferation and differentiation of oligodendrocyte lineage cells. *Molecular and Cellular Neuroscience*, 25(1), pp.111-123.

Rank	ID	p-value	t	B	Gene	Gene name
88	A_23_P210835	8.23E-59	23.295316	123.41854	MYT1	myelin transcription factor 1

Table 1: MYT1 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene
5775	ILMN_1721167	1.01E-02	-2.7162819	-2.97667	MYT1

Table 2: A single exposure to trastuzumab is sufficient to drive differential expression of MYT1 in the primary tumors of patients with breast cancer.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: MYT1 is expressed at significantly higher levels in the primary tumors of patients treated with trastuzumab.

The mRNA expression level of MYT1 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between MYT1 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.