SSX2 is differentially expressed in models of MERS coronavirus infection.

Shahan Mamoor, MS¹

¹Thomas Jefferson School of Law

San Diego, CA 92101

mamoorsk@tjssl.edu

The coronavirus COVID19 pandemic is an emerging biosafety threat to the nation and the world (1). There are no treatments approved for coronavirus infection in humans (2) and there is a lack of information available regarding the basic transcriptional behavior of human cells and mammalian tissues following coronavirus infection. We mined multiple independent public (3) or published datasets (4-8) containing transcriptome data from infection models of human coronavirus 229E, the severe acute respiratory syndrome (SARS) coronavirus and Middle East respiratory syndrome (MERS) coronavirus to discover genes whose differential expression was conserved across the coronavirus family. We identified SSX2 (9) as a differentially expressed gene following infection of human cells specifically with two types of MERS coronaviruses. and not after infection of human cells with human coronavirus 229E, or and in the lungs of mice and ferrets infected with SARS coronavirus. An SSX2 interacting protein, SSX2IP, was among the genes most differentially expressed in the ferret blood after infection with SARS coronavirus. The expression of SSX2 is modulated to a degree unlike most any other gene following infection with MERS coronaviruses.

Keywords: SSX2, coronavirus, MERS coronavirus, SARS coronavirus, human coronavirus 229E, SARS-CoV-2, COVID19, systems biology of viral infection.

Viruses are classified according to a system known as the “Baltimore” classification of viruses (10) wherein the characteristics of the viral genome - whether it is positive-sense or negative-sense, whether it is single-stranded or double-stranded, whether it is composed of RNA or DNA - are used to group viruses into families. Coronaviruses are single-stranded, positive-sense RNA viruses that contain an envelope surrounding their viral particle (11). Their genome is largest of all RNA viruses, ranging from 27 to 33 kb in size (12). They obtain their name from the crown-like appearance of the viral particle imparted by the structure of the large-surface glycoprotein (12). The coronaviridae family includes seven viruses capable of infecting humans, including the severe acute respiratory distress syndrome, or SARS coronavirus (13), the Middle East respiratory syndrome coronavirus, or MERS coronavirus (14), the human coronaviruses (HCoV) 229E, OC43, HKU and NL63 (15-17), and the novel human coronavirus causing COVID19 infections now known as SARS-CoV-2 (18, 19). As of March 19, 2020, the World Health Organization reported 209,839 cases of COVID19 and 8778 deaths from SARS-CoV-2 infection world-wide (20). There are no FDA-approved treatments for human coronavirus infection.

We used a systems-level approach to identify the genes whose expression changes most significantly following infection of human cells with Middle East Respiratory Syndrome Coronavirus (MERS-CoV) using two independent datasets (3, 4) and compared these data to similar analyses of datasets generated using a human coronavirus 229E *in vitro* infection model (5) as well as two *in vivo* models of SARS coronavirus infection of the mouse lung (6, 8) and one *in vivo* model of SARS coronavirus infection of the ferret lung (7). Across both of the MERS-CoV datasets, we identified the SSX gene family member SSX2 as among the genes most differentially expressed following MERS coronavirus infection. SSX2 represents a transcriptional target of the host cell gene expression program following infection of human cells with MERS coronaviruses.

Methods

We used datasets GSE100509 (3) and GSE56677 (4), GSE89167 (5), GSE59185 (6), GSE22581 (7), and GSE68820 (8) for this systems-level differential gene expression analysis of coronavirus infections in conjunction with GEO2R.

GSE100509 was generated using Agilent-026652 Whole Human Genome Microarray 4x44K v2 technology. GSE56677 was generated using Agilent-039494 SurePrint G3 Human GE v2 8x60K Microarray 039381 technology. GSE89167 was generated using 039494 SurePrint G3 Human GE v2 8x60K Microarray 039381 technology. GSE59185 was generated using Agilent-028005 SurePrint G3 Mouse GE 8x60K Microarray technology. GSE68820 was generated using Agilent-014868 Whole Mouse Genome Microarray 4x44K G4122F technology. GSE22581 was generated using Affymetrix Canine Genome 2.0 Array technology.

The Benjamini and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used.

A statistical test was performed to evaluate the significance of difference in SSX2 mRNA expression levels in CALU3 2B4 cells with MERS-CoV infection as compared to CALU3 2B4 cells at baseline (0 hours) using a one-way ANOVA with Dunnett's multiple comparisons test. A statistical test was performed to evaluate the significance of difference in SSX2 mRNA expression levels in human primary microvascular endothelial cells with MERS-CoV as compared to baseline infection at 0 hours using a one-way ANOVA with Dunnett's multiple comparisons test. A statistical test was performed to evaluate the significance of difference between mRNA expression levels of SSXIP in uninfected ferret blood at day 0, and infected ferret blood at 2 days post-infection with SARS coronavirus using a two-tailed, unpaired t-test

with Welch's correction. Only p-values less than 0.05 were considered statistically significant.

We used PRISM for all statistical analyses (Version 8.4.0)(455).

Results

We mined two independent microarray datasets, public (3) and published (4) containing transcriptome data from models of MERS coronavirus infection in primary human cells and cell culture. We integrated this data with similar analyses of an *in vitro* infection model of human coronavirus 229E (5) and three *in vivo* models of SARS coronavirus infection, from the lungs of mice (6, 8) and in the blood of ferrets (7). Only in models of MERS coronavirus infection, we found that SSX2 was among the genes whose expression changed most significantly following infection with a coronavirus.

SSX2 is differentially expressed in primary human microvascular endothelial cells when comparing cells infected with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV EMC2012 and uninfected cells.

We identified SSX2 as differentially expressed following infection of primary human microvascular endothelial cells with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV EMC2012 when compared to non-infected cells (Table 1) (3). When sorting all of the transcripts expressed in human microvascular endothelial cells measured by microarray based on change in expression with and without infection, SSX2 ranked 2 out of 34127 transcripts. Differential expression of SSX2 in primary human microvascular endothelial cells following infection with MERS-CoV was statistically significant (Table 1; $p=1.41E-19$).

SSX2 is differentially expressed in the human cell line CALU3 2B4 when comparing cells infected with MERS-CoV London and uninfected cells.

We also identified SSX2 as differentially expressed in the human cell line CALU3 2B4 when comparing cells infected with MERS-CoV London from at 3 hours, 7 hours, 12 hours, 18 hours and 24 hours to cells infected at baseline (0 hours) (Table 2) (4). When sorting all of the transcripts expressed in CALU3 2B4 cells measured by microarray based on change in

expression with and without MERS-CoV London infection, SSX2 ranked 1 out of 28653 transcripts. Differential expression of SSX2 in CALU3 2B4 cells following infection with MERS-CoV London was statistically significant (Table 2; $p=5.63E-37$).

SSX2IP is differentially expressed in the blood of ferrets when comparing ferrets infected with SARS-CoV London and uninfected ferrets.

We also identified SSX2IP as differentially expressed in the blood of ferrets when comparing uninfected ferrets (day 0) to ferrets infected with SARS-CoV London 2 days post-infection (Table 3). When sorting all of the transcripts expressed in ferret blood measured by microarray based on change in expression with and without SARS-CoV infection, SSX2IP ranked 151 out of 43035 transcripts. Differential expression of SSX2IP in the blood of ferrets following infection with SARS-CoV was statistically significant (Table 3; $p=2.12E-03$).

SSX2 is transcriptionally induced following infection of primary human microvascular endothelial cells with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV EMC2012.

We extracted exact mRNA expression values for SSX2 from primary human microvascular endothelial cells infection with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV, and from uninfected primary human microvascular endothelial cells in order to compare expression levels of SSX2 between these two groups rather than relative to the rest of the transcriptome as assessed in differential gene expression analysis. This dataset contained transcriptome information from infection of primary human microvascular endothelial cells at 12 hours, 24 hours, 36 hours and 48 hours post-infection at compared to baseline (0 hours). We also performed a statistical test to evaluate whether the difference in expression of SSX2 in primary human microvascular endothelial cells infection with and without MERS-CoV infection was statistically significant. SSX2 was expressed at significantly higher levels at 12 hours, 24 hours, 36 hours, and 48 hours post-infection with MERS-CoV as compared to cells infected at baseline (0 hours) (Figure 1; $p<0.0001$ for all

comparisons). We calculated a fold change in SSX2 expression, relative to baseline (0 hours), of 1.7937 ± 0.0011 at 12 hours, of 1.7707 ± 0.0071 at 24 hours, of 1.7632 ± 0.0013 at 36 hours, and of 1.7746 ± 0.0021 at 48 hours post-infection (Table 1).

SSX2 is transcriptionally induced in the human cell line CALU3 2B4 following infection with MERS-CoV London.

We also extracted exact mRNA expression values for SSX2 from CALU3 2B4 cells following infection with MERS-CoV London and from CALU3 2B4 cells infected with MERS-CoV London at baseline in order to compare expression levels of SSX2 between these two groups. We also performed a statistical test to evaluate whether the difference in expression of SSX2 in CALU3 2B4 cells following infection with MERS-CoV London as compared to cells infected with MERS-CoV London at baseline was statistically significant. Consistent with the induction of SSX2 after infection of primary human microvascular endothelial cells with wild-type MERS-CoV, SSX2 was expressed at significantly higher levels post-infection MERS-CoV London at 7, 12, 18 and 24 hours (Figure 2; $p < 0.0001$ for 7, 12, 18 and 24 hours versus 0 hours). SSX2 was not significantly increased in expression at 3 hours post-infection (Figure 2; $p = .9999$). We calculated a fold change in SSX2 expression, relative to baseline (0 hours), of 1.0009 ± 0.0148 at 3 hours, of 1.5795 ± 0.0326 at 7 hours, of 1.7980 ± 0.0066 at 12 hours, of 1.8199 ± 0.0029 at 18 hours, and 1.8250 ± 0.0048 at 24 hours post-infection (Table 2).

SSX2IP is transcriptionally repressed in the blood of ferrets following infection with SARS-CoV.

We also extracted exact mRNA expression values for SSX2IP from the blood of ferrets following infection with SARS-CoV and from uninfected ferret blood in order to compare expression levels of SSX2IP between these two groups. We also performed a statistical test to evaluate whether the difference in expression of SSX2IP in the ferret blood 2 days post-infection with SARS-CoV was statistically significant. SSX2IP was expressed in the blood at significantly lower levels 2 days post-infection with SARS-CoV than in uninfected ferrets (Figure

3; $p=0.0026$). We calculated a fold change in SSX2IP expression, relative to baseline (day 0), of 0.4280 ± 0.1115 at 2 days post-infection (Table 3).

Thus, we found that SSX2 was among the genes whose expression changed most significantly after infection of human cells with MERS-CoV wild-type and MERS-CoV London, and that SSX2 was transcriptionally induced in primary human microvascular endothelial cells and in a human lung adenocarcinoma cell line following infection with wild-type MERS-CoV and MERS-CoV London, respectively. SSX2 was not significantly differentially expressed in a human cell line (Huh-7) following infection with human coronavirus 229E, and the transcript was not present or measured in a microarray dataset describing the transcriptional response *in vivo* to SARS coronavirus infections of the lung in ferrets or in mice (data not shown) (7, 6). An SSX2 interacting protein, SSX2IP, was among the genes most differentially expressed in blood of ferrets (7). Differential expression of SSX2 after coronavirus infection was specific to MERS coronaviruses, while differential expression of SSX2IP appeared to be specific to SARS coronavirus infection.

Discussion

COVID19 is a newly discovered coronavirus that infects humans and whose spread has lead to a global pandemic (15) with zero available therapeutic strategies. To facilitate target discovery and to further understanding of the basic transcriptional program of cells and tissues following coronavirus infection, we mined two independent microarray datasets containing transcriptome data of human cells infected with wild-type MERS-CoV and MERS-CoV London, in primary cells and in cell culture (3, 4) and integrated our findings with differential gene expression analyses using microarray data from *in vivo* models of SARS coronavirus infections of the mouse lung and in the ferret blood. Across two datasets, we identified SSX2 as one of the genes whose expression changed most significantly following infection with a MERS

1 coronavirus. SSX2 was not significantly differentially expressed in a human cell line (Huh-7)
2 infected with human coronavirus 229E. In both MERS experimental models analyzed, SSX2
3 expression significantly increased after infection with coronavirus. We also found differential
4 expression of the SSX2 interacting protein SSX2IP in the blood of ferrets following infection
5 with SARS coronavirus *in vivo*.

6
7 SSX2 belongs to the SSX gene family which is specifically expressed in
8 and the thyroid (20). Tumors from patients with synovial sarcoma are characterized by gene
9 fusion of either SSX1 or SSX2 to SYT, also known as SS18 (21). Along with RAB3IP, SSX2IP
10 was identified as an SSX2 interacting protein using a yeast two-hybrid screen (22). The SSX2-
11 SS18 gene fusion was found to modulate the expression of a number of genes including IGF2
12 and CD44; this involved changes in histone modifications at the promoters of IGF2 and CD44
13 (23). SSX2 has been demonstrated to antagonize the function of polycomb group proteins
14 BMI1 and EZH2, leading to derepression of polycomb target genes associated with negative
15 regulation of the trimethylation of lysine 27 of histone 3 (H3K27me3) (24). SSX2IP is a protein
16 associated with the mitotic spindle apparatus and appears to be involved in promoting the
17 maturation of and maintaining the integrity of the centrosome (25).

18
19 We found that SSX2 was differentially expressed in a human cell line, CALU3 2B4
20 following infection with the human coronavirus MERS-CoV London as well as in primary human
21 endothelial cells after infection with wild-type MERS-CoV. SSX2 was not significantly
22 differentially expressed in a human cell line (Huh-7) infected with human coronavirus 229E.
23 SSX2 and SSX2IP expression should be evaluated in the cells and tissues of patients infected
24 with the novel coronavirus SARS-CoV-2 to assess whether COVID19 infection similarly leads to
25 induction of SSX2IP. The fact that SSX2 and SSX2IP are both among the genes most
26 differentially expressed after infections with different members of the coronavirus family
27 suggests that these genes mediate some underlying process associated with coronavirus
28 infection.

References

1. World Health Organization, 2020. Coronavirus disease 2019 (COVID-19): situation report, 49.
2. Wang, Y. and Zhu, L.Q., 2020. Pharmaceutical care recommendations for antiviral treatments in children with coronavirus disease 2019. *World Journal of Pediatrics*, pp.1-4.
3. GSE100509. Baric R, Sims A, Heller N, Waters KM, Einfeld AJ, Kawaoka Y. PNNL. 902 Battelle Blvd. Richland, USA.
4. Selinger, C., Tisoncik-Go, J., Menachery, V.D., Agnihothram, S., Law, G.L., Chang, J., Kelly, S.M., Sova, P., Baric, R.S. and Katze, M.G., 2014. Cytokine systems approach demonstrates differences in innate and pro-inflammatory host responses between genetically distinct MERS-CoV isolates. *BMC genomics*, 15(1), p.1161.
5. Poppe, M., Wittig, S., Jurida, L., Bartkuhn, M., Wilhelm, J., Müller, H., Beuerlein, K., Karl, N., Bhujju, S., Ziebuhr, J. and Schmitz, M.L., 2017. The NF- κ B-dependent and-independent transcriptome and chromatin landscapes of human coronavirus 229E-infected cells. *PLoS pathogens*, 13(3), p.e1006286.
6. Regla-Nava, J.A., Nieto-Torres, J.L., Jimenez-Guardeño, J.M., Fernandez-Delgado, R., Fett, C., Castaño-Rodríguez, C., Perlman, S., Enjuanes, L. and DeDiego, M.L., 2015. Severe acute respiratory syndrome coronaviruses with mutations in the E protein are attenuated and promising vaccine candidates. *Journal of virology*, 89(7), pp.3870-3887.
7. Danesh, A., Cameron, C.M., León, A.J., Ran, L., Xu, L., Fang, Y., Kelvin, A.A., Rowe, T., Chen, H., Guan, Y. and Jonsson, C.B., 2011. Early gene expression events in ferrets in response to SARS coronavirus infection versus direct interferon-alpha2b stimulation. *Virology*, 409(1), pp.102-112.
8. Totura, A.L., Whitmore, A., Agnihothram, S., Schäfer, A., Katze, M.G., Heise, M.T. and Baric, R.S., 2015. Toll-like receptor 3 signaling via TRIF contributes to a protective innate immune response to severe acute respiratory syndrome coronavirus infection. *MBio*, 6(3), pp.e00638-15.
9. Piddini, E., Schmid, J.A., de Martin, R. and Dotti, C.G., 2001. The Ras-like GTPase SSX2 is involved in cell shape remodelling and interacts with the novel kinesin-like protein KIF9. *The EMBO journal*, 20(15), pp.4076-4087.

- 1 10. Baltimore, D., 1971. Expression of animal virus genomes. *Bacteriological reviews*, 35(3), p.
2 235.
- 3 11. Sawicki, S.G., Sawicki, D.L. and Siddell, S.G., 2007. A contemporary view of coronavirus
4 transcription. *Journal of virology*, 81(1), pp.20-29.
- 5 12. Lai, M.M. and Cavanagh, D., 1997. The molecular biology of coronaviruses. In *Advances in*
6 *virus research* (Vol. 48, pp. 1-100). Academic Press.
- 7 13. Peiris, J.S.M., Lai, S.T., Poon, L.L.M., Guan, Y., Yam, L.Y.C., Lim, W., Nicholls, J., Yee,
8 W.K.S., Yan, W.W., Cheung, M.T. and Cheng, V.C.C., 2003. Coronavirus as a possible
9 cause of severe acute respiratory syndrome. *The Lancet*, 361(9366), pp.1319-1325.
- 10 14. Lu, G., Hu, Y., Wang, Q., Qi, J., Gao, F., Li, Y., Zhang, Y., Zhang, W., Yuan, Y., Bao, J. and
11 Zhang, B., 2013. Molecular basis of binding between novel human coronavirus MERS-CoV
12 and its receptor CD26. *Nature*, 500(7461), pp.227-231.
- 13 15. van der Hoek, L., Pyrc, K., Jebbink, M.F., Vermeulen-Oost, W., Berkhout, R.J., Wolthers,
14 K.C., Wertheim-van Dillen, P.M., Kaandorp, J., Spaargaren, J. and Berkhout, B., 2004.
15 Identification of a new human coronavirus. *Nature medicine*, 10(4), pp.368-373.
- 16 16. Woo, P.C., Lau, S.K., Chu, C.M., Chan, K.H., Tsoi, H.W., Huang, Y., Wong, B.H., Poon,
17 R.W., Cai, J.J., Luk, W.K. and Poon, L.L., 2005. Characterization and complete genome
18 sequence of a novel coronavirus, coronavirus HKU1, from patients with
19 pneumonia. *Journal of virology*, 79(2), pp.884-895.
- 20 17. Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R.
21 and Niu, P., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *New*
22 *England Journal of Medicine*.
- 23 18. Bai, Y., Yao, L., Wei, T., Tian, F., Jin, D.Y., Chen, L. and Wang, M., 2020. Presumed
24 asymptomatic carrier transmission of COVID-19. *Jama*.
- 25 19. World Health Organization, 2020. Coronavirus disease 2019 (COVID-19): situation report,
26 59.
- 27 20. Gure, A.O., Türeci, Ö., Sahin, U., Tsang, S., Scanlan, M.J., Jäger, E., Knuth, A.,
28 Pfreundschuh, M., Old, L.J. and Chen, Y.T., 1997. SSX: a multigene family with several
members transcribed in normal testis and human cancer. *International journal of*
cancer, 72(6), pp.965-971.

21. Crew, A.J., Clark, J., Fisher, C., Gill, S., Grimer, R., Chand, A., Shipley, J., Gusterson, B.A. and Cooper, C.S., 1995. Fusion of SYT to two genes, SSX1 and SSX2, encoding proteins with homology to the Kruppel-associated box in human synovial sarcoma. *The EMBO journal*, 14(10), pp.2333-2340.
22. de Bruijn, D.R., dos Santos, N.R., Kater-Baats, E., Thijssen, J., van den Berk, L., Stap, J., Balemans, M., Schepens, M., Merks, G. and Geurts van Kessel, A., 2002. The cancer-related protein SSX2 interacts with the human homologue of a Ras-like GTPase interactor, RAB3IP, and a novel nuclear protein, SSX2IP. *Genes, Chromosomes and Cancer*, 34(3), pp. 285-298.
23. de Bruijn, D.R., Allander, S.V., van Dijk, A.H., Willemse, M.P., Thijssen, J., van Groningen, J.J., Meltzer, P.S. and van Kessel, A.G., 2006. The synovial sarcoma-associated SS18-SSX2 fusion protein induces epigenetic gene (de) regulation. *Cancer research*, 66(19), pp. 9474-9482.
24. Gjerstorff, M.F., Relster, M.M., Greve, K.B.V., Moeller, J.B., Elias, D., Lindgreen, J.N., Schmidt, S., Mollenhauer, J., Voldborg, B., Pedersen, C.B. and Brückmann, N.H., 2014. SSX2 is a novel DNA-binding protein that antagonizes polycomb group body formation and gene repression. *Nucleic acids research*, 42(18), pp.11433-11446.
25. Bärenz, F., Inoue, D., Yokoyama, H., Tegha-Dunghu, J., Freiss, S., Draeger, S., Mayilo, D., Cado, I., Merker, S., Klinger, M. and Hoeckendorf, B., 2013. The centriolar satellite protein SSX2IP promotes centrosome maturation. *Journal of Cell Biology*, 202(1), pp.81-95.

Rank	ID	p-value	F	Fold change	Gene	Gene name
2	A_24_P256243 NM_175698	5.63E-37	8990.3104	3hr: 1.7937 ± 0.0011 7hr: 1.7707 ± 0.0071 12hr: 1.7632 ± 0.0013 18hr: 1.7746 ± 0.0021 24hr: 1.7937 ± 0.0011	SSX2	synovial sarcoma, X breakpoint 2

Table 1: SSX2 is the most differentially expressed gene in primary human microvascular endothelial cells when comparing cells infected with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV EMC2012 and uninfected cells.

The rank of differential expression globally, the probe/transcript ID, the p-value of differential expression globally, F, a statistic used when more than two groups are compared in differential gene expression analysis (rather than the moderated t-statistic “t”), the gene and gene name are listed in this chart.

Rank	ID	p-value	F	Fold change	Gene	Gene name
1	A_24_P256243/ NM_175698	1.41E-19	1414.5025	3hr: 1.0009 ± 0.0148 7hr: 1.5795 ± 0.0326 12hr: 1.7980 ± 0.0066 18hr: 1.8199 ± 0.0029 24hr: 1.8250 ± 0.0048	SSX2	synovial sarcoma, X breakpoint 2

Table 2: SSX2 is among the most differentially expressed genes in human lung adenocarcinoma cells (CALU3 2B4) when comparing cells infected with MERS-CoV London from 3 to 24 hours and cells infected at baseline.

The rank of differential expression globally, the probe/transcript ID, the p-value of differential expression globally, F, a statistic used when more than two groups are compared in differential gene expression analysis (rather than the moderated t-statistic “t”), the gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Fold change	Gene	Gene name
151	CfaAffx. 31094.1.S1_s_at	2.12E-03	-4.886437	-1.1647	0.428 ± 0.1115	SSX2IP	synovial sarcoma, X breakpoint 2 interacting protein

Table 3: SSX2IP is among the most differentially expressed in the blood of ferrets infected with SARS coronavirus when compared to the lungs of ferrets on Day 0.

The rank of differential expression globally, the probe/transcript ID, the p-value of differential expression globally, t, a moderated t statistic, B, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: SSX2 is transcriptionally induced in primary human microvascular endothelial cells following infection with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV.

Expression of SSX2 at the mRNA level is graphically represented with the mean value marked and the p-value listed for each comparison relative to baseline infection at 0 hours.

Figure 2: SSX2 is transcriptionally induced in the human lung adenocarcinoma cell line CALU3 2B4 after infection with MERS-CoV London.

Expression of SSX2 at the mRNA level is graphically represented with the mean value marked and the p-value listed for each comparison relative to baseline infection at 0 hours.

Figure 3: SSX2IP is transcriptionally induced in the blood of ferrets infected with SARS coronavirus.

Expression of SSX2IP at the mRNA level is graphically represented with the mean value marked and the p-value listed for each comparison relative to baseline infection at 0 hours.