1 Opinion

2 Repurposing Amantadine and Memantine for COVID-19: Guanylate Kinases Might Pave
3 the Road towards Novel SARS CoV-2 and Anti Neoplastic Therapeutics.

4 Mina T. Kelleni, MD, PhD

5 Pharmacology Department, College of Medicine, Minia University, Egypt.

6 Mobile: +201200382422

7 drthabetpharm@yahoo.com, mina.kelleni@mu.edu.eg

8 <https://orcid.org/0000-0001-6290-6025>

9

10

11

12

13

14

15

16

17

18

19

20 *To the Editor,*

21 Guanylate kinase 1 (GK) is a highly specific cytosolic nucleoside monophosphate kinase
22 that is considered an essential enzyme in nucleotide metabolism pathways that solely
23 catalyzes the phosphorylation of guanosine monophosphate to diphosphate which acts as a
24 substrate for GTP/dGTP synthesis used for nucleic acid or cyclic GMP synthesis.
25 Importantly, GK plays a critical and essential role in metabolic activation of antiviral and
26 antineoplastic nucleoside-analogs prodrugs, including 6-thioguanine, 6-mercaptopurine,
27 acyclovir, ganciclovir and cyclopropavir. Interestingly, GK has been suggested to play a
28 fundamental role in SARS CoV-2 replication and in-vitro experiments testing the potential
29 benefits of GK inhibitors were suggested by Renz et al. (1) and other colleagues have
30 confirmed a GK role in SARS CoV-2 replication while adding more targets to be
31 considered(2).

32 However, we would like to emphasize that specific GK inhibitors, e.g. the nucleotide
33 analogue Ap5G, were only used to experimentally and owing to the various physiological
34 known and potentially yet to be discovered physiological functions of GK we suggest that
35 its specific inhibitors should be carefully assessed as regards to their safety to manage
36 COVID-19. On the other hand, GK gene was localized in 1980 on chromosome one and
37 later its encoding was described and it is currently readily available
38 [<https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=2987>] and
39 interestingly, genetic variants of other kinases have been described to negatively impact
40 their action (3), thus if GK was proved crucial for SARS CoV-2 replication, its genetic
41 polymorphisms might help to further explain some encountered idiosyncratic COVID-19
42 morbidity and mortality outcomes(4). Moreover, overexpression of GK gene was

43 previously suggested to be involved in the pathogenesis of some tumors together with an
44 unfulfilled call for exploration of some unexplored physiological functions that GK might
45 be involved (5) and we wish to repeat this call for the best interests of scientific medical
46 and pharmacological research. Additionally, we recommend testing another potential that
47 the non-authentic membrane-associated guanylate kinases, that contain GK like domains,
48 might share in SARS CoV-2 pathogenesis as these enzymes are known to regulate the
49 CNS NMDA receptors and their potential SARS CoV-2 interaction should be investigated
50 especially if the preliminary reports showing a potential benefit of the NMDA receptor
51 antagonist amantadine, which is currently used in treatment of parkinsonism and was used
52 in treatment and prophylaxis of influenza A virus infection before no longer recommended
53 because of viral resistance, proved clinically significant (6, 7) as we postulate that SARS
54 CoV-2 is still naïve and that amantadine uncompetitive NMDA antagonism might involve
55 inhibition of those membrane associated guanylate kinases and/or GK, to be also noted that
56 invitro antineoplastic properties of amantadine have been previously described.

57 Similarly, memantine, a dimethyl derivative of amantadine, an EMEA and FDA approved
58 drug for treatment of moderate to severe Alzheimer's disease, has been suggested to possess
59 beneficial effects in management of COVID-19 via several mechanisms (8). Memantine
60 is known to possess a higher affinity and more potent antagonism towards NMDA
61 receptors than amantadine, thus the same, and potentially better, suggested interaction with
62 GKs might be also applicable for memantine. Notably, both drugs have been suggested for
63 COVID-19 management basing on their anti-inflammatory and immune-modulatory
64 properties(9) as well as potential beneficial lysosomal dependent and neuroprotective
65 effects(10). Finally, we recommend that a short course of amantadine might be first

considered to be tested for selected pediatric severe to critical COVID-19 patients and of memantine for the adult and geriatric severe to critical cases while considering the potential contraindications, dosage modifications and drug-drug interactions.

Funding/Financial Exposure

None

Conflict of interest

The author declares that there is no conflict of interest

References

1. A. Renz, L. Widerspick, A. Dräger, FBA reveals guanylate kinase as a potential target for antiviral therapies against SARS-CoV-2. *Bioinformatics* **36**, i813-i821 (2020).
2. H. Delattre, K. Sasidharan, O. S. Soyer, Inhibiting the reproduction of SARS-CoV-2 through perturbations in human lung cell metabolic network. *Life Science Alliance* **4**, e202000869 (2021).
3. J. M. Lade, E. E. To, C. W. Hendrix, N. N. Bumpus, Discovery of Genetic Variants of the Kinases That Activate Tenofovir in a Compartment-specific Manner. *EBioMedicine* **2**, 1145-1152 (2015).
4. E. Pairo-Castineira *et al.*, Genetic mechanisms of critical illness in Covid-19. *Nature*, (2020).
5. A. A. da Rocha *et al.*, Hepatocyte growth factor-regulated tyrosine kinase substrate (HGS) and guanylate kinase 1 (GUK1) are differentially expressed in GH-secreting adenomas. *Pituitary* **9**, 83-92 (2006).
6. P. Grieb, M. Świątkiewicz, K. Prus, K. Rejdak, Amantadine for COVID-19. *The Journal of Clinical Pharmacology* **61**, 412-413 (2021).
7. R. Araújo, J. D. Aranda-Martínez, G. E. Aranda-Abreu, Amantadine Treatment for People with COVID-19. *Archives of Medical Research* **51**, 739-740 (2020).
8. S. Hasanagic, F. Serdarevic, Potential role of memantine in the prevention and treatment of COVID-19: its antagonism of nicotinic acetylcholine receptors (nAChR) and beyond. *European Respiratory Journal*, 2001610 (2020).
9. F. J. Jimenez-Jimenez, H. Alonso-Navarro, E. Garcia-Martin, J. A. G. Agundez, Anti-Inflammatory Effects of Amantadine and Memantine: Possible Therapeutics for the Treatment of Covid-19? *J Pers Med* **10**, (2020).
10. S. R. Brenner, The potential of memantine and related adamantanes such as amantadine, to reduce the neurotoxic effects of COVID-19, including ARDS and to reduce viral replication through lysosomal effects. *J Med Virol* **92**, 2341-2342 (2020).