Vulnerable groups at increased risk of COVID-19 in sub-Saharan Africa: The case of the HIV population

Marília R. Nepomuceno*

Max Planck Institute for Demographic Research, Rostock, Germany

The COVID-19 disease is spreading quickly around the globe. As the COVID-19 outbreak advances, governments are warning individuals at high risk to strictly follow restrictions to control the outbreak. To date, most information about the risk factors of the COVID-19 comes from China, South Korea, and Europe. Since the first outbreaks in China, more than 50 thousand individuals died.¹ Deaths are highest in Italy, Spain, China, France, and in the US.¹ Individuals above age 60 are at higher risk of dying if infected with the novel coronavirus.^{2,3} In Italy, the country that registered nearly one-fourth of the deaths worldwide, 90% of the deaths were above age 70, and 10% above age 90. Also, individuals with diabetes, cardiovascular and respiratory diseases, and cancer are at greater risk of death by COVID-19.^{3,4}

More recently, the COVID-19 disease is advancing fast towards different settings. Regions with a high burden of infectious diseases such as HIV, malaria, and dengue including Latin America and Africa are experiencing an increasing number of confirmed cases and deaths. Since the age structure and the distribution of relevant co-morbidities vary substantially by country, the risk profile for COVID-19 could be very different in some low income settings from that observed to date in China, South Korea and Europe.^{5,6}

Here, I want to draw attention to potential differences in the risk profile for COVID-19 in high HIV prevalence settings. Most people living with HIV are concentrated in sub-Saharan Africa, where the first death by COVID-19 was registered on the 19th of March.⁶ Since no country currently experiencing more advanced stages of the COVID-19 outbreak has a high prevalence of HIV, it is still unknown if the risk profile for COVID-19 derived from high-income settings will be similar in other regions such as sub-Saharan Africa.

*Corresponding author. E-mail: nepomuceno@demogr.mpg.de

In 2017, an estimated 36.8 million individuals were living with HIV in the globe, which makes this a major global public health issue.⁷ Despite the advancements in antiretroviral therapy, nearly a million of individuals died by HIV in 2017.⁷ Most HIV deaths are due to co-infections with other diseases including tuberculosis and hepatitis C.⁸ Also, individuals living with HIV are at increased risk of developing a range of noncommunicable diseases such as hypertension and cardiovascular disease.⁸ HIV-infected individuals are also at greater risk of dying related to influenza.⁹ This risk decreased for those who are on antiretroviral therapy, but it does not disappear. These are reasons that make the HIV population a potentially vulnerable group at increased risk of COVID-19.

Figure 1 shows the age distribution of the people living with HIV in different settings. Young adults (20–49 years) comprise most of the HIV population. Despite this similarity across countries, differences in population size are striking. In South Africa nearly 5.2 million individuals are living with HIV between ages 20 and 49, a number that is 50 times higher than the HIV-infected population of Italy, 8 times higher than that of Brazil, and 3 times higher than that of Mozambique.

There is no evidence as yet that people infected by HIV are at higher risk for severe illness or death from COVID-19. However, they are not immune to get infected by the novel coronavirus. Information from countries that have been testing both symptomatic and asymptomatic individuals such as South Korea and Iceland revealed that the age distribution of those infected by the novel coronavirus is similar to that of the total population.^{10,11} If the same is true in other settings, one can expect a massive number of infections at young and adult ages in low-income countries, which raises the relevance of the underlying co-morbidities among young adults.

In sub-Saharan Africa, HIV is the leading cause of morbidity at young and adult ages. More than 20% of individuals between age 15 and 49 are living with HIV in South Africa, Lesotho, and Botswana, and more than 10% in Mozambique and Namibia.⁸ This highlights the impact the COVID-19 outbreak may have in sub-Saharan African populations. A large number of deaths by COVID-19 at young and adult ages may occur as a consequence of the greater vulnerability of the young population in some sub-Saharan Africa countries compared to that of high-income settings.

Governments should be aware that differences in the distribution of co-morbidities by country can result in an increasing demand for hospitalization, and deaths by COVID-19 among young adults. In high HIV prevalence settings, particular attention should be given to young indi-

viduals that due their morbidity profile may be a more vulnerable group at increased risk of COVID-19.

Figure 1: Distribution of the HIV population by age and sex: Italy, Brazil, South Africa and Mozambique.
Source: *GBD(2017)*.

References

1. Johns Hopkins Center for Systems Science and Engineering. Coronavirus COVID-19 Global Cases. (2020). <https://systems.jhu.edu/research/public-health/ncov/> (accessed April 3, 2020).
2. Wu, D., Wu, T., Liu, Q. and Yang Z. (2020). The SARS-CoV-2 Outbreak: What We Know. *International Journal of Infectious Diseases*.
DOI: <https://doi.org/10.1016/j.ijid.2020.03.004>

3. Remuzzi, A and Remuzzi, G. (2020) COVID-19 and Italy: what next?. *Lancet*.
DOI: [https://doi.org/10.1016/S0140-6736\(20\)30627-9](https://doi.org/10.1016/S0140-6736(20)30627-9)
4. Zhou, F., Yu, t., Du, R., Fan, G., Liu, Y., Liu, Z., Xiang, J. et al. (2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*, 395: 1054–62.
DOI: [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3)
5. Dowd, J., Rotondi, V., Andriano, L., Brazel, D. M., Block, P., Ding, X., Liu, Y., Mills, M. (2020). Demographic science aids in understanding the spread and fatality rates of COVID-19. OSF Preprint.
https://osf.io/se6wy/?view_only=c2f00dfe3677493faa421fc2ea38e295
6. Walker, Whittaker, Watson et al. (2020) The Global Impact of COVID-19 and Strategies for Mitigation and Suppression. WHO Collaborating Centre for Infectious Disease Modelling, MRC Centre for Global Infectious Disease Analysis, Abdul Latif Jameel Institute for Disease and Emergency Analytics, Imperial College London.
7. Global Health Data Exchange (2017) Results. Seattle, United States: Institute for Health Metrics and Evaluation (IHME). <http://ghdx.healthdata.org/gbd-2017> (accessed March 29, 2020).
8. UNAIDS (2020) <https://www.unaids.org/en> (accessed March 31, 2020)
9. Cohen, C., Simonsen, L., Sample, J., Kang, J-W, Influenza-Related Mortality Among Adults Aged 25—54 Years With AIDS in South Africa and the United States of America. *Clinical Infectious Diseases*, 55(7):996–1003
DOI: <https://doi.org/10.1093/cid/cis549>
10. Backhaus, A. (2020). Coronavirus Why it’s so deadly in Italy. Demographics and why they are a warning to other countries. Medium. <https://medium.com/@andreasbackhausab/coronavirus-why-its-so-deadly-in-italy-c4200a15a7bf>
11. The National University Hospital of Iceland (2020). <https://www.covid.is/data> (accessed March 29, 2020).