Differential expression of CACHD1 in cancers of the breast.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY 11730

Breast cancer affects women at relatively high frequency¹. We mined published microarray datasets^{2,3} to determine in an unbiased fashion and at the systems level genes most differentially expressed in the primary tumors of patients with breast cancer. We report here significant differential expression of the gene encoding cache domain containing 1, CACHD1, when comparing primary tumors of the breast to the tissue of origin, the normal breast. CACHD1 mRNA was present at significantly lower quantities in tumors of the breast as compared to normal breast tissue. Analysis of human survival data revealed that expression of CACHD1 in primary tumors of the breast was correlated with post-progression survival in patients with luminal A subtype cancer, demonstrating a relationship between primary tumor expression of a differentially expressed gene and patient survival outcomes influenced by molecular subtype. CACHD1 may be of relevance to initiation, maintenance or progression of cancers of the female breast.

Keywords: breast cancer, CACHD1, cache domain containing 1, systems biology of breast cancer, targeted therapeutics in breast cancer.

Invasive breast cancer is diagnosed in over a quarter of a million women in the United States each year¹ and in 2018, breast cancer was the leading cause of cancer death in women worldwide⁴. While patients with localized breast cancer are provided a 99% 5-year survival rate, patients with regional breast cancer, cancer that has spread to lymph nodes or nearby structures, are provided an 86% 5-year survival rate^{5,6}. Patients with metastasis to distant sites, like the brain, are provided a 27% 5-year survival rate^{5,6}. Understanding how primary tumors are most transcriptionally different from the tissue from which they originate, the breast, can facilitate development of novel diagnostic and therapeutics to promote early detection and enhanced treatment, and contribute to efforts to prevent progression to metastatic stages. We mined published microarray data^{2,3} to understand at the transcriptome level and in an unbiased fashion genes most differentially expressed in primary tumors of the breast as compared to normal breast tissue. Cache domain containing 1 emerged as among the most differentially expressed genes in cancer of the female breast.

Methods

We utilized datasets GSE109169² and GSE42568³ for this differential gene expression analysis of female breast cancer. GSE109169 was generated using Affymetrix Human Exon 1.0 ST Array technology with $n=25$ normal breast tissue and $n=25$ tumors of the breast; analysis was performed using platform GPL5175. The tissues from this first dataset are paired tissues (25 tumors matching 25 breast tissues from 25 patients). GSE42568 was generated using Affymetrix Human Genome U133 Plus 2.0 array technology with $n=17$ normal breast tissue biopsies and $n=104$ primary breast tumor biopsies from patients with breast cancer; analysis was performed using platform GPL570; the majority of patients whose tumors were analyzed were age 50 or older. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether CACHD1 expression was significantly different between primary breast tumors and breast tissue using a two-tailed t -test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of CACHD1 mRNA expression levels with post-progression survival (PPS) in $n=36$ patients with basal subtype cancer, $n=76$ patients with luminal A subtype cancer, $n=38$ patients with luminal B subtype cancer, and $n=30$ patients with HER2+ cancer.

Results

We performed discovery of genes associated with breast cancer in females by mining two independently published microarray datasets^{2,3}.

CACHD1 is differentially expressed in primary tumors of the breast.

Comparison of 25 normal breast tissues to 25 tumors of the breast² from patients with early-onset breast cancer revealed that the gene encoding the cache domain containing 1, CACHD1, was among the genes most differentially expressed in tumors of the breast in human breast cancer (Chart 1). When sorting each of the genes expressed in tumors of the breast based on significance of difference as compared to normal breast tissue, CACHD1 ranked 105 out of 19076 total transcripts, equating to 99.4% differential expression (Chart 1). Differential expression of CACHD1 in female breast cancer was statistically significant (Chart 1; $p=1.46E-14$).

Analysis of a second microarray dataset³, here in 104 tumors from patients aged 31 to 89 at diagnosis, the majority of which were above age 50, again revealed significant differential expression of CACHD1 in human breast cancer (Chart 2). When sorting each of the genes expressed in the tumors of

patients with breast cancer based on significance of difference as compared to normal breast tissue, CACHD1 ranked 4365 out of 54675 total transcripts, equating to 92.0% differential expression (Chart 2). Differential expression of CACHD1 in the tumors of patients with breast cancer was statistically significant (Chart 2; $p=7.73E-08$). These data suggested that differential expression of CACHD1 was not an artifact of a single microarray dataset, nor was it strictly associated with early-onset breast cancer, rather a general feature of cancers of the breast.

CACHD1 is expressed at significantly lower levels in breast tumors as compared to the breast.

We obtained exact mRNA expression levels for CACHD1 from the breast and from breast tumors to understand the magnitude and direction of CACHD1 expression change. CACHD1 was expressed at lower levels in tumors of the breast as compared to normal breast tissue (Figure 1). Decreased expression of CACHD1 in primary breast tumors was statistically significant (Figure 1: $p<0.0001$). CACHD1 was expressed at 9.00 ± 0.45 arbitrary units (AU) in normal breast tissue, while it was expressed at 7.41 ± 0.62 AU in tumors of the breast. We calculated a mean fold change of 0.82 in CACHD1 mRNA levels when comparing primary tumors of the breast to normal breast tissues.

CACHD1 expression correlates with survival outcomes in luminal A subtype human breast cancer.

We performed Kaplan-Meier survival analysis in an attempt to correlate CACHD1 mRNA expression levels with survival outcomes in patients with breast cancer, evaluating correlations between molecular subtype and post-progression survival. We observed a correlation between CACHD1 expression and post-progression survival (PPS) in patients with luminal A subtype breast cancer which approached statistical significance (Figure 2; log rank p -value: 0.069 for post-progression survival, hazard ratio: 1.7 (0.95-3.02) (Fig. 2)). CACHD1 mRNA levels were a negative prognostic indicator in luminal A subtype breast cancer patients. Median PPS was 38.67 months for luminal A patients with low tumor expression of CACHD1 while median PPS was 26.8 months for luminal A patients with high tumor expression of CACHD1 (Chart 3). CACHD1 primary tumor expression was not correlated with post-progression survival in basal subtype (Figure 2; log rank p -value: 0.3 for PPS, hazard ratio: 1.51 (0.68-3.35) (Fig. 2), luminal B breast cancer (Figure 2; log rank p -value: 0.89 for PPS, hazard ratio: 0.95 (0.45-1.98) (Fig. 2)) or HER2+ breast cancer (Figure 2; log rank p -value: 0.2 for PPS, hazard ratio: 1.74 (0.74-4.14) (Fig. 2)).

Thus, through comparative transcriptome analysis of primary tumors of the breast and normal breast tissue, we found that differential expression and down-regulation of CACHD1 was among the most significant transcriptional features in primary tumors from patients with breast cancer. CACHD1 expression in primary tumors of the breast was correlated with post-progression survival in patients with luminal A disease, with mRNA levels of CACHD1 a negative prognostic indicator for luminal A breast cancer patients.

Discussion

Invasive breast cancer is a medical problem with a 27% 5-year survival rate for women whose disease has spread to distant sites^{5,6}. To facilitate understanding of the basic transcriptional differences between primary tumors of the breast and the tissues from which these tumors originate, normal breast tissues, we performed comparative transcriptome analysis using two independently published microarray datasets^{2,3}, providing evidence here that differential expression of cache domain containing 1, encoded by CACHD1, is a defining transcriptional feature of human breast cancer: in early onset-breast cancer, and in patients diagnosed after age 50. CACHD1 was expressed at significantly lower levels in primary tumors from patients with breast cancer as compared to normal breast tissue. Importantly, in patients with luminal

1 A breast cancer, expression of CACHD1 was correlated with post-progression survival. Cache domain
2 containing 1 and the molecular processes to which it pertains to may be relevant to the initiation or
3 progression of human breast cancer.
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

References

1. DeSantis, C.E., Ma, J., Goding Sauer, A., Newman, L.A. and Jemal, A., 2017. Breast cancer statistics, 2017, racial disparity in mortality by state. *CA: a cancer journal for clinicians*, 67(6), pp.439-448.
2. Chang, J.W., Kuo, W.H., Lin, C.M., Chen, W.L., Chan, S.H., Chiu, M.F., Chang, I.S., Jiang, S.S., Tsai, F.Y., Chen, C.H. and Huang, P.H., 2018. Wild-type p53 upregulates an early onset breast cancer-associated gene GAS7 to suppress metastasis via GAS7–CYFIP1- mediated signaling pathway. *Oncogene*, 37(30), pp.4137-4150.
3. Clarke, C., Madden, S.F., Doolan, P., Aherne, S.T., Joyce, H., O’driscoll, L., Gallagher, W.M., Hennessy, B.T., Moriarty, M., Crown, J. and Kennedy, S., 2013. Correlating transcriptional networks to breast cancer survival: a large-scale coexpression analysis. *Carcinogenesis*, 34(10), pp.2300-2308.
4. Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A. and Jemal, A., 2018. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*, 68(6), pp.394-424.
5. ACS Cancer Facts & Figures 2019.
<https://www.cancer.net/cancer-types/breast-cancer/metastatic/statistics>.
6. Survival Rates for Breast Cancer. <https://www.cancer.org/cancer/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-survival-rates.html>
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 105
Probe ID: 2340078
p-value: 1.46E-14
t: 1.06E+01
B: 22.96061
Gene: CACHD1
Gene name: cache domain containing 1

Chart 1: CACHD1 is differentially expressed in the primary tumors of patients with early-onset human breast cancer.

Rank of differential expression, probe ID, p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Rank: 4365
Probe ID: 225627_s_at
p-value: 7.73E-08
t: 5.73
B: 7.326281
Gene: CACHD1
Gene name: cache domain containing 1

Chart 2: CACHD1 is differentially expressed in the primary tumors of patients with breast cancer.

Rank of differential expression, probe ID, p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Figure 1: CACHD1 is expressed at significantly lower levels in primary breast tumors as compared to normal breast tissue.

The mRNA expression level of CACHD1 in normal breast tissue (left) and in primary tumors of the breast (right) is graphically depicted with the result of a statistical test evaluating significance of difference in CACHD1 expression between normal breast tissue and primary tumors of the breast, a *p*-value, is $p < 0.0001$.

Figure 2: CACHD1 expression correlates with post-progression survival in patients with luminal A subtype breast cancer, but not in patients with basal subtype, luminal B subtype, or HER2+ cancer.

Depicted in this Kaplan-Meier plot is the probability of post-progression survival (PPS) for $n=36$ patients with basal subtype breast cancer (upper left), $n=30$ patients with HER2+ breast cancer (upper right), $n=76$ patients with luminal A breast cancer (lower left), and $n=38$ patients with luminal B breast cancer (lower right), stratified into two groups, based on low or high expression of CACHD1, in patient primary tumors. The log rank p-value denoting statistical significance of difference in post-progression survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on CACHD1 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Luminal A subtype patients

Low CACHD1 expression: 38.67 months

High CACHD1 expression: 26.8 months

Chart 3: In luminal A subtype breast cancer, median post-progression survival is superior in patients with low primary tumor expression of CACHD1.

The post-progression survival of $n=76$ luminal A subtype breast cancer patients based on stratification into low or high tumor expression of CACHD1 is listed in this chart.