Nup160 is a differentially expressed gene in brain metastatic human breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with brain metastasis in humans with metastatic breast cancer. We found that nucleoporin 160, encoded by Nup160, was among the genes whose expression was most different in the brain metastases of patients with metastatic breast cancer as compared to primary tumors of the breast. Nup160 mRNA was present at decreased quantities in brain metastatic tissues as compared to primary tumors of the breast. Importantly, expression of Nup160 in primary tumors was correlated with patient overall survival. Modulation of Nup160 expression may be relevant to the biology by which tumor cells metastasize from the breast to the brain in humans with metastatic breast cancer.

Keywords: breast cancer, metastasis, brain metastasis, central nervous system metastasis, nucleoporin 160, Nup160, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with brain metastatic breast cancer^{4,5}. We discovered differential and decreased expression of the gene encoding nucleoporin 160, Nup160, in brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE52604⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=11$ primary breast tumors and $n=3$ brain metastases from patients with breast cancer; analysis was performed using platform GPL885. GSE52604 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer; analysis was performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether Nup160 gene expression was significantly different between primary tumors of the breast and brain metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of Nup160 mRNA expression levels with overall survival in $n=1402$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of brain metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

Nup160 is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified the nucleoporin 160, encoded by Nup160, as a differentially expressed gene in the breast metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, Nup160 ranked 93 out of 17418 total transcripts (Chart 1), equating to ___% differential expression. Differential expression of Nup160 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=1.82e-04$).

Differential gene expression analyses that utilize primary and metastatic tumor tissues to discover genes associated with metastasis can be challenging due to the relatively heterogeneous cellular composition and constitution of the tumor as compared to normal, untransformed (benign) tissues. Thus, to attempt to validate differential transcriptome-wide differential expression of Nup160 in brain metastases in human metastatic breast cancer, we queried a second microarray dataset⁵, here comparing normal breast tissues rather than primary tumors of the breast to brain metastases. Again, we identified Nup160 as a differentially expressed gene in the brain metastatic tissues of patients with breast cancer (Chart 2). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to normal breast tissues, Nup160 ranked 12684 out of 41093 total transcripts (Chart 2), equating to 80.1% differential expression. Differential expression of Nup160 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=1.41e-02$). Thus, differential expression of Nup160, transcriptome-wide, in the brain metastases of women with metastatic breast cancer was conserved across two independent microarray datasets, both when compared to benign and transformed breast tissues. This suggests that Nup160 differential expression in human brain metastatic breast cancer is a *bona fide* biological process.

Nup160 is expressed at lower levels in the brain metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for Nup160, in primary tumors of the breast and in brain metastasis of patients with brain metastatic breast cancer to determine direction and statistical significance of change in Nup160 expression in brain metastatic tissues. We observed significantly lower expression of Nup160 in the brain metastases of patients with breast cancer as compared to primary tumors of the breast: Nup160 was expressed at -0.21 ± 0.39 arbitrary units (AU) in primary tumors of the breast, while it was expressed at -1.50 ± 0.65 AU in brain metastatic tissues. The difference in Nup160 mRNA levels between primary tumors of the breast and brain metastatic tissues was statistically significant (Figure 1; $p=0.000366$).

Nup160 expression is correlated with survival outcomes in human breast cancer.

We performed Kaplan-Meier survival analysis⁷ in $n=1402$ breast cancer patients in total, to evaluate whether Nup160 tumor expression was correlated with survival outcomes in breast cancer. We observed a statistically significant correlation between primary tumor expression of Nup160 and overall survival (OS) in patients with breast cancer (Figure 2). Patients whose primary tumors expressed low levels of Nup160 possessed median OS of 89.04 months, while patients whose primary tumors expressed high levels of Nup160 possessed median OS of 173.72 months. This difference in OS based on Nup160 tumor expression in patients with breast cancer trended towards statistical significance (Figure 2, Chart 3; logrank p -value: 0.065; hazard ratio: 0.78 (0.6-1.02)).

Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified nucleoporin 160, encoded by Nup160, as among the genes whose expression was most different, transcriptome-wide, in the brain metastases of patients with breast cancer, both when compared to primary tumors of the breast and to normal breast tissues; we observed significantly decreased expression of Nup160 in brain metastases as compared to primary tumors of the breast. Further, we found a correlation between Nup160 expression and patient survival outcomes, as overall survival was greater in patients whose primary tumors expressed higher levels of Nup160 as compared to patients whose primary tumors expressed lower levels of Nup160.

Discussion

We provided evidence here that nucleoporin 160, encoded by Nup160, is among the genes whose expression is most different in the brain metastases of patients with brain metastatic breast cancer, that Nup160 mRNA is present at significantly decreased quantities in brain metastatic tissues as compared to primary tumors of the breast, and that primary tumor Nup160 expression is significantly correlated with patient survival outcomes in human breast cancer. Nucleoporin 160 is a subunit of the massive molecular assembly known as the nuclear pore complex which regulates transport into and out of the nucleus; Nup160 has functions in export of messenger RNAs and localizes to the nuclear basket⁸. Evaluation of the effects of genetic depletion of Nup160 in mouse models of metastatic breast cancer on metastasis to the central nervous system is merited. Modulation of Nup160 expression may be relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier and colonize the brain.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Salhia, B., Kiefer, J., Ross, J.T., Metapally, R., Martinez, R.A., Johnson, K.N., DiPerna, D.M., Paquette, K.M., Jung, S., Nasser, S. and Wallstrom, G., 2014. Integrated genomic and epigenomic analysis of breast cancer brain metastasis. *Plos one*, 9(1), p.e85448.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.
8. Vasu, S., Shah, S., Orjalo, A., Park, M., Fischer, W.H. and Forbes, D.J., 2001. Novel vertebrate nucleoporins Nup133 and Nup160 play a role in mRNA export. *The Journal of cell biology*, 155(3), pp.339-354.

Rank: 93
Probe ID: 13750
p-value: 1.82e-04
t: 4.86
B: 0.974684
Gene: Nup160
Gene name: nucleoporin 160

Chart 1: Nup160 is differentially expressed in brain metastatic breast cancer when comparing brain metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 12684
probe ID: A_24_P922606
p-value: 1.41e-02
t: -2.5500708
B: -3.662526
Gene: Nup160
Gene name: nucleoporin 160

Chart 2: Nup160 is differentially expressed in brain metastatic breast cancer when comparing brain metastases to normal breast tissues.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: Nup160 is expressed at significantly lower levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of Nup160 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in Nup160 expression between primary tumors of the breast and brain metastases, a p -value, is $p=0.000366$.

Figure 2: Correlation between Nup160 primary tumor expression and overall survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival for $n=1402$ total patients stratified into two groups, based on low or high expression of Nup160 in patient primary tumors. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on Nup160 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

1 Low Nup160 expression: 89.04 months
2 High Nup160 expression: 173.72 months

3
4 **Chart 3: Median overall survival is inferior in patients with low primary tumor expression of Nup160.**

5 The median OS (overall survival) of patients with low primary tumor expression of Nup160 and high
6 primary tumor expression of Nup160 is listed in this chart, for $n=1402$ breast cancer patients.