TYRO3 is differentially expressed in the tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab (Herceptin) is a monoclonal antibody targeting the extracellular domain of the human epidermal growth factor receptor 2 (HER2)¹ utilized for the treatment of adjuvant and metastatic breast cancer² in the United States and worldwide. We mined published microarray and gene expression data^{3,4} to discover in an unbiased manner the most striking transcriptional features of trastuzumab treatment. We identified the TAM family receptor protein tyrosine kinase TYRO3⁵ as among the genes most differentially expressed in the primary tumors of patients with breast cancer treated with trastuzumab. The primary tumors of breast cancer patients treated with trastuzumab expressed significantly higher levels of TYRO3 than did those of patients not treated with trastuzumab, implicating an additional cell surface receptor with tyrosine kinase capacity among the repertoire of genes whose expression is modulated following trastuzumab treatment.

Keywords: trastuzumab, breast cancer, TYRO3, systems biology of breast cancer, targeted therapeutics in breast cancer.

1 The mechanism of action ascribed to trastuzumab function is thought to be binding of the F_{ab}
2 domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2
3 (HER2)¹, but the transcriptional consequences resulting from signaling downstream of trastuzumab
4 binding to HER2 is less well understood. Through unbiased, systems-level comparative transcriptome
5 analysis of published microarray and gene expression data^{3,4}, we report here that TYRO3 is a
6 differentially expressed gene in the tumors of trastuzumab-treated patients: patients treated with
7 trastuzumab express significantly higher levels of TYRO3 than do those of patients not treated with
8 trastuzumab.

9 **Methods**

10 We used datasets GSE38057³ and GSE114082⁴ this differential gene expression analysis of the
11 transcriptional response to trastuzumab treatment in patients with breast cancer. GSE38057 was
12 generated using DASL Human Cancer Panel by Probe technology with $n=6$ primary breast tumors from
13 patients treated with induction chemotherapy and/or hormone therapy without trastuzumab and $n=35$
14 primary breast tumors from patients treated with induction chemotherapy and/or hormone therapy in
15 addition to trastuzumab, both of which developed brain metastases; analysis was performed using
16 platform GPL5826. GSE114082 was generated using Illumina HumanHT-12 WG-DASL V4.0 R2
17 expression beadchip technology with $n=17$ primary breast tumors from patients before trastuzumab
18 treatment and $n=17$ primary tumors from patients after trastuzumab treatment, analysis performed using
19 platform GPL14951. The Benjamini and Hochberg method of p -value adjustment was used for ranking
20 of differential expression but raw p -values were used to assess statistical significance of global or
21 multiplexed differential expression. Log-transformation of data was auto-detected, and the NCBI
22 generated category of platform annotation was used. A statistical test was performed to evaluate whether
23 TYRO3 gene expression was significantly between primary breast tumors of patients receiving
24 trastuzumab using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical
25 analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

26 **Results**

27 We harnessed the power of blind, comparative transcriptome analysis using published microarray
28 and gene expression data^{3,4} to identify the most significant transcriptional features of the primary breast
tumors of patients treated with trastuzumab (Herceptin).

29 **TYRO3 is differentially expressed in the tumors of patients treated with trastuzumab.**

30 We identified the TAM family receptor protein tyrosine kinase TYRO3 as among the genes
31 whose expression changed most significantly in the tumors of breast cancer patients treated with
32 trastuzumab (Table 1). When sorting each of the genes expressed in the primary tumors of breast cancer
33 patients treated with trastuzumab based on significance of change as compared to that of patients not
34 treated with trastuzumab, TYRO3 ranked 58 out of 1506 total transcripts (Table 1), equating to 96.1%
35 differential expression. TYRO3 differential expression in the tumors of patients treated with trastuzumab
36 was statistically significant (Table 1; $p=0.05411$).

37 The first dataset, although measuring the expression of over 1500 genes, was still limited. We
38 turned to a second microarray dataset, this measuring the expression of genes across the transcriptome -

29377 genes in total - but with another advantage. The tumors compared in this dataset were sampled from the same patient before and after a single, brief exposure to trastuzumab, not comparing tumors from different patients treated with a course of trastuzumab or not. Thus, while differentially expressed genes when comparing tumors based on trastuzumab treatment can result from a variety of sources, including gene expression changes resulting from tumor evolution over the course of treatment, changes identified in this dataset, occurring after only a single treatment of trastuzumab, are more likely to be a direct transcriptional result of trastuzumab binding events. We found again that TYRO3 was among the genes most differentially expressed in the tumors of patients administered a single treatment of trastuzumab (Table 2). When sorting each of the genes expressed in the tumors of patients administered a single treatment of trastuzumab, TYRO3 ranked 3632 out of 29377 total transcripts (Table 2), equating to 87.6% differential expression. Differential expression of TYRO3 in the primary tumors of breast cancer patients following treatment with trastuzumab was statistically significant (Table 2; $p=2.5E-03$).

TYRO3 is expressed at significantly higher levels in the tumors of breast cancer patients treated with trastuzumab than in the tumors of those not treated with trastuzumab.

We obtained exact mRNA expression levels for TYRO3 from tumors of patients treated with trastuzumab and from the tumors of patients not treated with trastuzumab to directly evaluate magnitude and change of expression in TYRO3 based on history of trastuzumab treatment. TYRO3 was expressed at significantly higher levels in the tumors of patients treated with trastuzumab as compared to those of patients not treated with trastuzumab (Figure 1; $p=0.0134$). We calculated a mean fold change of 1.75 ± 0.93 in TYRO3 expression in the primary tumors of patients treated with trastuzumab relative to that of patients not treated with trastuzumab.

Thus, we found that TYRO3 was among the genes most differentially expressed in the tumors of patients treated with trastuzumab, that TYRO3 was expressed at significantly higher levels in the tumors of patients treated with trastuzumab as compared to those not treated with trastuzumab, and that this transcriptional change was likely a direct result of trastuzumab treatment as opposed to a gene expression change associated with tumor evolution.

Discussion

TYRO3⁵ is a member of the TAM receptor tyrosine kinases (RTKs) that include MERTK and AXL; these RTKs are expressed in the brains of both mice and rats^{5,6} and they possess an extracellular domain that bears resemblance to that of cell adhesion molecules expressed in the nervous system⁶. In the central nervous system of the rat, the expression pattern of TYRO3 during development mirrors that of synaptogenesis⁶. Both Gas6 and the anti-coagulation factor protein S have been reported as ligands for TYRO3^{7,8}. TYRO3, MERTK and AXL share approximately 35% homology and their extracellular domains contain two immunoglobulin-like domains as well as two fibronectin type III repeats⁷. One study reported that depletion of TYRO3 using small interfering RNA in a number of breast cancer cell lines in vitro resulted in impairment of cell proliferation and cell cycle arrest⁹.

We provided evidence here that TAM family receptor tyrosine kinase TYRO3, a cell surface receptor broadly and most abundantly expressed in the mammalian central nervous system^{5,6}, is a differentially expressed gene in the tumors of patients treated with trastuzumab, and that breast cancer patients treated with trastuzumab expressed significantly higher levels of TYRO3 mRNA in their primary tumors than did patients not treated with trastuzumab.

References

1. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
2. Herceptin (Trastuzumab). Indications. <https://www.herceptin.com/hcp/indications.html>.
3. Duchnowska, R., Jassem, J., Goswami, C.P., Dundar, M., Gökmen-Polar, Y., Li, L., Woditschka, S., Biernat, W., Sosińska-Mielcarek, K., Czartoryska-Arłukowicz, B. and Radecka, B., 2015. Predicting early brain metastases based on clinicopathological factors and gene expression analysis in advanced HER2-positive breast cancer patients. *Journal of Neuro-oncology*, 122(1), pp.205-216.
4. Triulzi, T., Regondi, V., De Cecco, L., Cappelletti, M.R., Di Modica, M., Paolini, B., Lollini, P.L., Di Cosimo, S., Sfondrini, L., Generali, D. and Tagliabue, E., 2018. Early immune modulation by single-agent trastuzumab as a marker of trastuzumab benefit. *British journal of cancer*, 119(12), pp. 1487-1494.
5. Schulz, N.T., Paulhiac, C.I., Lee, L. and Zhou, R., 1995. Isolation and expression analysis of tyro3, a murine growth factor receptor tyrosine kinase preferentially expressed in adult brain. *Molecular brain research*, 28(2), pp.273-280.
6. Prieto, A.L., Weber, J.L. and Lai, C., 2000. Expression of the receptor protein-tyrosine kinases Tyro-3, Axl, and Mer in the developing rat central nervous system. *Journal of comparative neurology*, 425(2), pp.295-314.
7. Stitt, T.N., Conn, G., Goret, M., Lai, C., Bruno, J., Radzilewski, C., Mattsson, K., Fisher, J., Gies, D.R., Jones, P.F. and Masiakowski, P., 1995. The anticoagulation factor protein S and its relative, Gas6, are ligands for the Tyro 3/Axl family of receptor tyrosine kinases. *Cell*, 80(4), pp.661-670.
8. Godowski, P.J., Mark, M.R., Chen, J., Sadick, M.D., Raab, H. and Hammonds, R.G., 1995. Reevaluation of the roles of protein S and Gas6 as ligands for the receptor tyrosine kinase Rse/Tyro 3. *Cell*, 82(3), pp.355-358.
9. Ekyalongo, R.C., Mukohara, T., Funakoshi, Y., Tomioka, H., Kataoka, Y., Shimono, Y., Chayahara, N., Toyoda, M., Kiyota, N. and Minami, H., 2014. TYRO3 as a potential therapeutic target in breast cancer. *Anticancer research*, 34(7), pp.3337-3345.

Rank	ID	p-value	t	B	FC	Gene
58	GI_27597077-S-5	0.05411	1.980192	-4.37	1.75 ± 0.93	NM_006293.2

Table 1: TYRO3 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in TYRO3 between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Rank	ID	p-value	t	B	Gene
3632	ILMN_1740169	2.5E-03	3.2510661	-1.71807	TYRO3

Table 2: A single, brief exposure to trastuzumab is sufficient to drive differential expression of TYRO3 in the primary tumors of patients with breast cancer.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: TYRO3 is expressed at significantly higher levels in the primary tumors of patients treated with trastuzumab

The mRNA expression level of TYRO3 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between TYRO3 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.