Protective and Resilience Factors to Promote Mental Health among Indigenous Youth in Canada: A Scoping Review Protocol

Leah Carrier^a; Hwayeon Danielle Shin^b ; Melissa A. Rothfus^c; Janet A. Curran^d

^a Doctoral Student, School of Nursing, Dalhousie University, Halifax Canada.
<https://orcid.org/0000-0001-7632-2629>

^b Doctoral Student, Institute of Health Policy, Management and Evaluation, University of Toronto, Ontario, Canada.
<https://orcid.org/0000-0003-4037-4464>

^c Librarian Scientist, W.K. Kellogg Health Sciences Library, Dalhousie University, Halifax, Canada

^d Professor, School of Nursing, Dalhousie University, Halifax Canada. Affiliate Scientist, IWK Health Centre, Halifax, NS.

Correspondence to:

Hwayeon Danielle Shin
hdanielle.shin@mail.utoronto.ca
Institute of Health Policy, Management and Evaluation,
University of Toronto
155 College St 4th Floor, Toronto,
ON Canada M5T 3M6

Conflicts of Interest

There are no conflicts of interest to declare with this submission.

Word count

3000

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sector.

Contributors

LC and HDS designed the scoping review protocol including data collection and interpretation planning and drafted and revised the protocol for intellectual content. LC, HDS, MAR, JAC made a final approval for the submission of the protocol. LC and HDS made an equal contribution to the manuscript and claim first co-authorships.

Acknowledgement

We would also like to thank Dr. Margot Latimer for reviewing an earlier draft of this protocol

ABSTRACT

Introduction: Many mental health concerns emerge in adolescence and young adulthood, making this a critical period to initiate early interventions for mental health promotion and illness prevention. Although Indigenous young people in Canada are at a higher risk of mental health outcomes and faced with limited access to appropriate care and resources, they have unique strengths and resilience that promote mental health and wellness. Furthermore, resilience has been described as a ‘healing journey’ by Indigenous peoples, and interventions that account for the culture of these groups show promise in promoting mental health and wellness. As such, there is a need for innovative mental health interventions for Indigenous youth that transcend the Western biomedical model, use a strengths-based approach, and account for the cultural practices and belief systems of Indigenous peoples. This scoping review aims to explore the resilience and protective factors that promote mental health and wellness for Indigenous youth in present-day Canada with the aim of compiling and summarizing the available literature on this topic to date.

Methods and Analysis: The review will follow Joanna Briggs Institute methodology for conducting scoping reviews. The reporting will follow the Preferred Reporting Items for Systematic Reviews and Meta-analysis Extension for Scoping Reviews (PRISMA-ScR) checklist and guidelines. We will include both published and unpublished grey literature and search the following databases: PubMed, CINAHL, PsycInfo, ERIC, Embase and Scopus. The search of all databases was conducted on August 26, 2021. Further, we will use government and relevant Indigenous organization websites. Two reviewers will independently screen and select the articles and extract the data.

Ethics and Dissemination: No ethical approval is required for this study. We will share the results through conference presentations and an open-access publication in a peer-reviewed journal. A lay-language report will be created and disseminated to community organizations that work with Indigenous youth.

Key words

Indigenous Youth, Resilience, Protective Factors, Mental Health, Mental Wellness

Strengths and limitations of this study

- While there are numerous studies illustrating the risk factors for mental health concerns among Indigenous youth, this is a novel and timely topic to comprehensively identify the range of resilience and protective factors that promote mental health and wellness.
- The study selection, data extraction and charting will be performed by two independent reviewers to minimize the risk of bias or errors.
- This scoping review has limitations regarding the language and geography, and the findings of this review may be relevant in countries of a similar context.
- Since the purpose of our scoping review is to map and characterize the evidence, we will not be conducting critically appraisals to determine the quality of individual studies to assess the risk of bias.

INTRODUCTION

Although Indigenous children and youth face significant health inequities when compared with the rest of the Canadian population^{1,2}, they have inherent strengths and resilience that can be drawn upon to promote health and wellness. A review of critiques and recommendations for interventions promoting healthy youth development suggests that there is a need to focus on reducing risk and strengthening protective factors³. Risk factors for adverse mental health outcomes include poverty, family conflict, neglect in early childhood, and unsafe housing⁴. Fortunately, protective factors can positively modify a person's response to external and environmental stressors that result in maladaptive outcomes⁵. In other words, protective factors can minimize the likelihood of dysfunction and disorder in the presence of risks. Identified protective factors are, but not limited to, living in a safe community, living free of violence, supportive relationships, and education⁴. Protective factors are tightly connected to resilience, which is defined as one's capacity to respond and overcome risks in order to avoid negative outcomes^{6,7}. Furthermore, resilience has also been conceptualized as function of situational and contextual factors⁸ or a dynamic process⁹ that contribute to positive outcomes. A protective model of resilience explains how protective factors operate to influence the trajectory from risk exposure to adverse outcome and modify the impact of risk exposure^{5,6}. For example, a longitudinal study of homeless youth found that physical abuse was a risk factor for greater depressive symptoms¹⁰. However, having mentors, family, and friends that youth could rely on acted as a protective factor resulting in positive social support, which then lowered the risk for depressive symptoms¹⁰. In recognition of critical role that resilience and protective factors can play, research is needed to explore these factors for effective mental health promotion.

Resilience initiatives among Indigenous populations must address the trauma and cultural loss that occurs due to colonization^{11–13}. For example, the Cree have described resilience as a “healing journey”^{14(p790)}. Resilience identified by Indigenous youth in Canada (e.g., Metis and Inuit) includes, but are not limited to, coping strategies and health promoting practices that benefit youth across varying levels of adversity^{15,16}. Research among Indigenous youth and adults in Canada suggests that protective factors for mental wellness include culture, community, and family^{17–19}. For example, mental health interventions that account for the culture of Indigenous peoples show particular promise, culture acting as a protective factor, because cultural identity is foundational to holistic health outcomes and is nurtured through participation in cultural activities^{20,21}.

Mental health promotion and innovative approaches to promoting Indigenous youth mental health is a critical issue in Canada as mental health concerns are prevalent among youth, and various research has highlighted the high frequency of delayed or disrupted access to mental health services^{22,23}. Indigenous young people experience higher rates mental health concerns and more limited access to appropriate care and resources when compared to their settler Canadian counterparts²⁴. For example, suicide is the second leading cause of death among all youth in Canada²⁵, which has been consistent for the past 10 years²⁶; this finding is even more dire when one considers that suicide rates among Indigenous youth are 4–10 times higher than the Canadian average²⁷. Strikingly, a recent Canadian study found that First Nations youth had an elevated risk of suicidal ideation and attempts if they had a parent or grandparent who attended residential school¹⁸. Pain has been conceptualized from a mental and emotional perspective in research conducted with Indigenous youth in Mi’kma’ki (also known as Atlantic Canada)²⁸. Additionally, First Nations children and youth have higher rates of acute and chronic pain^{29–31}, mental illness,

and substance use disorders¹⁹. The intergenerational effect of residential school trauma and its impacts on multiple dimensions of mental health is stronger among younger youth, illustrating the importance of early intervention to promote mental health in this population^{17,18}. Children who have mental health problems are more likely to experience continued challenges to mental health problems in adolescence and adulthood³². Moreover, many mental health disorders have their onset in adolescence and young adulthood³³, making this a critical period to initiate early interventions for mental health promotion and illness prevention³⁴.

There is a demonstrated need for innovative mental health interventions for Indigenous youth that transcend the Western biomedical model, use a strengths-based approach, and account for the cultural practices and belief systems of Indigenous peoples^{16,35}. A systematic review done by Rowhani and Hatala¹⁶ included 17 peer-reviewed articles and investigated how resilience of Indigenous youth is referenced in the literature and implemented in Canada. They found that these 17 articles commonly included a collaborative research approach and demonstrated diversity in culture, language, and environment across the Indigenous communities in Canada¹⁶. Furthermore, major factors contributing to resilience among these groups included cultural continuity, community and family ties and empowerment¹⁶. Building on this previous work and specifically adding in the aspects of mental health and wellness, this current scoping review's primary objective is to explore how existing literature identifies resilience or protective factors that promote mental health and mental wellness among Indigenous youth in Canada. Furthermore, we hope to include all types of evidence that pertains to resilience and protective factors for Indigenous youth mental health. This holistic understanding of evidence is particularly useful in the current scoping review topic because Indigenous knowledge also exists in the form of grey literature. It is common for Indigenous writers to use personal narratives in

describing a ‘healing journey’¹⁴. For example, community knowledge related to mental health can differ from Western understandings of individual protective or resiliency factors, with a critical lens focused on the systems and structural processes that affect the health of Indigenous youth³⁶. While Western literature may focus on the individual factors that can be strengthened or supported with mental health interventions, it is important to reflect on whether this is the best place to focus efforts when there are structural barriers to health equity for Indigenous youth that need to be addressed ³⁶. Therefore, we will be including diverse evidence that systematic reviews would traditionally exclude, in addition to the published work about Indigenous knowledge in this area.

It is important to note that research has been historically conducted *on* Indigenous Peoples rather than *in collaboration with* them. As a result, there has previously been a failure to prioritize cultural conceptualizations and appropriate portrayals of Indigenous Peoples, as reflected in much of the published literature. This has resulted in detrimental practices associated with the ownership, utilization, and understanding of Indigenous knowledge.³⁷ However, it is important to note that there is a growing body of literature from research done in collaboration with and led by Indigenous communities in recent years^{36,38–42}. For example, Kirmayer and colleagues³⁶ discuss the concept of community resilience and how this aligns with Indigenous values and beliefs about how a person is interconnected with their environment and other people. This work defines the concept of community resilience, which may look at how individuals overcome challenges with the help of social or cultural resources in their communities. Community resilience may also refer to how communities as a whole demonstrate resiliency in the face of challenges. In another paper, Kirmayer and co-authors describe findings from resilience research with Inuit, Metis, Mi’kmaq, and Mohawk communities, describing resilience

sources as a dynamic process that comes from relationships and interactions between people and their communities and social systems³⁸. Recognizing that community involvement and priorities are critical for research related to Indigenous health and that community conceptualizations of resilience and mental health promotion may differ from Western conceptualizations, the secondary objective of this review is to identify and report the level of community involvement described in the included studies that pertain to resilience or protective factors for the mental health of Indigenous youth.

A preliminary search of the *JB**I Database of Systematic Reviews and Implementation Reports*, the Cochrane Database of Systematic Reviews, PubMed, CINAHL, and Embase was conducted and no current or underway scoping or systematic reviews on the topic were identified.

REVIEW QUESTIONS

1. What does the literature identify as resilience or protective factors for the mental health and well-being of Indigenous youth (ages 10-25) in Canada?
 - a. What is the level of community involvement reported in the included studies and how is community involvement described?

INCLUSION CRITERIA

Participants

The review will consider studies that include self-identified Indigenous youth in Canada who are between the ages of 10 and 25 years. This age bracket is consistent with how many Indigenous communities define ‘youth’ and with definitions used in current mental health literature^{43,44}. Articles will be included where participants have an official mental health diagnoses (DSM-5⁴⁵), self-identified mental health and wellness concerns, symptoms of

psychological distress, and/or are involved in mental health promotion and illness prevention initiatives. All genders will be included.

Concept

This scoping review will include all articles that examine resilience and/or protective factors for mental health and mental wellness among Indigenous youth. Articles that examine prevalence and incidence of mental health concerns without discussing resilience or protective factors will be excluded. For this review, the following operational terms and definitions will be used (Table 1). Articles may not explicitly use the terms, ‘resilience’ or ‘protective factors’. As such, this review will include articles if they discuss other mental health promoting factors and psychosocial factors that contribute to adaptive outcomes for mental health and wellness, all of which fall under the operationalized definitions for resilience or protective factors.

Table 1. Operational terms and definitions

Term	Definition
Protective factors	“Influences that modify, ameliorate, or alter a person's response to some environmental hazard that predisposes to a maladaptive outcome”. ^{5(p600)}
Resilience	Personality traits ⁴⁶ , function of situational and contextual factors ⁸ or a dynamic process ⁹ that contribute to positive mental health outcomes.
Youth	Individuals with ages of 10 and 25 years. ^{43,44}

Context

This review will consider articles that are conducted in the health care, education, or community settings in Canada. International articles where the data related to Indigenous youth in Canada can be extracted will also be included. Settings may be urban, rural, or remote. Articles with Indigenous communities or Indigenous participants who live in settler communities

will be included. Articles will be excluded when Canadian data is not clearly identified or able to be extracted.

Types of Sources

This scoping review will consider both quantitative and qualitative designs, as well as those studies and articles that use Indigenous ways of knowing. Quantitative designs include any experimental and quasi-experimental study designs including randomized controlled trials, non-randomized controlled trials, before and after studies and interrupted time-series studies. In addition, analytical observational studies including prospective and retrospective cohort studies, case-control studies and analytical cross-sectional studies will be considered for inclusion. This review will also consider descriptive observational study designs including case series, individual case reports and descriptive cross-sectional studies for inclusion. Qualitative studies will also be considered that focus on qualitative data including, but not limited to, designs such as phenomenology, grounded theory, ethnography, qualitative description, action research and feminist research. In addition, scoping and systematic reviews that meet the inclusion criteria will also be considered and their references lists will be reviewed for primary articles. Text and opinion papers will also be considered for inclusion in this scoping review. Grey literature will also be considered, and may include papers, reports, or policy documents related to youth mental health promotion and protective factors. Studies in English will be included. No parameters around dates will be applied.

METHODS

Scoping reviews determine the scope of existing evidence on a specific topic and identify key concepts and research gaps^{47,48}. The findings of the reviews can be used to inform future research that is focused on important knowledge gaps⁴⁹. The findings can also provide critical

information for decision makers, policy makers, and clinicians in the area of Indigenous youth mental health. Therefore, a scoping review is an appropriate methodology to systematically explore all literature including peer-reviewed articles and grey literature about resilience and protective factors for Indigenous youth's mental health. To ensure replicability and rigorous methodology, the proposed scoping review will strictly adhere to the Joanna Briggs Institute methodology for scoping reviews^{50,51}. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for scoping reviews (PRISMA-ScR) checklist will be used to guide the reporting of this scoping review⁵². Major steps comprised in our scoping review methods are: 1) identifying the research question; 2) searching for relevant studies; 3) screening and selecting relevant studies; 4) charting data; and 5) summarizing and presenting key findings.

Search Strategy

The final search strategy and data search was developed in consultation with a JBI-trained librarian scientist. A preliminary search of PubMed and Google Scholar was undertaken to identify target articles on the topic. The text words contained in the titles and abstracts of relevant articles, the index terms, headings and MESH terms used to describe the articles were used to develop the final search strategy for PubMed. The search strategy, including all identified keywords, headings, MESH terms, and index terms, was adapted for each included information source. The final search strategies conducted on August 26, 2021 for all included databases can be found in Appendix 1. A second librarian will complete a peer review of the final search strategies using the Peer Review of Electronic Search Strategies (PRESS) guidelines.⁵³

Information Sources

PubMed, Cumulative Index of Nursing and Allied Health Literature, Education Resources Information Center (ERIC), Embase, PsycInfo and Scopus will be included as

information sources. The search for unpublished literature will include: Grey Literature Report, conference abstracts and presentations and dissertations, government documents and relevant Indigenous organization websites including, but not limited to Native Women's Association of Canada, Assembly of First Nations, and the First Nations Inuit Governance Centre. Lastly, a hand search of three high impact journals (Canadian Journal of Community Mental Health, Journal of Psychiatric and Mental Health Nursing, and International Journal of Mental Health Nursing) will be carried out.

Study Selection

Following the search, all identified citations will be collated and uploaded into Covidence, an online software program⁵⁴, and duplicates will be removed. Titles and abstracts will then be screened by two independent reviewers for assessment against the inclusion criteria for the review. Potentially relevant studies will be retrieved in full and their citation details imported into Covidence. The full text of selected citations will be assessed in detail against the inclusion criteria by two independent reviewers. Reasons for exclusion of full text studies that do not meet the inclusion criteria will be recorded and reported in the full scoping review. Any discrepancies between the reviewers at each stage of the study selection process will be resolved through discussion, or with a third reviewer. Scoping reviews do not require methodological assessment,⁵⁰ so critical appraisal will not be conducted. The results of the search will be reported and presented in a PRISMA 2020 flow diagram⁵⁵.

Data Charting

A draft of the data extraction instrument (Appendix 2) has been developed by the reviewers to extract the following study information:

- Authors

- 287 • Year of publication
- 288 • Aim of study
- 289 • Type of study
- 290 • Literature type
- 291 • Country of publication
- 292 • Study population (participants)
- 293 • Geographical setting
- 294 • Type of service (healthcare, education, community, etc.)
- 295 • Description of community involvement
- 296 • Involved community, knowledge users, or stakeholders
- 297 • Type of mental health concerns
- 298 • Outcome measurement tool
- 299 • Methodology/methods
- 300 • Use of a theory or framework
- 301 • Definition of resilience/protective factors/coping skills used
- 302 • Description of resilience or protective factors (this can also include other mental health
- 303 promoting factors or psychosocial factors that contribute to adaptive outcomes for mental
- 304 health and wellness)
- 305 • Reported outcomes
- 306 • Study main findings and conclusion
- 307 • Study future implications
- 308 Two reviewers will independently extract data using the developed data extraction tool.
- 309 Reviewers will meet after data extraction has been completed for the first three studies to

identify any discrepancies and ensure consistency. The draft data extraction tool will be modified and revised as necessary during the process of extracting data from each included study. Modifications and revisions will be detailed in the full scoping review report. After finalizing data extraction tool, the remaining data will be extracted by two independent reviewers. Any conflicts in the data extraction will be resolved through discussion, or with a third reviewer. Authors of papers will be contacted to request missing or additional data, where required.

Drawing on Bronfenbrenner's ecological systems theory⁵⁶ as well as Indigenous knowledge about the interacting systems that influence behaviour, narrative descriptions of resilience and protective factors will be mapped on the appropriate level of ecological system (individual, family, community, cultural, and systems level). Secondly, community involvement described in the identified citations will be characterized appropriately. As data analysis is expected to be an iterative process, necessary changes will be made and reported in the full review.

Data Presentation

The extracted data will be, where appropriate, presented in a tabular form with the narratives in a manner that aligns with the objective of this scoping review. The tabular summary will highlight the major categories of the data extraction form as described above. A narrative summary will accompany the tabulated and/or charted results and will describe how the results relate to the reviews objective and questions.

A narrative description of protective factors identified by the studies and community definitions of protective factors will be provided to discuss apparent gaps in current knowledge. The use of theory and framework to understand the protective factors for Indigenous communities will also be reported, which will help illuminate contextual factors that may influence the use and implementation of future interventions in this area.

333 **Patient and Public Involvement**

334 Patients or the public were not involved in this scoping review design, including data
335 collection and interpretation planning. However, a lay-language report will be created and
336 disseminated to community organizations that work with Indigenous youth.

337 **Ethics and Dissemination**

338 This is a scoping review on existing literature, and therefore, no ethical approval is
339 required for this study. The goal of this scoping review is to map and characterise the evidence
340 related to resilience and protective factors for mental health and wellness among Indigenous
341 Youth in Canada. The findings of the scoping review may inform future research directions as
342 well as mental health intervention designs that account for the cultural and belief systems of
343 Indigenous peoples. As such, not only we will share the results through conference presentations
344 and an open-access publication in a peer-reviewed journal but also share with key academic and
345 community organizations (e.g., Indigenous Health Interest Group at Dalhousie University,
346 Friendship Centres, community health centres) and health systems through lay-language
347 summaries and informal presentations.

REFERENCES

1. Smylie J, Fell D, Ohlsson A, Joint Working Group on First Nations I Inuit, and Métis Infant Mortality of the Canadian Perinatal Surveillance System. A Review of Aboriginal Infant Mortality Rates in Canada: Striking and Persistent Aboriginal/Non-Aboriginal Inequities. *Can J Public Health*. 2010;101(2):143-148. doi:10.1007/BF03404361
2. Peters PA, Oliver LN, Kohen DE. Mortality among children and youth in high-percentage First Nations identity areas, 2000-2002 and 2005-2007. *Rural Remote Health*. 2013;13(3):2424-2424.
3. Catalano RF, Hawkins JD, Berglund ML, Pollard JA, Arthur MW. Prevention science and positive youth development: competitive or cooperative frameworks? *J Adolesc Health*. 2002;31(6):230-239. doi:10.1016/S1054-139X(02)00496-2
4. Government of Canada. Protective and risk factors for mental health. Published 2019. <https://www.canada.ca/en/public-health/services/protective-risk-factors-mental-health.html>
5. Rutter M. Resilience in the face of adversity. Protective factors and resistance to psychiatric disorder. *Br J Psychiatry J Ment Sci*. 1985;147:598-611. doi:10.1192/bjp.147.6.598
6. Rutter M. Psychosocial Resilience and Protective Mechanisms. *Am J Orthopsychiatry*. 1987;57(3):316-331. doi:10.1111/j.1939-0025.1987.tb03541.x
7. Rak CF, Patterson LE. Promoting resilience in at-risk children. *J Couns Dev*. 1996;74(4):368-373. doi:10.1002/j.1556-6676.1996.tb01881.x
8. Masten AS, Best KM, Garmezy N. Resilience and development: Contributions from the study of children who overcome adversity. *Dev Psychopathol*. 1990;2(4):425-444. doi:10.1017/S0954579400005812
9. Luthar SS, Cicchetti D, Becker B. The Construct of Resilience: A Critical Evaluation and Guidelines for Future Work. *Child Dev*. 2000;71(3):543-562. doi:10.1111/1467-8624.00164
10. Tyler KA, Schmitz RM, Ray CM. Role of Social Environmental Protective Factors on Anxiety and Depressive Symptoms Among Midwestern Homeless Youth. *J Res Adolesc*. 2018;28(1):199-210. doi:10.1111/jora.12326
11. Brooks CM, Daschuk MD, Poudrier J, Almond N. First Nations youth redefine resilience: listening to artistic productions of 'Thug Life' and hip-hop. *J Youth Stud*. 2015;18(6):706-725. doi:10.1080/13676261.2014.992322
12. Goulet L, Linds W, Episknew J-A, Schmidt K. Creating a Space for Decolonization: Health through Theatre with Indigenous Youth. *Native Stud Rev*. 2011;20(1):89-116.
13. Michel Tousignant, Nibisha Sioui. Resilience and Aboriginal Communities in Crisis: Theory and Interventions. *J Aborig Health*. 2009;5(1):43.

14. Isaak CA, Stewart DE, Mota NP, Munro G, Katz LY, Sareen J. Surviving, healing and moving forward: Journeys towards resilience among Canadian Cree adults. *Int J Soc Psychiatry*. 2015;61(8):788-795. doi:10.1177/0020764015584648
15. Kral MJ, Salusky I, Inuksuk P, Angutimarik L, Tulugardjuk N. Tunngajuq: Stress and resilience among Inuit youth in Nunavut, Canada. *Transcult Psychiatry*. 2014;51(5):673-692. doi:10.1177/1363461514533001
16. Rowhani M, Hatala A. A Systematic Review of Resilience Research among Indigenous Youth in Contemporary Canadian Contexts. *Int J Health Wellness Soc*. 2017;7:45-58. doi:10.18848/2156-8960/CGP/v07i04/45-58
17. Bombay A, Matheson K, Anisman H. Decomposing identity: Differential relationships between several aspects of ethnic identity and the negative effects of perceived discrimination among First Nations adults in Canada. *Cultur Divers Ethnic Minor Psychol*. 2010;16(4):507-516. doi:10.1037/a0021373
18. Bombay A, McQuaid RJ, Schwartz F, Thomas A, Anisman H, Matheson K. Suicidal thoughts and attempts in First Nations communities: links to parental Indian residential school attendance across development. *J Dev Orig Health Dis*. 2019;10(1):123-131. doi:10.1017/S2040174418000405
19. Health Canada and Assembly of First Nations. *First Nations Mental Wellness Continuum Framework*. Ottawa: Health Canada.; 2015. Accessed January 18, 2021. https://thunderbirdpf.org/wp-content/uploads/2015/01/24-14-1273-FN-Mental-Wellness-Framework-EN05_low.pdf
20. Reyhner J. Indigenous Language Immersion Schools for Strong Indigenous Identities. *Herit Lang J*. 2010;7(2):138-152.
21. LaFromboise TD, Hoyt DR, Oliver L, Whitbeck LB. Family, community, and school influences on resilience among American Indian adolescents in the upper midwest. *J Community Psychol*. 2006;34(2):193-209. doi:10.1002/jcop.20090
22. Gravel R, Béland Y. The Canadian Community Health Survey: Mental Health and Well-Being. *Can J Psychiatry*. 2005;50(10):573-579. doi:10.1177/070674370505001002
23. Canadian Mental Health Association. Fast Facts about Mental Illness - CMHA National. Accessed February 4, 2020. <https://cmha.ca/fast-facts-about-mental-illness>
24. Statistics Canada. Aboriginal peoples in Canada: First Nations people, Metis and Inuit: National Household Survey, 2011. Published May 8, 2013. Accessed January 18, 2021. <https://www150.statcan.gc.ca/n1/daily-quotidien/130508/dq130508a-eng.htm>
25. Kutcher SP, Szumilas M. Youth suicide prevention. *CMAJ Can Med Assoc J J Assoc Medicales Can*. 2008;178(3):282-285. doi:10.1503/cmaj.071315

26. Public Health Agency of Canada. Suicide in Canada: Key Statistics (infographic). Government of Canada. Published August 1, 2019. Accessed September 28, 2019. <https://www.canada.ca/en/public-health/services/publications/healthy-living/suicide-canada-key-statistics-infographic.html>
27. Government of Canada SC. Suicide among First Nations people, Métis and Inuit (2011-2016): Findings from the 2011 Canadian Census Health and Environment Cohort (CanCHEC). Published June 28, 2019. Accessed April 25, 2021. <https://www150.statcan.gc.ca/n1/pub/99-011-x/99-011-x2019001-eng.htm>
28. Latimer M, Sylliboy JR, Francis J, et al. Co-creating better healthcare experiences for First Nations children and youth: The FIRST approach emerges from Two-Eyed seeing. *Paediatr Neonatal Pain*. 2020;2(4):104-112. doi:10.1002/pne2.12024
29. Latimer M, Simandl D, Finley A, et al. Understanding the Impact of the Pain Experience on Aboriginal Children's Wellbeing: Viewing Through a Two-Eyed Seeing Lens. *First Peoples Child Fam Rev*. 2014;9(1):22-37.
30. Latimer M, Rudderham S, Lethbridge L, et al. Occurrence of and referral to specialists for pain-related diagnoses in First Nations and non-First Nations children and youth. *CMAJ Can Med Assoc J J Assoc Medicale Can*. 2018;190(49):E1434-E1440. doi:10.1503/cmaj.180198
31. Latimer M, Sylliboy JR, MacLeod E, et al. Creating a safe space for First Nations youth to share their pain. *PAIN Rep*. 2018;3. https://journals.lww.com/painrpts/Fulltext/2018/09001/Creating_a_safe_space_for_First_Nations_youth_to.6.aspx
32. Smetanin P, Stiff D, Briante C, Adair CE, Ahmed S, Khan M. The Life and Economic Impact of Major Mental Illnesses in Canada: 2011 to 2041. Published online 2011.
33. Jones P. Adult mental health disorders and their age at onset. *Br J Psychiatry*. 2013;202(s54):s5-s10. doi:10.1192/bjp.bp.112.119164
34. Mental Health Commission of Canada. Changing directions, changing lives: The mental health strategy for Canada. Published online 2012. https://www.mentalhealthcommission.ca/sites/default/files/MHStrategy_Strategy_ENG.pdf
35. Sasakamoose J, Scerbe A, Wenaus I, Scandrett A. First Nation and Métis Youth Perspectives of Health: An Indigenous Qualitative Inquiry. *Qual Inq*. 2016;22(8):636-650. doi:10.1177/1077800416629695
36. Kirmayer LJ MD, FRCPC, Sehdev M, Whitley R PhD, Dandeneau SF PhD, Isaac C. Community Resilience: Models, Metaphors and Measures. *J Aborig Health*. 2009;5(1):62-117.
37. First Nations Information Governance Centre. *Ownership, Control, Access and Possession (OCAP™): The Path to First Nations Information Governance.*; 2014. Accessed August 2,

2021. <http://books.scholarsportal.info/en/read?id=/ebooks/ebooks4/cpdc4/2018-12-11/3/10095457>

38. Kirmayer LJ, Dandeneau S, Marshall E, Phillips MK, Williamson KJ. Rethinking Resilience from Indigenous Perspectives. *Can J Psychiatry*. 2011;56(2):84-91. doi:10.1177/070674371105600203
39. Andersson N, Ledogar RJ. The CIET Aboriginal Youth Resilience Studies: 14 Years of Capacity Building and Methods Development in Canada. *Pimatisiwin*. 2008;6(2):65-88.
40. Tousignant M PhD, Sioui N PhD. Resilience and Aboriginal Communities in Crisis: Theory and Interventions. *J Aborig Health*. 2009;5(1):43-61.
41. Njeze C, Bird-Naytowhow K, Pearl T, Hatala AR. Intersectionality of Resilience: A Strengths-Based Case Study Approach With Indigenous Youth in an Urban Canadian Context. *Qual Health Res*. 2020;30(13):2001-2018. doi:10.1177/1049732320940702
42. Isbister-Bear O, Hatala A, Sjoblom E. Strengthening Âhkamêyimo among Indigenous youth: The social determinants of health, justice, and resilience in Canada's north. *J Indig Well-Being*. 2017;2:76-89.
43. World Health Organization. Atlas : child and adolescent mental health resources : global concerns: implications for the future. *Child Adolesc Ment Health Atlas*. Published online 2005. <https://apps.who.int/iris/handle/10665/43307>
44. Malla A, Shah J, Iyer S, et al. Youth Mental Health Should Be a Top Priority for Health Care in Canada. *Can J Psychiatry*. 2018;63(4):216-222. doi:10.1177/0706743718758968
45. American Psychiatric Association., American Psychiatric Association., DSM-5 Task Force. *Diagnostic and Statistical Manual of Mental Disorders : DSM-5*. American Psychiatric Association; 2013.
46. Friberg O, Barlaug D, Martinussen M, Rosenvinge JH, Hjemdal O. Resilience in relation to personality and intelligence. *Int J Methods Psychiatr Res*. 2005;14(1):29-42. doi:10.1002/mpr.15
47. Armstrong R, Hall BJ, Doyle J, Waters E. 'Scoping the scope' of a cochrane review. *J Public Health*. 2011;33(1):147-150. doi:10.1093/pubmed/fdr015
48. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005;8(1):19-32. doi:10.1080/1364557032000119616
49. Wickremasinghe D, Kuruvilla S, Mays N, Avan BI. Taking knowledge users' knowledge needs into account in health: an evidence synthesis framework. *Health Policy Plan*. 2015;31(4):527-537. doi:10.1093/heapol/czv079

50. Peters MDJ, Marnie C, Tricco AC, et al. Updated methodological guidance for the conduct of scoping reviews. *JBIEvid Synth*. 2020;18(10). doi:<https://doi.org/10.11124/JBIES-20-00167>
51. Peters M, Godfrey C, McInerney P, Baldini Soares C, Khalil H, Parker D. Chapter 11: Scoping Reviews. In: Aromataris E, Munn Z, eds. *Joanna Briggs Institute Reviewer's Manual*. The Joanna Briggs Institute; 2017. <https://reviewersmanual.joannabriggs.org/>
52. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018;169(7):467-473. doi:10.7326/M18-0850
53. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol*. 2016;75:40-46. doi:10.1016/j.jclinepi.2016.01.021
54. *Covidence Systematic Review Software*.; 2019. www.covidence.org
55. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi:10.1136/bmj.n71
56. Bronfenbrenner U. Ecological models of human development. In: *The International Encyclopedia of Education*. Vol 3. 2nd ed. Oxford: Elsevier; 1994.

APPENDIX 1: Search strategy

PubMed

Searches conducted on August 26, 2021

Search	Query	Records retrieved
#1	("Adolescent"[Mesh] OR "Young Adult"[Mesh] OR "Child"[Mesh]) AND "Indigenous Canadians"[Mesh]	1,498
#2	(child*[tiab] OR youth*[tiab] OR adolescent*[tiab] OR "school age"[tiab] OR "middle school"[tiab] OR "junior high"[tiab] OR "high school"[tiab] OR teen*[tiab] OR "young adult"[tiab] OR "emerging adult"[tiab]) AND ("first nations"[tiab] OR "inuit"[tiab] OR metis[tiab] OR aboriginal*[tiab] OR indigenous[tiab] OR "first people"[tiab] OR native[tiab])	13,565
#3	#1 OR #2	14,648
#4	(Arctic[tw]) OR (circumpolar[tw]) OR ("Prince Edward Island"[tw]) OR ("Newfoundland and Labrador"[tw]) OR ("Nova Scotia"[tw]) OR (Canadian*[tw]) OR (Canada[tw]) OR ("P.E.I."[tw]) OR ("New Brunswick"[tw]) OR ("Quebec"[tw]) OR ("Ontario"[tw]) OR ("Manitoba"[tw]) OR ("Saskatchewan"[tw]) OR ("Alberta"[tw]) OR ("British Columbia"[tw]) OR ("B.C."[tw]) OR ("Northwest Territories"[tw]) OR ("N.W.T."[tw]) OR ("Nunavut"[tw]) OR ("Yukon Territory"[tw])	285,973
#5	"Arctic regions"[Mesh] OR "Canada"[Mesh] OR "Nova Scotia"[Mesh] OR "Newfoundland and Labrador"[Mesh] OR "Prince Edward Island"[Mesh] OR "New Brunswick"[Mesh] OR "Quebec"[Mesh] OR "Ontario"[Mesh] OR "Manitoba"[Mesh] OR "Saskatchewan"[Mesh] OR "Alberta"[Mesh] OR "British Columbia"[Mesh] OR "Northwest Territories"[Mesh] OR "Nunavut"[Mesh] OR "Yukon Territory"[Mesh]	174,128

#6	#4 OR #5	285,973
#7	(“Suicide”[Mesh]) OR (“Mental Health”[Mesh]) OR (“Mental Disorders”[Mesh]) OR (“Psychology”[Mesh:NoExp]) OR (“Psychology, Adolescent”[Mesh]) OR (“Resilience, Psychological”[Mesh]) OR (“Community Mental Health Services”[Mesh]) OR (“Culturally Competent Care”[Mesh]) OR (“Health Services”[Mesh]) OR (“Health Services Accessibility”[Mesh]) OR (“Psychological Distress”[Mesh]) OR (“Behavioral Symptoms”[Mesh]) OR (“Substance-Related Disorders”[Mesh])	3,713,983
#8	“mental* ill*”[tiab] OR suicid*[tiab] OR resilen*[tiab] OR adaptability[tiab] OR "protective factors"[tiab] OR "mental health"[tiab] OR coping[tiab] OR cope[tiab] OR "mental wellness"[tiab] OR "mental wellbeing" OR psycho-social[tiab] OR psychosocial[tiab] OR trauma*[tiab] OR “cultural safety” [tiab] OR “cultural competence”[tiab] OR "social support"[tiab] OR “mental toughness”[tiab] OR “community involvement”[tiab] OR “community support*”[tiab] OR “psychosis”[tiab] OR “mental disorder*”[tiab] OR “substance-related disorder*”[tiab] OR “substance abuse” [tiab] OR “substance use*”[tiab] OR “drug use*”[tiab] OR “drug abuse”[tiab] OR “drug dependence”[tiab] OR alcoholism[tiab] OR “alcohol abuse”[tiab] OR “disruptive behavior*”[tiab] OR “disruptive behaviour*”[tiab] OR "Addiction"[tiab] OR "Alcohol Use"[tiab] OR "Cannabis Use*”[tiab] OR “cannabis abuse”[tiab] OR "Drug Dependenc*”[tiab] OR "Inhalant Abuse"[tiab] OR “inhalant use*”[tiab] OR “opioid abuse”[tiab] OR "Opioid Use*”[tiab] OR "Tobacco Use*”[tiab] OR “prescription abuse”[tiab] OR “chronic stress”[tiab] OR “acute stress”[tiab] OR “adjustment disorder*”[tiab] OR “stress disorder*”[tiab] OR depression[tiab] OR “depressive symptoms”[tiab]	1,441,683

#9	#7 OR #8	4,429,175
#10	#3 AND #6 AND #9	1,024

EMBASE

Searches conducted on August 26, 2021

Search	Query	Records Retrieved
#1	('adolescent'/exp OR 'young adult'/exp OR 'child'/exp OR 'middle school student'/exp OR 'high school student'/exp) AND ('Inuit'/exp OR 'First Nation'/exp)	1,252
#2	(child*:ti,ab OR youth*:ti,ab OR adolescent*:ti,ab OR 'school age':ti,ab OR 'middle school':ti,ab OR 'junior high':ti,ab OR 'high school':ti,ab OR teen*:ti,ab OR 'young adult':ti,ab OR 'emerging adult':ti,ab) AND (inuit:ti,ab OR metis:ti,ab OR aboriginal*:ti,ab OR indigenous:ti,ab OR 'first people':ti,ab OR native:ti,ab)	16,938
#3	#1 OR #2	17,642
#4	'Canada'/exp OR 'arctic region'/exp	200,995
#5	Arctic:ti,ab,kw OR circumpolar:ti,ab,kw OR 'Prince Edward Island':ti,ab,kw OR 'Newfoundland and Labrador':ti,ab,kw OR 'Nova Scotia*':ti,ab,kw OR Canadian*:ti,ab,kw OR Canada:ti,ab,kw OR 'P.E.I.':ti,ab,kw OR 'New Brunswick':ti,ab,kw OR Quebec:ti,ab,kw OR Ontario:ti,ab,kw OR Manitoba*:ti,ab,kw OR Saskatchewan:ti,ab,kw OR Alberta*:ti,ab,kw OR 'British Columbia*':ti,ab,kw OR 'B.C.':ti,ab,kw OR 'Northwest Territories':ti,ab,kw OR 'N.W.T.':ti,ab,kw OR Nunavut:ti,ab,kw OR 'Yukon Territory':ti,ab,kw	288,706
#6	#4 OR #5	350,592
#7	'Suicide'/exp OR 'Mental Health'/exp OR 'Mental Disease'/exp OR 'Psychology'/de OR 'Psychological Resilience'/exp OR 'community	8,104,052

	mental health service'/exp OR 'cultural competence'/exp OR 'health service'/exp OR 'health care access'/exp OR 'drug dependence'/exp OR 'child psychiatry'/exp OR 'disruptive behavior'/exp OR 'acute psychosis'/exp	
#8	'mental* ill*':ti,ab OR 'suicide':ti,ab OR 'child psychology':ti,ab OR resilen*:ti,ab OR 'protective factors':ti,ab OR 'mental health':ti,ab OR coping:ti,ab OR cope:ti,ab OR 'mental wellness':ti,ab OR 'mental wellbeing':ti,ab OR 'psycho-social':ti,ab OR psychosocial:ti,ab OR 'cultural safety':ti,ab OR 'cultural competence':ti,ab OR 'social support':ti,ab OR trauma*:ti,ab OR 'cultur* connectedness':ti,ab OR 'mental toughness':ti,ab OR adaptability:ti,ab OR 'community involvement':ti,ab OR 'community support*':ti,ab OR 'psychosis':ti,ab OR 'mental disorder*':ti,ab OR 'substance-related disorder*':ti,ab OR 'substance abuse':ti,ab OR 'substance use*':ti,ab OR 'drug use*':ti,ab OR 'drug abuse':ti,ab OR 'drug dependence':ti,ab OR alcoholism:ti,ab OR 'alcohol abuse':ti,ab OR 'disruptive behavior*':ti,ab OR 'disruptive behaviour*':ti,ab OR 'Addiction':ti,ab OR 'Alcohol Use':ti,ab OR 'Cannabis Use':ti,ab OR 'cannabis abuse':ti,ab OR 'Drug Dependenc*':ti,ab OR 'Inhalant Abuse':ti,ab OR 'inhalant use*':ti,ab OR 'opioid abuse':ti,ab OR 'Opioid Use':ti,ab OR 'Tobacco Use Disorder':ti,ab OR 'prescription abuse':ti,ab OR 'chronic stress':ti,ab OR 'acute stress':ti,ab OR 'adjustment disorder*':ti,ab OR 'stress disorder*':ti,ab OR depression:ti,ab OR 'depressive symptoms':ti,ab	1,841,855
#9	#7 OR #8	8,728,982
#10	#3 AND #6 AND #9	1,165

CINAHL

Searches conducted on August 26, 2021

Search	Query	Records Retrieved

#1	<p>TI ((child* OR youth* OR adolescent* OR “school age” OR “middle school” OR “junior high” OR “high school” OR teen* OR “young adult” OR “emerging adult”) AND (“first nations” OR “inuit” OR “metis” OR “aboriginal*” OR “indigenous” OR “first people” OR native))</p> <p>OR AB ((child* OR youth* OR adolescent* OR “school age” OR “middle school” OR “junior high” OR “high school” OR teen* OR “young adult” OR “emerging adult”) AND (“first nations” OR “inuit” OR “metis” OR “aboriginal*” OR “indigenous” OR “first people” OR native))</p>	5,614
#2	(((MH "Adolescence+") OR (MH "Young Adult") OR (MH "Child"))) AND (MH “Indigenous Peoples+”))	5,431
#3	S1 OR S2	9,555
#4	(MH "Canada+") OR (MH "Arctic Regions")	106,724
#5	TX (Arctic OR circumpolar OR “Newfoundland and Labrador” OR “Nova Scotia*” OR Canadian* OR Canada OR “New Brunswick” OR Quebec* OR Ontario OR Manitoba* OR Saskatchewan OR Alberta* OR “British Columbia*” OR “B.C.” OR “Northwest Territories” OR “N.W.T.” OR Nunavut OR “Yukon Territory”)	509,284
#6	S4 OR S5	509,311
#7	(MH "Suicide+") OR (MH "Mental Health") OR (MH "Mental Disorders+") OR (MH "Adolescent Psychology") OR (MH "Child Psychology") OR (MH "Psychology, Social+") OR (MH "Psychology") OR (MH "Hardiness") OR (MH "Community Mental Health Services") OR (MH "Cultural Competence”) OR (MH "Health Services Accessibility+") OR (MH "Health Services+") OR (MH "Substance Use Disorders+") OR (MH "Child Psychiatry") OR (MH "Behavioral Symptoms")	2,437,255

#8	<p>TI (“mental* ill*” OR suicid* OR resilen* OR “protective factors” OR “mental health” OR coping OR cope OR “mental wellness” OR “mental wellbeing” OR “psycho-social” OR psychosocial OR “cultural safety” OR “cultural competence” OR “social support” OR trauma* OR “cultur* connectedness” OR “mental toughness” OR adaptability OR “community involvement” OR “community support*” OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR "Addiction" OR "Alcohol Use" OR "Cannabis Use*" OR “cannabis abuse” OR "Drug Dependenc*" OR "Inhalant Abuse" OR “inhalant use*” OR “opioid abuse” OR "Opioid Use*" OR "Tobacco Use*” OR “prescription abuse” OR “chronic stress” OR “acute stress” OR “adjustment disorder*” OR “stress disorder*” OR depression OR “depressive symptoms”)</p> <p>OR AB (“mental* ill*” OR suicid* OR resilen* OR “protective factors” OR “mental health” OR coping OR cope OR “mental wellness” OR “mental wellbeing” OR “psycho-social” OR psychosocial OR “cultural safety” OR “cultural competence” OR “social support” OR trauma* OR “cultur* connectedness” OR “mental toughness” OR adaptability OR “community involvement” OR “community support*” OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR "Addiction" OR "Alcohol Use" OR "Cannabis Use*" OR “cannabis abuse” OR "Drug Dependenc*" OR "Inhalant Abuse" OR “inhalant use*” OR “opioid abuse” OR "Opioid Use*" OR "Tobacco Use*” OR “prescription abuse” OR “chronic</p>	592,034
----	--	---------

	stress" OR "acute stress" OR "adjustment disorder*" OR "stress disorder*" OR depression OR "depressive symptoms")	
#9	S6 OR S7	2,752,063
#10	S3 AND S6 AND S9	2,003

PsycInfo

Searches conducted on August 26, 2021

Search	Query	Records Retrieved
#1	(DE "Adolescent Development" OR DE "Adolescent Behavior" OR DE "Emerging Adulthood" OR DE "Early Adolescence") AND (DE "Indigenous Populations" OR DE "Inuit")	123
#2	TI (child* OR youth* OR adolescent* OR "school age" OR "middle school" OR "junior high" OR "high school" OR teen* OR "young adult" OR "emerging adult") AND (inuit OR metis OR aboriginal* OR indigenous OR "first people" OR "first nation*" OR native) OR AB (child* OR youth* OR adolescent* OR "school age" OR "middle school" OR "junior high" OR "high school" OR teen* OR "young adult" OR "emerging adult") AND (inuit OR metis OR aboriginal* OR indigenous OR "first people" OR "first nation*" OR native)	11,751
#3	S1 OR S2	11,756
#4	TX ("Arctic Regions" OR arctic OR circumpolar OR "Prince Edward Island" OR "P.E.I." OR "Newfoundland and Labrador" OR "Nova Scotia*" OR Canadian* OR Canada OR "New Brunswick" OR Ontario OR Manitoba* OR Saskatchewan OR Alberta* OR "British Columbia*" OR "B.C." OR "Northwest Territories" OR "N.W.T." OR "Nunavut" OR "Yukon Territory")	347,047

#5	((((((((DE "Suicidal Behavior" OR DE "Youth Suicide" OR DE "Mental Disorders" OR DE "Self-Injurious Behavior" OR DE "Suicide Prevention") OR (DE "Mental Health")) OR (DE "Psychology")) OR (DE "Adolescent Psychology")) OR (DE "Resilience (Psychological)")) OR (DE "Emotional Adjustment")) OR (DE "Adaptability (Personality)")) OR (DE "Community Mental Health Services")) OR (DE "Cultural Competence")) OR (DE "Health Care Access")) OR (DE "Mental Health Services" OR DE "Substance Use Disorder" OR DE "Addiction" OR DE "Alcohol Use Disorder" OR DE "Cannabis Use Disorder" OR DE "Drug Abuse" OR DE "Drug Dependency" OR DE "Inhalant Abuse" OR DE "Opioid Use Disorder" OR DE "Tobacco Use Disorder" OR DE "Child Psychiatry" OR DE "Acute Psychosis" OR DE "Acute Schizophrenia")	399,746
#6	<p>TI (“mental* ill*” OR suicid* OR resilen* OR “protective factors” OR “mental health” OR coping OR cope OR “mental wellness” OR “mental wellbeing” OR “psycho-social” or psychosocial OR “cultural safety” OR “cultural competence” OR “social support” OR trauma* OR “cultur* connectedness” OR “mental toughness” OR adaptability OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR "Addiction" OR "Alcohol Use" OR "Cannabis Use*" OR “cannabis abuse” OR "Drug Dependenc*" OR "Inhalant Abuse" OR “inhalant use*” OR “opiod abuse” OR "Opioid Use*" OR "Tobacco Use*” OR “prescription abuse” OR “chronic stress” OR “acute stress” OR “adjustment disorder*” OR “stress disorder*” OR depression OR “depressive symptoms”)</p> <p>OR AB (“mental* ill*” OR suicid* OR resilen* OR “protective factors” OR “mental health” OR coping OR cope OR “mental</p>	943,972

	wellness” OR “mental wellbeing” OR “psycho-social” or psychosocial OR “cultural safety” OR “cultural competence” OR “social support” OR trauma* OR “cultur* connectedness” OR “mental toughness” OR adaptability OR “community involvement” OR “community support*” OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR "Addiction" OR "Alcohol Use" OR "Cannabis Use*" OR “cannabis abuse” OR "Drug Dependenc*" OR "Inhalant Abuse" OR “inhalant use*” OR “opioid abuse” OR "Opioid Use*" OR "Tobacco Use*” OR “prescription abuse” OR “chronic stress” OR “acute stress” OR “adjustment disorder*” OR “stress disorder*” OR depression OR “depressive symptoms”)	
#7	S5 OR S6	1,076,814
#8	S3 AND S4 AND S7	619

ERIC

Searches conducted on August 26, 2021

Search	Query	Results Retrieved
#1	(MAINSUBJECT.EXACT("Late Adolescents") OR MAINSUBJECT.EXACT("Early Adolescents") OR MAINSUBJECT.EXACT("Preadolescents") OR MAINSUBJECT.EXACT.EXPLODE("Youth") OR MAINSUBJECT.EXACT.EXPLODE("Secondary School Students") OR MAINSUBJECT.EXACT("Adolescents") OR MAINSUBJECT.EXACT("Young Adults")) AND (MAINSUBJECT.EXACT("Canada Natives") OR MAINSUBJECT.EXACT("Eskimos") OR MAINSUBJECT.EXACT("Indigenous Populations"))	615

#2	<p>ti ((child* OR youth* OR adolescent* OR “school age” OR “middle school” OR “junior high” OR “high school” OR teen* OR “young adult” OR “emerging adult”) AND (“first nations” OR “inuit” OR metis OR aboriginal* OR indigenous OR “first people” OR native))</p> <p>OR ab ((child* OR youth* OR adolescent* OR “school age” OR “middle school” OR “junior high” OR “high school” OR teen* OR “young adult” OR “emerging adult”) AND (“first nations” OR “inuit” OR metis OR aboriginal* OR indigenous OR “first people” OR native))</p>	8,635
#3	1 OR 2	8,913
#4	<p>Canada OR Canadian* OR “Arctic Regions” OR arctic OR circumpolar OR “Prince Edward Island” OR “P.E.I.” OR “Newfoundland and Labrador” OR “Nova Scotia*” OR “New Brunswick” OR Quebec* OR Ontario OR Manitoba* OR Saskatchewan OR Alberta* OR “British Columbia*” OR “B.C.” OR “Northwest Territories” OR “N.W.T.” OR Nunavut OR “Yukon Territory”</p>	65,760
#5	<p>MAINSUBJECT.EXACT("Coping") OR</p> <p>MAINSUBJECT.EXACT.EXPLODE("Mental Disorders") OR</p> <p>MAINSUBJECT.EXACT("Resilience (Psychology)") OR</p> <p>MAINSUBJECT.EXACT("Access to Health Care") OR</p> <p>MAINSUBJECT.EXACT("Mental Health") OR</p> <p>MAINSUBJECT.EXACT("Suicide") OR</p> <p>MAINSUBJECT.EXACT("Child Psychology") OR</p> <p>MAINSUBJECT.EXACT("Community Health Services") OR</p> <p>MAINSUBJECT.EXACT.EXPLODE("Substance Abuse") OR</p> <p>MAINSUBJECT.EXACT.EXPLODE("Child Psychology") OR</p> <p>MAINSUBJECT.EXACT.EXPLODE("Psychosis")</p>	78,985
#6	<p>ti (“mental* ill*” OR suicid* OR coping OR cope OR resilen* OR “protective factors” OR “mental health” OR “mental wellness” OR</p>	85,642

	<p> “mental wellbeing” OR “psycho-social” OR psychosocial OR “cultural safety” OR “cultural competence” OR “social support” OR “community mental health services” OR “culturally competent care” OR “health services” OR “health services accessibility” OR trauma* OR “cultur* connectedness” OR “mental toughness” OR adaptability OR “community involvement” OR “community support*” OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR "Addiction" OR "Alcohol Use" OR "Cannabis Use*" OR “cannabis abuse” OR "Drug Dependenc*" OR "Inhalant Abuse" OR “inhalant use*” OR “opioid abuse” OR "Opioid Use*" OR "Tobacco Use*” OR “prescription abuse” OR “chronic stress” OR “acute stress” OR “adjustment disorder*” OR “stress disorder*” OR depression OR “depressive symptoms”) </p> <p> OR ab (“mental* ill*” OR suicid* OR coping OR cope OR resilen* OR “protective factors” OR “mental health” OR “mental wellness” OR “mental wellbeing” OR “psycho-social” OR psychosocial OR “cultural safety” OR “cultural competence” OR “social support” OR “community mental health services” OR “culturally competent care” OR “health services” OR “health services accessibility” OR trauma* OR “cultur* connectedness” OR “mental toughness” OR adaptability OR “community involvement” OR “community support*” OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR "Addiction" OR "Alcohol Use" OR "Cannabis Use*" OR “cannabis abuse” OR "Drug Dependenc*" OR "Inhalant Abuse" OR “inhalant use*” OR </p>	
--	---	--

	“opioid abuse” OR "Opioid Use*" OR "Tobacco Use*" OR “prescription abuse” OR “chronic stress” OR “acute stress” OR “adjustment disorder*” OR “stress disorder*” OR depression OR “depressive symptoms”)	
#7	5 OR 6	126,622
#8	3 AND 4 AND 7	153

Scopus

Searches conducted on August 26, 2021

Search	Query	Results Retrieved
#1	TITLE-ABS-KEY ((Adolescent* OR “young adult” OR child OR children OR youth* OR “school age” OR “middle school” OR “junior high” OR “high school” OR teen* OR “young adult” OR “emerging adult”) AND (“Indigenous Canadian*” OR inuit OR metis OR “first nations” OR aboriginal* OR indigenous OR “first people” OR native))	42,666
#2	TITLE-ABS-KEY (Canada OR Canadian* OR “Arctic regions” OR Arctic OR {Prince Edward Island} OR “Newfoundland and Labrador” OR "Nova Scotia*" OR {P.E.I.} OR {New Brunswick} OR Quebec* OR Ontario OR Manitoba* OR Saskatchewan OR Alberta* OR “British Columbia*” OR {B.C.} OR “Northwest Territories” OR {N.W.T.} OR Nunavut OR “Yukon Territory”)	725,032
#3	TITLE-ABS-KEY (“mental* ill*” OR Suicid* OR {mental health} OR “mental disorder*” OR {adolescent psychology} OR {psychological resilience} OR resilen* OR “protective factors” OR coping OR cope OR {mental wellness} OR “mental wellbeing” OR {psycho-social} OR psychosocial OR “cultural safety” OR “cultural competence” OR “social support*” OR “community mental health services” OR “culturally competent care” OR “health services” OR “health services accessibility” OR trauma* OR “cultur* connectedness” OR {mental	3,570,197

	toughness} OR adaptability OR “community involvement” OR “community support*” OR {Behavioral Symptoms} OR {Adolescent Psychiatry} OR {Substance-Related Disorders} OR “psychosis” OR “mental disorder*” OR “substance-related disorder*” OR “substance abuse” OR “substance use*” OR “drug use*” OR “drug abuse” OR “drug dependence” OR alcoholism OR “alcohol abuse” OR “disruptive behavior*” OR “disruptive behaviour*” OR “Addiction” OR “Alcohol Use” OR “Cannabis Use*” OR “cannabis abuse” OR “Drug Dependenc*” OR “Inhalant Abuse” OR “inhalant use*” OR “opioid abuse” OR “Opioid Use*” OR “Tobacco Use*” OR “prescription abuse” OR “chronic stress” OR “acute stress” OR “adjustment disorder*” OR “stress disorder*” OR depression OR “depressive symptoms”)	
#4	#1 AND #2 AND #3	1,393

APPENDIX 2: Data Extraction Instrument

Study title	
Authors	
Year of publication	
Aim of study	
Type of study	
Literature type	
Country of publication	
Study population (participants)	
Geographical setting	
Type of service (healthcare, education, community)	
Description of community involvement	
Involved community, knowledge users, or stakeholders	
Use of a theory or framework	
Methodology/methods	
Outcome measurement tool	
Type of mental health studied	
Definition of resiliency/protective factors/coping skills used	
Description of resilience or protective factors	
Reported outcomes	
Study main findings and conclusion	
Future implications	
Other notes	
Study title	
Authors	
Year of publication	
Aim of study	
Type of study	
Literature type	
Country of publication	
Study population (participants)	
Geographical setting	
Type of service (healthcare, education, community)	
Level of community involvement	

Use of a theory or framework	
Methodology/methods	
Measurement tool	
Type of mental health studied	
Definition of resiliency/protective factors/coping skills	
Resilience or protective factors identified	
Author classification factor	
Level of resiliency factors (individual, family, community, system)	
Reported outcomes	
Study main findings and conclusion	
Future implications	
Other notes	

2
3
4