Matrix metalloproteinase 14 (MMP14) is differentially expressed in breast cancer metastases.

Shahan Mamoor, MS¹

¹Thomas Jefferson School of Law

San Diego, CA 92101

mamoorsk@tjssl.edu

Breast cancer is a leading cause of death for women (1, 2). We mined public datasets to understand the most significant transcriptional and epigenomic changes between primary tumors of the breast and the metastases that they generate (3-6). In metastases to both the brain (3) and the bones (4), the matrix metalloproteinase 14 (MMP14) was among the genes whose expression changes most significantly between the transcriptomes of metastatic tissue to that of primary breast tumor. In metastases to the lymph nodes, a site at the MMP14 genomic locus contained significantly less methyl marks than the same site in primary tumors. The MMP14 gene was hypomethylated during spread of the breast cancer to the lymph node, MMP14 among the genes whose expression changed most significantly across the entire brain and bone transcriptome, and MMP14 messenger RNA was expressed at significantly lower levels in metastases across both tissues. Interestingly, primary tumors with metastatic potential expressed higher levels of MMP14 than primary tumors without metastatic potential. MMP14 is a target of interest in metastatic breast cancer.

Keywords: breast cancer, metastasis, MMP14, matrix metalloproteinase 14, systems oncology, tumor transcriptome analysis, targeted therapeutics in breast cancer

1 In 2018, over a quarter of a million women were diagnosed with breast cancer in the
2 United States (1). While the 5-year survival rate for women with stage I disease, cancer that
3 has not spread beyond the breast, is 98% (1), the 5-year survival rate for women diagnosed
4 with metastatic cancer that has spread to organs outside of the breast such as the lungs, the
5 liver, the bone and the brain is 36% (2). The rational design or discovery of novel therapeutics
6 for women with metastatic breast cancer requires identification of specific target genes and
7 pathways that can be acted upon. In order to determine in a systematic and unbiased manner
8 which genes are expressed most differently between primary tumors of the breast and
9 metastasis to foreign sites, we compared total transcription (3, 4, 6) and total DNA methylation
10 data (5) using four independently published datasets generated using patient tumor and
11 metastasis tissues. We identified matrix metalloproteinase MMP14 as the one of the most
12 differentially expressed genes when comparing primary breast tumors and metastasis to the
13 bone and brain at the transcriptome-level (3, 4). Analysis of a separate dataset revealed that in
14 lymph node metastases, the MMP14 genomic locus was one of the most differentially
15 methylated sites in the entire genome (5). MMP14 was expressed at significantly lower levels
16 in metastases to the brain and the bone. However, when comparing the primary tumors of
17 patients whose disease did not metastasize to the primary tumors of patients whose disease did
18 metastasize, MMP14 was expressed at higher levels in the primary breast tumors with
19 metastatic potential (6). These data support a notion that MMP14 is a target of interest in
20 understanding the fundamental biology of breast cancers and the deadly process of metastasis.
21
22
23

24 **Methods**

25 We used GSE125989 (3), GSE124648 (4), GSE59000 (5), and GSE102484 (6) for this
26 analysis, in conjunction with GEO2R. GSE125989 was generated using Affymetrix Human
27 Genome U133A 2.0 Array, with n=16 breast cancer primary tumors and n=16 brain metastases
28 (paired samples). GSE124648 was generated using Affymetrix Human Genome U133A Array
with n=124 breast cancer primary tumor and n=11 bone metastases. GSE59000 was generated

using Illumina HumanMethylation450 BeadChip with n=36 primary tumors of the breast and n=36 lymph node (LN) metastases. GSE102484 was generated using Affymetrix Human Genome U133 Plus 2.0 Array with n=582 primary breast tumors that did not metastasize and n=101 primary breast tumors that did metastasize. The “Benjamini and Hochberg” method of p-value adjustment was used for transcriptome rank, but raw p-values were used to determine whether a gene was differentially expressed with respect to the transcriptome. For log-transformation of data on GEO2R, the “auto-detect” option was utilized, and the NCBI generated category of platform annotation was utilized. We analyzed, independent of whether a gene was differentially expressed transcriptome-wide, whether the difference in mRNA expression levels was different between the two groups in which the gene was differentially expressed (e.g., tumor and metastasis). We performed this statistical analysis using PRISM (8.1.2)(227) using a two-tailed unpaired t-test with Welch’s correction and the results of each test is shown on the corresponding graph as a p-value. P-values less than 0.05 were considered statistically significant.

Results

We performed whole transcriptome comparative analysis of primary and metastatic breast tumors of the brain (3) and the bone (4) using two independent published datasets. This was complemented by analysis of global DNA methylation in lymph node metastases using a public dataset (5), and whole transcriptome analysis of primary tumors from patients whose tumors did or did not metastasize using a fourth dataset (6).

MMP14 is differentially expressed in metastasis to the bone and to the brain when compared to primary tumors of the breast

We identified MMP14 as one of the most differentially expressed genes in metastasis to both the brain and bones in patients with breast cancer when compared to primary breast tumors. In the brain, MMP14 was the 45th most differentially expressed gene out of 22277 total

transcripts. In the bone, MMP14 was the 172nd most differentially expressed gene out of 22284 total transcripts. This was statistically significant for both metastases to the brain (Table 1; $p=2.09E-05$) as well as the bones (Table 1; $p=6.91E-14$).

MMP14 is expressed at lower levels in the bone and the brain when compared to primary tumors of the breast

We then obtained exact mRNA expression values for MMP14 in each primary tumor of the breast, in metastases to the brain and in metastases to the bone. When comparing 16 breast tumors to paired brain metastases from the same patient, MMP14 was expressed at significantly lower levels in brain metastasis tissue (Figure 1; $p<0.0001$). When comparing primary tumors of the breast to bone metastases, MMP14, again, was expressed at significantly lower levels in metastatic tissue (Figure 2; $p<0.0001$).

In metastasis to the lymph nodes, a site at the MMP14 genomic locus is among the most differentially methylated genome-wide

Next, we probed a dataset that analyzed global DNA methylation in 36 primary tumors of the breast and 36 lymph node metastases. We identified MMP14 as one of the most differentially methylated genes across the genome. A site at the genomic locus of MMP14 was the 88th most differentially methylated locus out of 22277 regions probed. This was statistically significant (Table 1; $p=9.33E-05$). When comparing the level of DNA methylation present at this differentially methylated site between lymph node metastases and primary tumors of the breast, the MMP14 gene possessed significantly less methyl marks in lymph node metastases than the same site in primary breast tumors (Figure 3, $p=0.0001$). Thus, in metastases to the lymph nodes from patients with breast cancer, MMP14 is among the genes whose DNA methylation changes the most across the entire genome, and this differentially methylated site is significantly hypomethylated when compared to primary breast tumors.

MMP14 is expressed at higher levels in the primary breast tumors with metastatic potential when compared to primary breast tumors that did not metastasize

Finally, we compared transcriptome data from 582 primary tumors of the breast that did not metastasize to transcriptome data from 101 primary tumors of the breast that did metastasize. MMP14 was one of the genes whose expression changed most significantly between primary tumors with and without metastatic potential, ranking 165 out of 54675 total transcripts. This was statistically significant (Table 1; $p=2.13E-06$). Primary tumors from patients that subsequently developed metastatic disease expressed significantly higher levels of MMP14 than primary tumors from patients did not develop metastases (Figure 4; $p<0.0001$).

Discussion

The 5-year survival rate for patients with metastatic breast cancer is 36%. Understanding the most significant changes in the transcriptomes of metastases when compared to primary tumors in breast cancer can facilitate discovery of actionable targets for development of novel therapeutics. We analyzed transcriptome and epigenome data from four separate published (3, 4, 6) or public (5) datasets and identified MMP14 as among the most differentially expressed genes in metastatic breast cancer. The MMP14 genomic locus was significantly less methylated in metastases to the lymph node than in primary breast tumors. However, expression of MMP14 in breast tumors from patients who developed metastases was significantly higher than in breast tumors from patients who did not develop metastases.

MMP14 over-expression in the tumors of patients with breast cancer as compared to primary tissues of the breast has previously been reported, and invasive behavior of breast cancer cells has been attributed to MMP14 expression. One study found that there was a relationship between higher MMP14 expression in primary breast tumors and a poorer prognosis but that this relationship did not reach the level of significance (7). Another study assessed three separate breast cancer patient cohorts and found that higher MMP14 mRNA expression in primary tumors did significantly correlate with decreased overall survival (8).

1 Matrix metalloproteinases, including MMP14, have a conserved homopexin-like or PEX domain
2 that possesses a beta-propeller domain with four blades (8). The PEX domain of MMP14 is
3 important for association with CD44 and subsequent activation of proMMP2 by MMP14 (8).
4 Synthetic peptides that contain sequences from outer portions of the four blades of MMP14
5 inhibit migratory behavior of breast cancer cells *in vitro* and interfere with metastasis in the
6 MDA-MB-435 xenograft mouse model (8). A humanized antibody that binds MMP14 has limiting
7 effects on primary tumor growth as well as on metastasis, in the MDA-MB-231 orthotopic
8 xenograft mouse models and in the 4T1 syngeneic mouse model of triple negative breast
9 cancer (9). Another study using a different antibody targeting MMP14, DX-2400, significantly
10 inhibited primary tumor growth in two separate xenograft models (4T1 and E0771) (10). A study
11 using the mammary tumor virus-polyoma middle T mouse breast cancer model isolated tumor-
12 initiating cells (TIC) and found that Mmp14 was significantly up-regulated in TIC (11). Together,
13 these published studies show that MMP14 inhibition is effective in multiple xenograft mouse
14 models of breast cancer in either reducing tumor growth or metastasis to foreign sites. Our
15 findings show that MMP14 is among the genes whose expression changes most significantly
16 out of all of the genes expressed cell when comparing metastases and primary breast tumors.
17 MMP14 is expressed at significantly lower levels in metastases than in the primary tumor from
18 which it originated (3). This could suggest that while MMP14 up-regulation may be critical for
19 initiation of the metastatic process (Figure 4) (8), down-regulation of MMP14 in metastatic tissue
20 might be similarly critical for establishing and maintaining the metastasis.
21
22

23 Biologically speaking, MMP14 is one of the most significantly different genes when
24 comparing breast cancers to metastases they generate at the level of gene expression and at
25 the level of DNA methylation.
26
27
28

References

1. DeSantis, C.E., Ma, J., Gaudet, M.M., Newman, L.A., Miller, K.D., Goding Sauer, A., Jemal, A. and Siegel, R.L., 2019. Breast cancer statistics, 2019. *CA: a cancer journal for clinicians*, 69(6), pp.438-451.
2. Mariotto, A.B., Etzioni, R., Hurlbert, M., Penberthy, L. and Mayer, M., 2017. Estimation of the number of women living with metastatic breast cancer in the United States. *Cancer Epidemiology and Prevention Biomarkers*, 26(6), pp.809-815.
3. Iwamoto T, Niikura N, Ogiya R, Yasojima H et al. Distinct gene expression profiles between primary breast cancers and brain metastases from pair-matched samples. *Sci Rep* 2019 Sep 16;9(1):13343
4. Sinn BV, Fu C, Lau R, Litton J et al. SET_{ER/PR}: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ Breast Cancer* 2019;5:16.
5. GSE59000. "Methylome remodeling in regional breast cancer metastasis". Marsha Reyngold, Timothy Chan Lab. Memorial Sloan-Kettering Cancer Center, Radiation Oncology
6. Cheng, S.H.C., Huang, T.T., Cheng, Y.H., Tan, T.B.K., Horng, C.F., Wang, Y.A., Brian, N.S., Shih, L.S. and Yu, B.L., 2017. Validation of the 18-gene classifier as a prognostic biomarker of distant metastasis in breast cancer. *PloS one*, 12(9).
7. Têtu, B., Brisson, J., Wang, C.S., Lapointe, H., Beaudry, G., Blanchette, C. and Trudel, D., 2006. The influence of MMP-14, TIMP-2 and MMP-2 expression on breast cancer prognosis. *Breast Cancer Research*, 8(3), p.R28
8. Zarrabi, K., Dufour, A., Li, J., Kuscu, C., Pulkoski-Gross, A., Zhi, J., Hu, Y., Sampson, N.S., Zucker, S. and Cao, J., 2011. Inhibition of matrix metalloproteinase 14 (MMP-14)-mediated cancer cell migration. *Journal of Biological Chemistry*, 286(38), pp.33167-33177.
9. Ling, B., Watt, K., Banerjee, S., Newsted, D., Truesdell, P., Adams, J., Sidhu, S.S. and Craig, A.W., 2017. A novel immunotherapy targeting MMP-14 limits hypoxia, immune suppression and metastasis in triple-negative breast cancer models. *Oncotarget*, 8(35), p. 58372.
10. Ager, E.I., Kozin, S.V., Kirkpatrick, N.D., Seano, G., Kodack, D.P., Askoxylakis, V., Huang, Y., Goel, S., Snuderl, M., Muzikansky, A. and Finkelstein, D.M., 2015. Blockade of MMP14

1 activity in murine breast carcinomas: implications for macrophages, vessels, and
2 radiotherapy. *JNCI: Journal of the National Cancer Institute*, 107(4).

- 3 11. Hillebrand, L.E., Wickberg, S.M., Gomez-Auli, A., Follo, M., Maurer, J., Busch, H., Boerries,
4 M. and Reinheckel, T., 2019. MMP14 empowers tumor-initiating breast cancer cells under
5 hypoxic nutrient-depleted conditions. *The FASEB Journal*, 33(3), pp.4124-4140.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Rank	ID	p-value	t	B	Gene	Gene Name
GSE125989						
45	202827_s_at	2.09E-05	-4.8588886	2.6937	MMP14	matrix metalloproteinase 14
GSE124649						
172	202827_s_at	6.91E-14	-8.3498205	21.10693	MMP14	matrix metalloproteinase 14
GSE59000						
88	160020_at	9.33E-05	-4.137874	1.27295	MMP14	matrix metalloproteinase 14
GSE102484						
165	202827_s_at	2.13E-06	4.78163766	4.599928	MMP14	matrix metalloproteinase 14

Table 1: MMP14 is differentially expressed in the bone and the brain, and differentially methylated in the lymph nodes in metastatic breast cancer.

The rank of global differential gene expression for the MMP14 mRNA message or the MMP14 gene (GSE59000), the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expressed between the two groups compared, the gene symbol and gene name are shown in this chart (obtained from GEO2R).

Figure 1: MMP14 is expressed at lower levels in metastasis to the brain when compared to primary tumors of the breast

The mRNA expression level of the MMP14 gene in primary breast tumors (blue) and in metastases to the brain (red) is graphically represented by dots denoting expression in individual tumors with the horizontal black bar denoting the mean mRNA expression level of MMP14 (left), and as a bar graph with the error bars denoting the standard error of the mean (right). The results of a statistical test evaluating the difference in expression between primary tumor and brain metastasis, a p-value, is listed on the graph.

Figure 2: MMP14 is expressed at lower levels in metastases to the bone when compared to primary tumors of the breast.

The mRNA expression level of the MMP14 gene in primary breast tumors (blue) and in metastases to the bones (red) is graphically represented by dots denoting expression in individual tumors with the horizontal black bar denoting the mean mRNA expression level of MMP14 (left), and as a bar graph with the error bars denoting the standard error of the mean (right). The results of a statistical test evaluating the difference in expression between primary tumor and bone metastasis groups, a p-value, is listed on the graph.

Figure 3: A differentially methylated site at the MMP14 genomic locus manifests a significant loss of methyl marks in lymph node metastases.

The DNA methylation level of a site at the MMP14 gene in primary breast tumors (blue) versus lymph node metastases (red) is graphically represented by dots denoting DNA methylation in individual tumors with the horizontal black bar denoting the mean DNA methylation level of MMP14 (left), and as a bar graph with the error bars denoting the standard error of the mean (right). The results of a statistical test evaluating the difference in expression between primary tumor and lymph node metastasis groups, a p-value, is listed on the graph.

Figure 4: MMP14 is expressed at higher levels in the primary breast tumors with metastatic potential when compared to primary breast tumors that did not metastasize

The mRNA expression level of the MMP14 gene in primary breast tumors without metastatic potential (blue) and in primary breast tumors with metastatic potential (red) is graphically represented by a violin plot with the dotted black line representing the median MMP14 mRNA expression level, and the dotted colored lines representing the 25th and 75th quartiles of expression (left), and with dots denoting expression in individual tumors with the horizontal black bar denoting the mean mRNA expression level of MMP14 (right). The results of a statistical test evaluating the difference in expression between non-metastatic and metastatic groups, a p-value, is listed on the graph.