POT1 is a differentially expressed gene in brain metastatic human breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with brain metastasis in humans with metastatic breast cancer. We found that protection of telomeres 1, encoded by POT1, was among the genes whose expression was most quantitatively different in the brain metastases of patients with metastatic breast cancer. An antisense non-coding RNA transcribed at the POT1 locus was also found to be differentially expressed in brain metastatic tissues of patients with metastatic breast cancer. POT1 mRNA was present at decreased quantities in brain metastatic tissues as compared to primary tumors of the breast. Importantly, expression of POT1 in primary tumors was significantly correlated with patient overall survival in patients with breast cancer. Modulation of POT1 expression may be relevant to the biology by which tumor cells metastasize from the breast to the brain.

Keywords: breast cancer, metastasis, brain metastases, central nervous system metastases, protection of telomeres 1, POT1, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with brain metastatic breast cancer^{4,5}. We discovered differential and decreased expression of the gene encoding protection of telomeres 1, POT1, in brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE52604⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=11$ primary breast tumors and $n=4$ brain metastases from patients with breast cancer; analysis was performed using platform GPL885. GSE52604 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer; analysis was performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether POT1 gene expression was significantly different between primary tumors of the breast and brain metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of POT1 mRNA expression levels in primary tumors of the breast with overall survival in $n=1402$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic and primary tumor tissues of patients with metastatic breast, as well as normal tissues of the breast using published microarray data^{4,5} to describe the transcriptional landscape of brain metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

POT1 is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified protection of telomeres 1, encoded by POT1, as a differentially expressed gene in the brain metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, POT1 ranked 54 out of 17418 total transcripts (Chart 1), equating to 99.7% differential expression. Differential expression of POT1 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=1.01e-04$).

A POT1 non-coding RNA is differentially expressed in the brain metastases of patients with breast cancer.

Differential gene expression analyses that utilize primary and metastatic tumor tissues to discover genes associated with metastasis can be challenging due to the relatively heterogeneous cellular composition and constitution of the tumor as compared to normal, untransformed (benign) tissues. Thus, to attempt to validate differential transcriptome-wide differential expression of DACT1 in brain metastases in human metastatic breast cancer, we queried a second microarray dataset⁵, here comparing normal breast tissues rather than primary tumors of the breast to brain metastases. Here we identified an antisense non-coding POT1 RNA, POT-AS1, as differentially expressed in the brain metastatic tissues of patients with breast cancer (Chart 2). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to normal breast tissues, POT1 ranked 16619 out of 41093 total transcripts (Chart 2), equating to 59.6% differential expression. Differential expression of POT1 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=4.26e-02$). Thus, differential expression of POT1, transcriptome-wide, in metastatic tissues of women with metastatic breast cancer was conserved across two independent microarray datasets, in the brain and in the lymph nodes.

POT1 is expressed at lower levels in the brain metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for POT1, in primary tumors of the breast and in brain metastasis of patients with brain metastatic breast cancer to determine direction and statistical significance of change in POT1 expression in brain metastatic tissues. We observed significantly decreased expression of POT1 in the brain metastases of patients with breast cancer as compared to primary tumors of the breast: POT1 was expressed at -0.42 ± 0.31 arbitrary units (AU) in primary tumors of the breast, while it was expressed at -1.47 ± 0.39 AU in brain metastatic tissues (Figure 1). The difference in POT1 mRNA levels between primary tumors of the breast and brain metastatic tissues was statistically significant (Figure 1; $p=0.000121$).

POT1 expression is correlated with survival outcomes in human breast cancer.

We performed Kaplan-Meier survival analysis⁷ in 1402 breast cancer patients in total, to evaluate whether POT1 tumor expression was correlated with survival outcomes in breast cancer. We observed a correlation between primary tumor expression of POT1 and overall survival (OS) in patients with breast cancer, in the upper survival quartile (Figure 2). Patients whose primary tumors expressed low levels of POT1 possessed median OS of 85.2 months, while patients whose tumors expressed high levels of POT1 possessed median OS of 143 months. This difference in OS based on POT1 tumor expression in patients with breast cancer was statistically significant (Figure 2, Chart 3; logrank p -value 0.018; hazard ratio: 0.72 (0.55-0.95)).

Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified protection of telomeres 1, encoded by POT1, as among the genes whose expression was most different, transcriptome-wide, in the brain and lymph node metastases of patients with breast cancer, when compared to primary tumors of the breast and to normal breast tissues, respectively; we observed significantly decreased expression of POT1 in brain metastases as compared to primary tumors of the breast. Further, we found a correlation between POT1 expression and patient survival outcomes, as overall survival was greater in patients whose primary tumors expressed higher levels of POT1 as compared to patients whose primary tumors expressed lower levels of POT1.

Discussion

We provided evidence here that POT1 is among the genes whose expression is most different in the brain metastases of patients with metastatic breast cancer, that POT1 mRNA is present at significantly decreased quantity in brain metastatic tissues as compared to primary tumors of the breast, that a POT1 antisense non-coding RNA is also differentially expressed in brain metastases of humans with metastatic breast cancer, and that primary tumor POT1 expression is significantly correlated with patient survival outcomes in human breast cancer. Evaluation of the effects of genetic depletion of POT1 in mouse models of metastatic breast cancer on metastasis to the central nervous system is merited. Modulation of POT1 expression may be relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier and colonize the brain.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Salhia, B., Kiefer, J., Ross, J.T., Metapally, R., Martinez, R.A., Johnson, K.N., DiPerna, D.M., Paquette, K.M., Jung, S., Nasser, S. and Wallstrom, G., 2014. Integrated genomic and epigenomic analysis of breast cancer brain metastasis. *Plos one*, 9(1), p.e85448.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 54
Probe ID: 4415
p-value: 1.01e-04
t: 5.15
B: 1.513241
Gene: POT1
Gene name: protection of telomeres 1

Chart 1: POT1 is differentially expressed in brain metastatic breast cancer when comparing brain metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 16619
probe ID: A_24_P24867
p-value: 4.26e-02
t: 2.0844878
B: -4.634879
Gene: POT1-AS1
Gene name: protection of telomeres 1, antisense RNA 1

Chart 2: An antisense non-coding RNA produced at the POT1 locus is differentially expressed in the lymph node metastases of patients with breast cancer when comparing brain metastases to normal breast tissues.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: POT1 is expressed at significantly lower levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of POT1 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in POT1 expression between primary tumors and brain metastases is $p=0.000121$.

Figure 2: Significant correlation between POT1 primary tumor expression and overall survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival for $n=1402$ total breast cancer patients stratified into two groups, based on low or high expression of POT1 in patient primary tumor, in the upper survival quartile. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on POT1 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

1 Low POT1 expression: 85.2 months

2 High POT1 expression: 143 months

3
4 **Chart 3: Median overall survival is inferior in patients with low primary tumor expression of POT1.**

5 The median OS (overall survival) of patients with low primary tumor expression of POT1 and high
6 primary tumor expression of POT1 is listed in this chart, for $n=1402$ breast cancer patients, in the upper
7 survival quartile.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28