A Review on Therapeutic strategies of Relapsed and Refractory Multiple Myeloma

Dattatreya Mukherjee², MBBS ^{1,2,3}

1. MBBS Student and Research Assistant, Jinan University, Guangzhou, P.R China

2. Researcher at iCART, All India Institute of Medical Science, New Delhi

3. Researcher in the clinical trial and research Division, Larkin Community Hospital, Miami, USA

Article Type: Literature Review

Total Word: 7522 with Reference

References: 87

Corresponding Author: Dattatreya Mukherjee, dattatreyamukherjee4u@outlook.com

CC-BY Attribution-Non-Commercial-No Derivatives 4.0 International

DOI: 10.31219/osf.io/bfj4z

Abstract:

Improvement in treatment options in multiple myeloma (MM) makes the survival rate into double during recent years but most of the patient eventually relapses. The treatment of relapse and refracting MM is still now very complex. At the time of relapse, the use of alternative drugs is the current practice. Using Protease inhibitors like Bortezomib and immune-modulators like thalidomide gives a satisfactory treatment prognosis in MM. Now many new options are available like second and third generation protease inhibitor like carfilzomib and ixazomib, the immune-modulator like lenalidomide, pomalidomide, the monoclonal antibody like daratumumab, elotuzumab, Histone Deacetylase inhibitor, Kinesin Spindle inhibitor and NF- κ B, MAPK, AKT inhibitors. Daratumumab, targeting CD38, has multiple mechanism of action like modulation immunosuppressive bone marrow micro-environment. Daratumumab has a good progression free survival when it is given with lenalidomide/ dexamethasone or bortezomib/dexamethasone. Autologous stem cell transplantation (ASCT) is important treatment in relapsing and refracting MM. Donor lymphocytic infusion is a new and effective technique in the treatment of relapsing and refracting MM. The advancement in the treatment of relapsed and refractory MM has been discussed in this article. The final goal should be finding a balance between efficacy, toxicity and cost.

Keywords: Relapsed, Refractory, Multiple Myeloma, Treatment advancement

Introduction:

Multiple Myeloma (MM) is characterized by the excess amount of monoclonal plasma cell in the bone marrow (BM) associated with monoclonal protein in blood and urine. About 50 years earlier, the use of melphalan and prednisone formulations demonstrated a 2-3 years expansion of the median survival of MM patients. (1) The high dose of melphalan in 1980 followed by bone marrow (BM) transplant and peripheral blood stem cell rescue in 1990 further improved the median survival to 3-4 years. (1) After 1998, due to a new paradigm of drug development towards targeting the tumor cell in their microenvironment, the median survival increased to 7-8 years. (2) The improvement in median survival happens through the administration immunomodulatory drugs (IMiDs) thalidomide, lenalidomide and Proteasome inhibitor (PI) Bortezomib. Application of high dose conventional therapy with autologous stem cell transplant

(HDT-ASCT) to transplant eligible patient is a common treatment method. If the patient is not eligible for transplant then thalidomide or bortezomib associated with melphalan is recommended. (3) These novel agents are the backbone of the treatment of multiple myeloma for both in the diagnosis and relapse setting.

All most all patients with MM will eventually relapse. The choice of a treatment regimen at relapse depends upon many factors including timing of relapse, response to prior therapy, aggressiveness of the relapse and performance status (TRAP). (4) In newly diagnose Case Bortezomib-Lenalidomide- Dexamethasone (VRd), Bortezomib-Cyclophosphamide-Dexamethasone (VCd), Bortezomib-thalidomide-dexamethasone (VTd) are active regimen in the case of relapse disease. (4)

The poor prognosis is shown by the MM patient who had received three prior line therapy, lenalidomide/pomalidomide, Bortezomib/carfilzomib, and has been exposed to an alkylating agent. (5)

Three Daratumumab based regimen shown a good efficacy, Daratumumab with lenalidomide and dexamethasone (DRd), Daratumumab with bortezomib and dexamethasone (DVd), Daratumumab with pomalidomide and dexamethasone. (6)

Other than this regimen Carfilzomib-lenalidomide- Dexamethasone (KRd), ixazomib-lenalidomide-dexamethasone (IRd), elotuzumab-lenalidomide, dexamethasone (ERd), various pomalidomide based regimen like daratumumab-pomalidomide-dexamethasone (DPd) and Carfilzomib, pomalidomide, dexamethasone (KPd) is used and for aggressive treatment anthracycline based regimen is useful. (4)

The new findings about the disease makes a good development in the novel tumor targeting agents which targets a specific pathway of tumor growth and survival. Due to their effectiveness and tolerability, some of this drug research has entered into the phase III and IV clinical studies. (7). The treatment regimen and optimal combination for relapsed and refractory MM is still unknown. It's a challenge to identify the effective regimen and combination for that particular patient. In future study, we should give more concentration on immune based therapy and more precisely tumor microenvironment targeting drugs in case of both frontline and relapse-refractory type of MM

According to NCCN guidelines version 4 2021, for the treatment of RRMM are conventional therapy, clinical trial, autologous hematopoietic cell transplant and allogenic hematopoietic cell transplant. Refractory cases are treated with palliative care.

This paper is a review paper on the last 10 years treatment development in case of relapse and refractory type of MM.

What is Relapse and Refractory MM?

In 2006, the international myeloma working group (IMWG) modified the definition of relapse and refractory MM with the context to the new response criteria. (8,9). Relapse and refractory myeloma is defined as relapse in patients who must have achieved at least minimal response (MR), which either becomes nonresponsive while on salvage therapy or progresses within 60 days of last treatment. (8,9) Primary refractory myeloma is defined as disease that is non-responsive, patients who have never achieved a MR or better with any therapy. (9) Double refractory MM refers to disease refractory to both PI and IMiDs.

Symptoms such as CRAB (increased calcium, renal failure, anemia and bone disease) shows the relapse of the disease. For a patient with non-secretary myeloma, the increase in the level plasma cell in the marrow or bone disease shows the progression where symptomatic relapse is defined as the same way as the patients for secretary myeloma. (10)

Table 1: Criteria of detecting Relapse according to NCCN Guidelines 4.2021

Clinical Relapse	<p>Clinical Relapse need one or more of the following criteria:</p> <ol style="list-style-type: none"> 1. Direct indicators of increasing disease and/or end organ dysfunction (calcium elevation, renal failure, anemia, lytic bone lesion [CRAB]) related to the underlying clonal plasma cell disorder. It is not used in calculations of time to progression or progression free survival but is listed as something that can be reported optionally or for use of clinical practice. 2. Development of new soft tissue plasmacytomas or bone lesions. 3. Definite increase in size of existing plasmacytomas or bone lesions. A definite increase is defined as 50% and more than 1cm increase as measured serially by the SPD of the measurable lesion. 4. Hypercalcemia (>11mg/dl) 5. Decrease in hemoglobin of more than 2g/dL not related to the therapy or other non -myeloma related condition 6. Rise in serum creatinine by 2mg/dL or more 7. Hyper viscosity with serum para-protein
------------------	--

Relapse from Complete Relapse (to be used only if the endpoint is disease free survival)	Any one or more of the following criteria: <ol style="list-style-type: none">1. Reappearance of serum or urine M-protein by immunofixation or electrophoresis.2. Development of more than 5% plasma cells in the bone marrow.3. Appearance of any other sign of progression (new plasmacytoma, lytic bone lesion, hypercalcemia)
Relapse from MRD negative (to be used only if the endpoint is disease free survival)	Any one or more of the following criteria: <ol style="list-style-type: none">1. Loss of MRD negative state2. Reappearance of serum or urine M protein3. Development of more than 5% clonal plasma cell in the bone marrow4. Appearance of any other sign of progression

Clonal Evolution in multiple myeloma:

Clonal Evolution drives tumor progression, chemo resistance and relapse of cancer. According to a study of Jill Corre et al. KRAS, NRAS, DIS3, BRAF and LRP1B are the major mutated gene and during the relapse the mutational burden is same. (11) In the same study double event deletion of 17p/TP3 mutation has been found in relation to clonal evolution in multiple myeloma.

Keats et al. also discovered that about one-third of MM patients have a healthy gene and are at low risk of potentially positive health results. (12) A linear progression was influential in another one-third and another one-third shifts in the genome over time and both are at high risk and potentially bad result in therapy. (12) The evolution of adverse prognostic relapse markers, including MAPK mutation and biallelic inactivation of the tumour suppressor gene, has been shown in additional research. (13,14,1,16)

Keats et al. also discovered that about one-third of MM patients have a healthy gene and are at low risk of potentially positive health results. (17) The main factor in drug resistance and recurrence is this clonal mutation. In CML Philadelphia chromosomal mutation is the key clonal factor in treatment failure. In MM sometime Lenalidomide-dexamethasone combination gives a better result. Few studies have analyzed clonal evolution after treatment (18,19) no-one gives the same treatment in myeloma so the gap is still unfilled. However, in MM evolution, changes in clonal substructure play a very important part and gives the idea of progression of smoldering myeloma to symptomatic myeloma. (20) In the pre-existing state, chemoresistance and mutation occur due to novel mutations and sub clonal mutations.

It's always challenging to treat in clonal level correctly. PI and DMiDs are very helpful is the treatment of frontline as well as in relapse. Now days next generation sequencing technologies have come. With that researchers started to unravels the genomic landscape of this disease. This many portions of this fields are still unraveled in case of relapse and refractory MM and in coming future this may lead to personalized treatment care for MM. (21)

According to NCCN Guidelines Version 4. 2021,

Relapse or Progressive disease —————> Therapy, Clinical Trial, Autologous/ allogenic
Hematopoietic Cell transplant————> Refractory disease with lack of treatment options —————>
Palliative Care

Prognostic Factor:

Among the relapsed patient, the challenge is to select the best approach for each patient with the balance in efficacy and toxicity. Still now the biomarker-based treatment approach is limited. In a relapse and refractory situation, what medication a clinician can give is still variable and depends on the prior treatment regimen and previous treatment resistance. The prognostic value of chromosomal abnormalities in del13q, t (4,14), del17p, t (14,16), t (14,20) and gain of 1q21 are taken as high risk cytogenetic abnormalities. Apart from this according to International staging system (ISS) high serum albumin, serum beta 2 microglobulin and a high serum lactate dehydrogenase (LHD) count, presence of extra-medullary disease, presence of circulating plasma cell shows rapid progression on relapse during or after the treatment. (22,23)

Avet-Loiseau et al showed that the presence of t(4,14) and del 17p defines by FISH shows different event free survival and overall survival within each ISS stage.(24) The Mayo Stratification of Myeloma and risk adapted therapy(mSMART) criteria shows high risk, intermediate risk and standard risk condition.(25)

High Risk: FISH: del17p, t(14,16),t(14,20)

GEP: high risk signature

Intermediate risk: FISH:t(4,14)

Cytogenetics del13

Hypodiploidy with PCLI greater than equal to 3%

Standard Risk: FISH t(11,14),t(6,14)

Table 2. mSMART(25)

Proliferation of malignant plasma cell can be measured by plasma cell labeling index and Ki67 antigen expression (26,27). In the relapse and refractory MM patients undergoing bortezomib-based treatment, the Canadian community registered an amplification of 1q21 which is associated with a slightly worse outcome and no statistical variation in OS was found for del13q, del17p, t (4,14) or del1p21. (28) In a post-hoc sub analysis of an expanded access program treatment with lenalidomide-dexamethasone overcome the poor prognosis conferred by deletion of chromosome 13q or t(4,14) with similar median TTP or OS to patient without this chromosomal abnormality.(29) With another Lenalidomide based combination, the overall response rate (ORR) was similar for the patients with or without del 13q or t(4,14) but in case of del17p the response is poor and significantly shorter median TTP with respective of the patient without cytological abnormality (30)

Bruton Kinase (BTK) regulates many vital signal regulating pathways. A small molecule KS99 in a tubulin polymerization inhibitor. This in vivo inhibits the tumor growth in MM by inhibiting BTK. (31)

In MM the prognostic score is yet unknown. The Gene expression profiling (GEP) may identify a poor outcome patient. ECM92, UAMS70, UAMS80 gene signature helps to detect the prognosis of MM. (32,33). Cytogenetics is important in case of MM frontline treatment as well as relapse and refractory treatment. In this treatment a combination therapy is needed. Lenalidomide, pomalidomide, PI based combination shows significant prognosis. Pomalidomide based regimen is better than Lenalidomide based regimen in case of 17p deletion patient and pomalidomide and PI is now the best approach towards to the relapse case of MM. Cytogenetics is very important and stem cell transplantation, donor lymphocytic infusion can be very effective in current future. But the most important treatment which will come in future that is personalized genetic treatment and gene correction therapy. In a study it has been shown that CD 226 will play a key role in future regarding MM treatment, Action of CD137 and mAB(4-1BB) shows strong anti-myeloma activity while inhibition of PD1 and CTLA-4 shows no effect. (34)

Treatment Indication and Treatment selection in Relapse and refractory MM

The biochemical relapse is defined as the significant increase amount of M protein. Amount of M protein will be more than equal to 10mg/L or in urine 500mg/24hours. Clinical relapse is defined as CRAB criteria (hyper calcemic, renal failure, anemia and bone deformity) and increase in plasmacytomas. It can also be defined as increase in free light chain or FLC of more than 200 mg/L in a 2 months' interval test. (35)

Treatment should be considered on the basis of high-risk stratification and the regimen of drug which is used in the frontline setting.

The key drug for the treatment of diagnostic as well as relapse and refractory MM is bortezomib which is approved by the FDA in May 13, 2003. In July 1998 Thalidomide, immunomodulatory drug, got the approval to use in multiple myeloma patient. Lenalidomide, another immunomodulatory drug got its FDA approval in 27 December 2005. Carfilzomib is approved in 20 July 2012 by FDA. Pomalidomide got FDA approval in 2013. In November 21, 2016 Daratumumab is approved by FDA.

Treatment selection is very important in relapse and refractory MM. The age and the previous regimen of drug used in frontline treatment is important. Palumbo A et al (36) describes a scoring system (0-5) to identify 3 groups of patients on the basis of age, comorbidities, cognition and physical condition. Fit is score 0, Intermediate fitness is score 1 and frail is score 2 or more. The overall survival is most for fit patient, intermediate for intermediate fit patient and lowest is for other patients. This score predicts the mortality and chance of toxicity is higher in the elderly patients (36). The fit patient can get the full regimen of 3 drugs and frail patient will get the double drug regimen and dose reduction according to the guidelines provided. (7) The motto of the treatment is different to these groups. The motto of fit patient is to go towards complete remission but in case of frail the primary motto is palliativeness i.e. to provide a good quality of life. (36,37,38)

Currently in clinical practice, physician use to suggest different regimen, not the same regimen provided in the frontline therapy. Although several studies reported the retreatment with Bortezomib which gives an overall response rate (ORR) of 40% and time to progression (TTP) of 8.4 months with a common adverse effect of thrombocytopenia (grade 3-4) (39)

In 2010, Bortezomib, a first call PI and two DMiDs thalidomide and lenalidomide are the backbone of the relapse and refractory multiple myeloma. In 2010, National Comprehensive cancer network(NCCN) guidelines recommended the treatment of relapsed and refractory MM by using single agent of bortezomib or Lenalidomide or regimen with bortezomib-dexamethasone, bortezomib- doxorubicin, lenalidomide- dexamethasone, lenalidomide-bortezomib-dexamethasone.(40) With the Bortezomib, lenalidomide and thalidomide regimen the response rate goes up to 85% (41) In 2010, Sundar Jagannath et al described in their paper that use of PI like bortezomib with the use of histone deacetylase inhibitor may be a valuable treatment for relapse and refractory MM. (42) In 2010, Melphalan, Prednisone, lenalidomide, thalidomide regimen was used to treat relapse and refractory MM where 1 year PFS was 51% and 1 year OS was 72% with a grade 4 hematologic adverse events neutropenia 18%, thrombocytopenia 7%, anemia 2% and grade 3 non hematologic adverse effect infection 14%, neurological toxicity 4.5% and fatigue 7% is seen.(43)

A Neils WC paper published in 2011 indicated that therapy with lenalidomide is the safest option for relapsed and refractory neuropathy-associated MM, and bortezomib demonstrates maximum effectiveness in patients with renal insufficiency rather than thromboembolism risk. (44) It is also mentioned in this paper that allogeneic stem cell transplantation can be offered to certain patients if autologous stem cell transplantation fails. (44) In Relapsed case melphalan associated autologous stem cell transplantation can be used. Radiotherapy and osteoplastic procedure can be used adjunct to systemic therapy in case of Relapsed MM with osteolytic bone lesions. (45). In 2012, carfilzomib entered in the treatment regimen. Carfilzomib shows a good selectivity which arrests the cell growth and apoptosis. (46) It shows a median response of 7.8 months and median PFS of 3.7 months and median OS 15.6 months. (46) In case of Relapse and refractory MM Carfilzomib has well tolerability with low incidence of grade 3 or higher peripheral neuropathy. (46)

In 2013 Jesus Sam Meguel (47) described in their paper that Pomalidomide with low dose dexamethasone (oral regimen) can be the new treatment in the relapse and refractive MM. Their study shows a median PFS of 4.0 months over High dose dexamethasone treatment where median PFS was 1.4 months. The common adverse events showed are neutropenia, anemia thrombocytopenia. (47)

A study of Silvia Park et (published in 2013), Dexamethasone, cyclophosphamide, etoposide and cisplatin regimen was used to relapsed and refractory MM patient with a median age of 58 years and response rate was 45.1%. (48)

Bortezomib and lenalidomide refractory patient was treated with Carfilzomib mono therapy where ORR was 23.7% with median duration response was 7.8 months. (49) The most common adverse effects were thrombocytopenia, anemia, fatigue, diarrhea.

A study (PANORAMA 1) (50) has been done Panobinostat- bortezomib-dexamethasone to bortezomib-dexamethasone. Median follow up of Panobinostat group was 6.47 months whereas for control group 5.59 months. Median PFS was 11.9 months to 8.08 months for control group. Median OS was 33.64 months for Panobinostat group 30.39 months for control group.

Eloquent 2 Trial (51) (Elotuzumab-lenalidomide-dexamethasone to lenalidomide-dexamethasone) (n=646, elotuzumab group 321 and 325 in control group). 1year PFS in 68% for elotuzumab group and 57% for control group. 2 years PFS is 41% for elotuzumab group and 27% for control group. Median PFS is 19.4 month for elotuzumab group and 14.9 months for control group. ORR is 79% for elotuzumab group and 66% for control group. Common adverse effect was grade 3 or 4 lymphocytopenia, neutropenia fatigue or pneumonia. So for relapsed and refractory patient the Elotuzumab regimen shows a prolonged PFS as well as OS.

Tourmaline MM1 trial(52) (oral Ixazomib-lenalidomide-dexamethasone to lenalidomide-dexamethasone) (n=772)

Median progression free survival 20.6 months to Ixazomib group and 14.7 months to control group. ORR 78% to Ixazomib group to 72% to control group. Severe adverse effects occur to ixazomib group that is grade 4 thrombocytopenia. 27% patients show peripheral neuropathy in ixazomib group where 22% shows the same in control group. So, the PFS and ORR is better for the Ixazomib group but the adverse effects are more severe in case of Ixazomib group also.

Castor trial (53) (Daratumumab-bortezomib-dexamethasone to bortezomib-dexamethasone) (n=498) 12 months progression free survival is 60.7% in Daratumumab group and 26.9% for control group. ORR is 82.9% in Daratumumab group to 63.2% in the control group and complete response is 19.2% in daratumumab group and 9.0% to control group. Most severe adverse effect is seen in grade 3 or 4 thrombocytopenia, anemia and neutropenia. So daratumumab group has a prolonged PFS, ORR and CSS than only bortezomib-dexamethasone regimen.

Pollux trial (54) (Daratumumab-Lenalidomide-Dexamethasone to Lenalidomide-Dexamethasone) (n=569). 12 months PFS was 83.2% in Dara group where as 60.1% in control group. ORR is 93% in Dara group to 76.4% in control group. ($p<0.001$) Most common adverse effect was grade 3 or 4 neutropenia (51.9% in Dara group), thrombocytopenia and anemia. Daratumumab based treatment regimen has shown better PFS, ORR.

ASPIRE Trial (55) (Carfilzomib-lenalidomide-dexamethasone to Lenalidomide-dexamethasone) Median OS was 48.3 months for Carfilzomib group to 40.4 months for control group. In patient who received one prior line of therapy median OS was 11.4 months longer for Carfilzomib over control group and 6.5 months longer who received two or more than two prior line therapies.

Carfilzomib group shows a higher OS and lesser death with an increase in median survival of 7.9 months. Carfilzomib regimen is good for first relapse.

Carfilzomib-lenalidomide-dexamethasone has a favorable benefit risk profile for patients with relapse and refractory MM, even with more than 70years old patients (56)

Endeavor Trial (57) (n= 929, 464 in carfilzomib group and 465 to bortezomib group) (Carfilzomib-dexamethasone to Bortezomib-dexamethasone) Median OS was 47.6 months in carfilzomib group and 40 months in bortezomib group. Carfilzomib patient shows serious adverse effect than bortezomib group like anemia, pneumonia and hypertension, thrombocytopenia (Grade 3).

OPTIMISMM trial (58) (Pomalidomide-bortezomib-dexamethasone to Bortezomib dexamethasone) (n=559, 281 are on pomalidomide group and 279 are in control group) Median PFS is 11.2 months for pomalidomide group and 7.1 is for controlled group. Most common adverse effects are grade 3-4 neutropenia, infection and thrombocytopenia. This study suggests that pomalidomide-bortezomib- dexamethasone can be an effective treatment in lenalidomide refractory cases.

In lenalidomide refractory cases physician can use bortezomib, pomalidomide regimen or pomalidomide plus low dose dexamethasone regimen. (59). In lenalidomide refractory case isatuximab-pomalidomide-dexamethasone can be used. (60) In Relapse and Refractory MM treatment histone deacetylase inhibitor, kinetin spindle protein inhibitor and mTOR inhibitors can be used (44).

In MM many immune mechanism are involved including reduced expression of tumor antigen and human leukocyte antigen (HLA) expression of inhibitory ligands like program cell death 1 (PDL1) by plasma cell and TREG cell and myeloid derived suppressor cell can contribute in the immune suppression (61-73) So to treat MM, immune based therapy is also important like monoclonal antibody, Chimeric antigen receptor (CAR-T), checkpoint inhibition and vaccination is important.

CAR T Cell therapy in Relapsed and Refractory Multiple Myeloma:

A preclinical study suggests that bb 2121, a CAR T treatment that inhibits B cell maturation antigen (BCMA) and a future treatment for relapsed and refractory MM.(74) Early phase of CAR-T treatment shows a promising result in the relapse and refractive MM.(75) So the mechanism of CAR T treatment is to targeting BCMA and kill the MM cells expressing BCMA.(76)

Checkpoint inhibitors:

Checkpoint inhibitors are the inhibitors of PD L1, PD L2. But PD L1 inhibition with immunomodulatory drugs shows severe toxicity in the treatment. (77) In initial trial AMG 420

shows a good response in case of relapse and refractory MM. (77). Many MM cells show PD L1 positive, still now more research is needed in the checkpoint inhibition treatment regarding relapse and refractory MM.

So, CAR-T treatment, Immune checkpoint and Vaccines are the future of the treatment of Relapse and refractory MM.

Current treatment in case of Lenalidomide refractory cases:

CANDOR: (78) (Carfilzomib-daratumumab-dexamethasone to carfilzomib-dexamethasone) (n=466) This regimen can be used in lenalidomide and bortezomib refractory. The daratumumab group have prolonged PFS than controlled group in case of relapse and refractory MM.

ICARIA MM: (79) (Isatuximab-pomalidomide-low dose dexamethasone to pomalidomide-low dose dexamethasone) (n=307, 154 to isatuximab group and 153 to controlled group) median PFS was 11.5 months to isatuximab compared to 6.5 months to controlled group. Most common adverse reaction was infusion reaction and upper respiratory tract infection.

Eloquent-3 Trial (80) (elotuzumab-pomalidomide-dexamethasone to pomalidomide-dexamethasone) (n=117, 60 to elotuzumab group, 57 to controlled group) median progression free survival is 10.3 months for elotuzumab group compared to 4.3 for controlled group. ORR is 53% to elotuzumab group compared to 26% in controlled group. In adverse effect grade 3-4 neutropenia, anemia and hyperglycemia is seen.

IKEMA trial: (81) (Isatuzumab-carfilzomib-dexamethasone to carfilzomib-dexamethasone) this trial shows a higher OSS (87% vs 83%), VGPR (72% vs 56%) CR (29% vs 13%)

Table 3: Regimen for Relapsed and refractory Multiple Myeloma (NCCN Ver 4.2021)

Preferred Regimen:

- Bortezomib/Lenalidomide/Dexamethasone
- Carfilzomib/Lenalidomide/Dexamethasone
- Daratumumab/Bortezomib/Dexamethasone
- Daratumumab/Carfilzomib/Dexamethasone
- Daratumumab/Lenalidomide/Dexamethasone
- Isatuximab-irfc/Pomalidomide/Dexamethasone
- Ixazomib/Lenalidomide/Dexamethasone
- Ixazomib/Pomalidomide/Dexamethasone
- Pomalidomide/Bortezomib/Dexamethasone

Other Recommended Regimen:

- Belantamab Mafodotin-blmf

- Bendamastine/ Bortezomib/Dexamethasone
- Bendamastine / Lenalidomide/ Dexamethasone
- Bortezomib//Liposomal Doxorubicin/ Dexamethasone
- Bortezomib/ Cyclophosphamide/ Dexamethasone
- Carfilzomib/Cyclophosphamide/Dexamethasone
- Carfilzomib (Twice Weekly)/Dexamethasone
- Cyclophosphamide/Lenalidomide/Dexamethasone
- Daratumumab/ Cyclophosphamide/ Bortezomib/Dexamethasone
- Daratumumab/Pomalidomide/Dexamethasone
- Elotuzumab/ Bortezomib/Dexamethasone
- Elotuzumab/Lenalidomide/Dexamethasone
- Elotuzumab/pomalidomide/Dexamethasone
- Ixazomib/Cyclophosphamide/Dexamethasone
- Panobinostat/Bortezomib/Dexamethasone
- Pomalidomide/Cyclophosphamide/Dexamethasone
- Pomalidomide/Carfilzomib/Dexamethasone
- Selinexor/ Bortezomib/Dexamethasone

Useful in Certain Circumstances:

- Bendamustine
- Bortezomib/Dexamethasone
- Carfilzomib/Cyclophosphamide/Thalidomide/Dexamethasone
- Carfilzomib/ Dexamethasone
- Daratumumab
- Dexamethasone/Cyclophosphamide/Etoposide/Cisplatin
- Dexamethasone/Thalidomide/Cisplatin/Doxorubicin/Cyclophosphamide/etoposide
- High dose Cyclophosphamide
- Ixazomib/Dexamethasone
- Lenalidomide/Dexamethasone
- Panobinostat/ Carfilzomib
- Panobinostat/Lenalidomide/Dexamethasone
- Pomalidomide/Dexamethasone
- Selinexor/Dexamethasone
- Venetoclax /Dexamethasone
- Selinexor/ Daratumumab/ Dexamethasone
- Selinexor/ Pomalidomide/ Dexamethasone

Fig 1: Mechanism of Actions of the Drugs used in the treatment of RRMM

*The picture is reconstructed from the paper of C.S Chim et. al. "Management of relapsed and refractory multiple myeloma: novel agents, antibodies, immunotherapies and beyond" Published in Leukaemia. 2018 Feb; 32(2): 252–262.

Transplant in Relapsed and Refractory Multiple Myeloma (RRMM)

Emerging drugs, such as inhibitors of deacetylase, monoclonal antibodies, and new proteasome inhibitors, look exciting and may modify the RRMM therapeutic environment. The benefit of ASCT rescue in patients with RRMM has been seen in a small number of trials, but there is continuing discussion and debates regarding its timing and efficacy. (82) A conversation on the timing and potential role of salvage ASCT and allogeneic stem cell transplantation in RRMM has been re-ignited with increase in transplant performance.

In a study, (83) it's reported that 64 stem cell patients were obtained within 1 year of diagnosis, none of whom had a transplant at the plateau level of their illness. Patients who were seen within 12 months of diagnosis underwent four cycles of regular chemotherapy with vincristine, doxorubicin, and dexamethasone (VAD) and then mobilized stem cells. Patients were then put on 12-cycle maintenance chemotherapy with vincristine, BCNU, melphalan, cyclophosphamide, and prednisone or melphalan and prednisone.

Another Study (84) concluded that the role of allogenic transplant is controversial. The introduction into transplant protocols of proteasome inhibitors and immunomodulatory drugs has

the ability to enhance outcomes and optimize the function of allogeneic HCT for the treatment of multiple myeloma as a platform for long-term management of diseases.

Salvage ASCT has been shown to be a successful replacement following reinduction therapy in several retrospective trials of patients who relapsed from a previous ASCT, which found that the most significant factor affecting PFS after salvage ASCT was the period of remission from the first ASCTT. (86,87) According a study of Sergio Giralt, it has been mentioned that transplantation is very effective whose PFS is more than 18 months. (86)

In a study, RRMM patients, who achieved at least stable disease after re-induction of bortezomib/ doxorubicin/dexamethasone were randomized to receive salvage ASCT ($N=89$) or weekly oral cyclophosphamide ($N=85$) as consolidation, which showed a superior time to disease progression (19 months versus 11 months, HR: 0.45; $P<0.0001$), and recently OS (67 months versus 52 months, HR: 0.56; $P=0.022$) in those receiving salvage ASCT compared with oral cyclophosphamide consolidation. (87)

So, timing and regimen of Transplantation is still under controversy and more Clinical trial is needed on transplant eligible candidates.

Table 4: Preferred therapy of Transplant Candidates according to guidelines of NCCN 4.2021

Preferred regimen: Bortezomib/Lenalidomide/Dexamethasone and
Bortezomib/Cyclophosphamide/Dexamethasone

Other Recommended Regimen:

- Carfilzomib/ Lenalidomide/Dexamethasone
- Daratumumab/Lenalidomide/Dexamethasone
- Ixazomib/Lenalidomide/Dexamethasone

Useful in certain circumstances:

- Bortezomib/Doxorubicin/Dexamethasone
- Carfilzomib/Cyclophosphamide/Dexamethasone
- Ixazomib/Cyclophosphamide/Dexamethasone
- Bortezomib/ thalidomide/ dexamethasone
- Cyclophosphamide/Lenalidomide/Dexamethasone
- Daratumumab/ Cyclophosphamide/Bortezomib/Dexamethasone
- Daratumumab/Bortezomib/Thalidomide/Dexamethasone
- Dexamethasone/Thalidomide/Cisplatin/ Doxorubicin/Cyclophosphamide/ Etoposide/ Bortezomib

Maintenance Therapy:

- Preferred: Lenalidomide
- Other Recommended: Ixazomib and Bortezomib
- Useful in certain situation: Bortezomib/Lenalidomide

Conclusion:

The choice of treatment regimen for relapsed and refractory multiple myeloma is always very difficult. Many research trials show different results and choosing a best treatment for a particular patient is really challenging but day by day the PFS and OS is improving for both high risk and low risk patient. PI, Immunomodulators are showing a good response in the relapse and refractory MM. Other Immune checkpoint inhibitors, CAR-T and Vaccines are on research and they are the future of the treatment. Stem cell transplantation is used in relapsed and refractive MM treatment and Donor lymphocytic infiltration can be the future of the treatment also.

Targeted therapies are important in BRAF mutated MM (BRAF inhibitor vemurafenib) and BCL2 mutated MM (venetoclax). Lenalidomide refractory cases are very challenging and many clinical trials is going on and pomalidomide showing an impressive result. In future many more drugs will come. The main moto of this treatment regimen are giving a good efficacy and safety in an affordable cost with low adverse effect. In this review we have seen the better the drug regimen is, higher is the severity of adverse effects. So, in future studies researchers should consider adverse effects also more deeply and strictly. So, the final goal should be to find a balance among efficacy, toxicity, cost and achieving a long lasting control on the disease and eventually cure to a subset of patients.

Acknowledgement:

We want to acknowledge Jinan University and Chittaranjan National Cancer Institute and special thanks to Sastha Sahay for supporting us.

Funding: No funding is present for this research.

Conflict of Interest: no conflict of interest.

Authors Contribution: KM and UC is the project head. DM has wrote the manuscript under guidance of KM and UM.

Reference:

1. Anderson, Kenneth C. "The 39th David A. Karnofsky Lecture: bench-to-bedside translation of targeted therapies in multiple myeloma." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* vol. 30,4 (2012): 445-52. doi:10.1200/JCO.2011.37.8919
2. Improved survival in multiple myeloma and the impact of novel therapies. Kumar SK, Rajkumar SV, Dispenzieri A, Lacy MQ, Hayman SR, Buadi FK, Zeldenrust SR, Dingli D, Russell SJ, Lust JA, Greipp PR, Kyle RA, Gertz MA *Blood*. 2008 Mar 1; 111(5):2516-20.

3. Barosi G, Merlini G, Billio A, et al. SIE, SIES, GITMO evidence-based guidelines on novel agents (thalidomide, bortezomib, and lenalidomide) in the treatment of multiple myeloma. *Ann Hematol*. 2012;91(6):875-888. doi:10.1007/s00277-012-1445-y
4. Rajkumar, S Vincent. "Multiple myeloma: Every year a new standard?." *Hematological oncology* vol. 37 Suppl 1,Suppl 1 (2019): 62-65. doi:10.1002/hon.2586
5. Kumar SK, Dimopoulos MA, Kastiris E, et al. Natural history of relapsed myeloma, refractory to immunomodulatory drugs and proteasome inhibitors: a multicenter IMWG study. *Leukemia*. 2017;31(11):2443–8.
6. Rajkumar SV, Kyle RA. Progress in Myeloma - A Monoclonal Breakthrough. *The New England journal of medicine* 2016;375:1390–2.
7. Nijhof, I.S., van de Donk, N.W.C.J., Zweegman, S. et al. Current and New Therapeutic Strategies for Relapsed and Refractory Multiple Myeloma: An Update. *Drugs* 78, 19–37 (2018).
<https://doi.org/10.1007/s40265-017-0841-y>
8. Durie BG, Harousseau JL, Miguel JS, et al. International uniform response criteria for multiple myeloma. *Leukemia*. 2006;20:1467–1473. doi: 10.1038/sj.leu.2404284.
9. Anderson KC, Kyle RA, Rajkumar SV, Stewart AK, Weber D, Richardson P. Clinically relevant end points and new drug approvals for myeloma. *Leukemia*. 2008;22:231–239. doi: 10.1038/sj.leu.2405016.
10. Silvia Gentili, Sagar Lonial Relapsed and Refractory Multiple Myeloma new therapeutic strategy, *hemtato oncol clin N Am* 28(2014) 861-890
11. Corre J, Cleyne A, Robiou du Pont S, et al. Multiple myeloma clonal evolution in homogeneously treated patients. *Leukemia*. 2018;32(12):2636-2647. doi:10.1038/s41375-018-0153-6
12. Keats JJ, Chesi M, Egan JB, et al. Clonal competition with alternating dominance in multiple myeloma. *Blood*. 2012;120:1067–76
13. Bolli N, Avet-Loiseau H, Wedge DC, et al. Heterogeneity of genomic evolution and mutational profiles in multiple myeloma. *Nat Commun*. 2014;5:2997.
14. Chavan SS, He J, Tytarenko R, et al. Bi-allelic inactivation is more prevalent at relapse in multiple myeloma, identifying RB1 as an independent prognostic marker. *Blood Cancer J*. 2017;7:e535.
15. Kortuem KM, Braggio E, Bruins L, Barrio S, Shi CS, Zhu YX, Tibes R, Viswanatha D, Votruba P, Ahmann G, Fonseca R, Jedlowski P, Schlam I, Kumar S, Bergsagel PL, Stewart AK. Panel sequencing for clinically oriented variant screening and copy number detection in 142 untreated multiple myeloma patients. *Blood Cancer J*. 2016;6:e397.
16. Weinhold N, Ashby C, Rasche L, et al. Clonal selection and double-hit events involving tumor suppressor genes underlie relapse in myeloma. *Blood*. 2016;128:1735–44.

17. Biological and therapeutic impact of intratumor heterogeneity in cancer evolution. McGranahan N, Swanton C *Cancer Cell*. 2015 Jan 12; 27(1):15-26.
18. Ding L, Ley TJ, Larson DE, Miller CA, Koboldt DC, Welch JS, et al. Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* 2012. January 11; 481(7382): 506–510.
19. Weinhold N, Ashby C, Rasche L, Chavan SS, Stein C, Stephens OW, et al. Clonal selection and double-hit events involving tumor suppressor genes underlie relapse in myeloma. *Blood* 2016. September 29; 128(13): 1735–1744.
20. Walker BA, Wardell CP, Melchor L, Brioli A, Johnson DC, Kaiser MF, et al. Intracлонаl heterogeneity is a critical early event in the development of myeloma and precedes the development of clinical symptoms. *Leukemia* 2014. February; 28(2): 384–390.
21. Rollig C, Knop S, Bornhauser M. Multiple myeloma. *Lancet*. 2015;385:2197–208
22. Mikhael JR, Dingli D, Roy V, et al. Management of newly diagnosed symptomatic multiple myeloma: updated Mayo Stratification of Myeloma and Risk-Adapted Therapy (mSMART) consensus guidelines 2013. *Mayo Clin Proc*. 2013;88:360–76.
23. Moreau P, Cavo M, Sonneveld P, et al. Combination of international scoring system 3, high lactate dehydrogenase, and t(4;14) and/or del(17p) identifies patients with multiple myeloma (MM) treated with front-line autologous stem-cell transplantation at high risk of early MM progression-related death. *J Clin Oncol*. 2014;32:2173–80.
24. Avet-Loiseau, Attal M, Moreau et al genetic abnormalities and survival in multiple myeloma: the experience of the intergroupe Francophone du myeloma. *Blood* 2007;109(8):3489-95
25. Michel JR, dingli D, Roy V et al, management of newly diagnosed symptomatic multiple myeloma: updated mayo stratification of myeloma and risk adapted therapy(mSMART) consensus guidelines 2013. *Mayoclinic proc* 2013;88(4):360-76
26. Greipp PR, Kumar S. Plasma cell labeling index. *Methods Mol Med*. 2005;113:25–35.
27. Gastinne T, LeLeu X, Duhamel A, et al. Plasma cell growth fraction using Ki-67 antigen expression identifies a subgroup of multiple myeloma patients displaying short survival within the ISS stage I. *Eur J Haematol*. 2007;79:297–304.
28. Chang H, Trieu Y, Qi X yet al. Impact of cytogenetics in patient with relapsed or refractory multiple myeloma treatmed with bortezomib: adverse effect of 1q21gains. *Leuk Res* 2011;35(1): 95-8
29. Chen C, Reece DE, Siegel Det al. Expanded safety experience with linalidomide plus dexamathasone in relapsed or refractory myeloma. *BrJHaematol* 2009,146(2):164-70

30. Knop S, Gerecke C, Liebisch P, et al. linalidomide, adriamycine and dexamethasone(RAD) in patient with relapse and refractory multiple myeloma: a report from german myeloma study group DSMM(Deutsche Studiengruppe Multiples Myelom) Blood 2009;113(18): 4137-43
31. Pandey MK, Gowda K, Sung SS, et al. A novel dual inhibitor of microtubule and Bruton's tyrosine kinase inhibits survival of multiple myeloma and osteoclastogenesis. Exp Hematol. 2017;53:31-42. doi:10.1016/j.exphem.2017.06.003
32. Kuiper R, Broyl A, de Knecht Y, et al. A gene expression signature for high-risk multiple myeloma [published correction appears in Leukemia. 2014 May;28(5):1178-80. doi: 10.1038/leu.2014.53.]. Leukemia. 2012;26(11):2406-2413. doi:10.1038/leu.2012.127
33. van Laar, Ryan et al. "Translating a gene expression signature for multiple myeloma prognosis into a robust high-throughput assay for clinical use." BMC medical genomics vol. 7 25. 17 May. 2014, doi:10.1186/1755-8794-7-25
34. Guillerey, Camille et al. "Immunosurveillance and therapy of multiple myeloma are CD226 dependent." The Journal of clinical investigation vol. 125,5 (2015): 2077-89. doi:10.1172/JCI77181
35. Rajkumar SV, Harousseau JL, Durie B, et al. Consensus recommendations for the uniform reporting of clinical trials: report of the International Myeloma Workshop Consensus Panel 1. Blood. 2011;117:4691-5.
36. Palumbo, Antonio et al. "Geriatric assessment predicts survival and toxicities in elderly myeloma patients: an International Myeloma Working Group report." Blood vol. 125,13 (2015): 2068-74. doi:10.1182/blood-2014-12-615187
37. Engelhardt M, Dold SM, Ihorst G, Zober A, Moller M, Reinhardt H, Hieke S, Schumacher M, Wasch R. Geriatric assessment in multiple myeloma patients: validation of the International Myeloma Working Group (IMWG) score and comparison with other common comorbidity scores. Haematologica. 2016;101:1110-9.
38. Larocca A, Palumbo A. How I treat fragile myeloma patients. Blood. 2015;126:2179-85.
39. M. T. Petrucci, P. Giraldo, P. Corradini et al., "A prospective, international phase 2 study of bortezomib retreatment in patients with relapsed multiple myeloma," British Journal of Haematology, vol. 160, no. 5, pp. 649-659, 2013.
40. Richardson PG, Laubach J, Mitsiades C, et al. Tailoring treatment for multiple myeloma patients with relapsed and refractory disease. Oncology (Williston Park). 2010;24(3 Suppl 2):22-29.
41. Richards T, Weber D. Advances in treatment for relapses and refractory multiple myeloma. Med Oncol. 2010;27 Suppl 1:S25-S

42. doi:10.1007/s12032-009-9407-5 42. Jagannath S, Dimopoulos MA, Lonial S. Combined proteasome and histone deacetylase inhibition: A promising synergy for patients with relapsed/refractory multiple myeloma. *Leuk Res.* 2010;34(9):1111-1118. doi:10.1016/j.leukres.2010.04.001
43. Palumbo A, Larocca A, Falco P, et al. Lenalidomide, melphalan, prednisone and thalidomide (RMPT) for relapsed/refractory multiple myeloma. *Leukemia.* 2010;24(5):1037-1042. doi:10.1038/leu.2010.58
44. van de Donk NW, Lokhorst HM, Dimopoulos M, et al. Treatment of relapsed and refractory multiple myeloma in the era of novel agents. *Cancer Treat Rev.* 2011;37(4):266- 283. doi:10.1016/j.ctrv.2010.08.008
45. Moehler T, Goldschmidt H. Therapy of relapsed and refractory multiple myeloma. *Recent Results Cancer Res.* 2011;183:239-271. doi:10.1007/978-3-540-85772-3_11
46. McCormack PL. Carfilzomib: in relapsed, or relapsed and refractory, multiple myeloma. *Drugs.* 2012;72(15):2023-2032. doi:10.2165/11209010-000000000-00000
47. Miguel JS, Weisel K, Moreau P, et al. Pomalidomide plus low-dose dexamethasone versus high-dose dexamethasone alone for patients with relapsed and refractory multiple myeloma (MM-003): a randomised, open-label, phase 3 trial. *Lancet Oncol.* 2013;14(11):1055-1066. doi:10.1016/S1470-2045(13)70380-2
48. Park S, Lee SJ, Jung CW, et al. DCEP for relapsed or refractory multiple myeloma after therapy with novel agents. *Ann Hematol.* 2014;93(1):99-105. doi:10.1007/s00277-013- 1952-5
49. Thompson JL. Carfilzomib: a second-generation proteasome inhibitor for the treatment of relapsed and refractory multiple myeloma. *Ann Pharmacother.* 2013;47(1):56-62. doi:10.1345/aph.1R561
50. San-Miguel JF, Hungria VT, Yoon SS, et al. Panobinostat plus bortezomib and dexamethasone versus placebo plus bortezomib and dexamethasone in patients with relapsed or relapsed and refractory multiple myeloma: a multicentre, randomised, double-blind phase 3 trial [published correction appears in *Lancet Oncol.* 2015 Jan;16(1):e6]. *Lancet Oncol.* 2014;15(11):1195-1206. doi:10.1016/S1470-2045(14)70440-1
51. Lonial S, Dimopoulos M, Palumbo A, et al. Elotuzumab Therapy for Relapsed or Refractory Multiple Myeloma. *N Engl J Med.* 2015;373(7):621-631. doi:10.1056/NEJMoa1505654
52. Moreau P, Masszi T, Grzasko N, et al. Oral ixazomib, lenalidomide, and dexamethasone for Multiple Myeloma. *N Engl J Med.* 2016;374(17):1621-1634. doi:10.1056/NEJMoa1516282
53. Palumbo A, Chanan-Khan A, Weisel K, et al. Daratumumab, Bortezomib, and Dexamethasone for Multiple Myeloma. *N Engl J Med.* 2016;375(8):754-766. doi:10.1056/NEJMoa1606038
54. Dimopoulos MA, Oriol A, Nahi H, et al. Daratumumab, Lenalidomide, and Dexamethasone for Multiple Myeloma. *N Engl J Med.* 2016;375(14):1319-1331. doi:10.1056/NEJMoa1607751

55. Siegel DS, Dimopoulos MA, Ludwig H, et al. Improvement in Overall Survival With Carfilzomib, Lenalidomide, and Dexamethasone in Patients With Relapsed or Refractory Multiple Myeloma. *J Clin Oncol*. 2018;36(8):728-734. doi:10.1200/JCO.2017.76.5032
56. Dimopoulos, Meletios A et al. "Carfilzomib, lenalidomide, and dexamethasone in patients with relapsed multiple myeloma categorised by age: secondary analysis from the phase 3 ASPIRE study." *British journal of haematology* vol. 177,3 (2017): 404-413. doi:10.1111/bjh.14549
57. Dimopoulos MA, Goldschmidt H, Niesvizky R, et al. Carfilzomib or bortezomib in relapsed or refractory multiple myeloma (ENDEAVOR): an interim overall survival analysis of an open-label, randomised, phase 3 trial [published correction appears in *Lancet Oncol*. 2017 Oct;18(10):e562]. *Lancet Oncol*. 2017;18(10):1327-1337. doi:10.1016/S1470-2045(17)30578-8
58. Richardson PG, Oriol A, Beksac M, et al. Pomalidomide, bortezomib, and dexamethasone for patients with relapsed or refractory multiple myeloma previously treated with lenalidomide (OPTIMISMM): a randomised, open-label, phase 3 trial. *Lancet Oncol*. 2019;20(6):781-794. doi:10.1016/S1470-2045(19)30152-4
59. Siegel DS, Schiller GJ, Song KW, et al. Pomalidomide plus low-dose dexamethasone in relapsed refractory multiple myeloma after lenalidomide treatment failure. *Br J Haematol*. 2020;188(4):501-510. doi:10.1111/bjh.16213
60. Cavo M. Facing lenalidomide-refractory myeloma. *Blood*. 2019;134(2):99-101. doi:10.1182/blood-2019-04-901157
61. Klippel ZK, Chou J, Towler AM, Voong LN, Robbins P, Bensinger WI, Warren EH. Immune escape from NY-ESO-1-specific T-cell therapy via loss of heterozygosity in the MHC. *Gene Ther*. 2014;21:337-42
62. Racanelli V, Leone P, Frassanito MA, Brunetti C, Perosa F, Ferrone S, Dammacco F. Alterations in the antigen processing-presenting machinery of transformed plasma cells are associated with reduced recognition by CD8 + T cells and characterize the progression of MGUS to multiple myeloma. *Blood*. 2010;115:1185-93.
63. Brown RD, Pope B, Yuen E, Gibson J, Joshua DE. The expression of T cell related costimulatory molecules in multiple myeloma. *Leuk Lymphoma*. 1998;31:379-84
64. Benson DM Jr, Bakan CE, Mishra A, Hofmeister CC, Efebera Y, Becknell B, Baiocchi RA, Zhang J, Yu J, Smith MK, Greenfield CN, Porcu P, Devine SM, Rotem-Yehudar R, Lozanski G, Byrd JC, Caligiuri MA. The PD-1/PD-L1 axis modulates the natural killer cell versus multiple myeloma effect: a therapeutic target for CT-011, a novel monoclonal antiPD-1 antibody. *Blood*. 2010;116:2286-94.
65. Liu J, Hamrouni A, Wolowiec D, Coiteux V, Kuliczowski K, Hetuin D, Saudemont A, Quesnel B. Plasma cells from multiple myeloma patients express B7-H1 (PD-L1) and increase expression after stimulation with IFN- γ and TLR ligands via a MyD88-, TRAF6-, and MEK-dependent pathway. *Blood*. 2007;110:296-304.

66. Paiva B, Azpilikueta A, Puig N, et al. PD-L1/PD-1 presence in the tumor microenvironment and activity of PD-1 blockade in multiple myeloma. *Leukemia*. 2015;29:2110–3.
67. Tamura H, Ishibashi M, Yamashita T, Tanosaki S, Okuyama N, Kondo A, Hyodo H, Shinya E, Takahashi H, Dong H, Tamada K, Chen L, Dan K, Ogata K. Marrow stromal cells induce B7-H1 expression on myeloma cells, generating aggressive characteristics in multiple myeloma. *Leukemia*. 2013;27:464–72.
68. Dosani T, Carlsten M, Maric I, Landgren O. The cellular immune system in myelomagenesis: NK cells and T cells in the development of myeloma [corrected] and their uses in immunotherapies. *Blood Cancer J*. 2015;5:e306.
69. Sze DM, Giesajtis G, Brown RD, Raitakari M, Gibson J, Ho J, Baxter AG, Fazekas de St GB, Basten A, Joshua DE. Clonal cytotoxic T cells are expanded in myeloma and reside in the CD8(+)CD57(+)CD28(-) compartment. *Blood*. 2001;98:2817–27.
70. Malek E, de Lima M, Letterio JJ, Kim BG, Finke JH, Driscoll JJ, Giralto SA. Myeloidderived suppressor cells: The green light for myeloma immune escape. *Blood Rev*. 2016;30:341–8.
71. Franssen LE, van de Donk NW, Emmelot ME, Roeven MW, Schaap N, Dolstra H, Hobo W, Lokhorst HM, Mutis T. The impact of circulating suppressor cells in multiple myeloma patients on clinical outcome of DLIs. *Bone Marrow Transpl*. 2015;50:822–8.
72. Giallongo C, Tibullo D, Parrinello NL, La CP, Di RM, Bramanti V, Di RC, Conticello C, Chiarenza A, Palumbo GA, Avola R, Romano A, Di RF. Granulocyte-like myeloid derived suppressor cells (G-MDSC) are increased in multiple myeloma and are driven by dysfunctional mesenchymal stem cells (MSC). *Oncotarget*. 2016;7:85764–75.
73. Lesokhin AM, Hohl TM, Kitano S, Cortez C, Hirschhorn-Cymerman D, Avogadri F, Rizzuto GA, Lazarus JJ, Pamer EG, Houghton AN, Merghoub T, Wolchok JD. Monocytic CCR2(+) myeloid-derived suppressor cells promote immune escape by limiting activated CD8 T-cell infiltration into the tumor microenvironment. *Cancer Res*. 2012;72:876–86.
74. Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-Cell Therapy bb2121 in Relapsed or Refractory Multiple Myeloma. *N Engl J Med*. 2019;380(18):1726-1737. doi:10.1056/NEJMoa1817226
75. Feinberg D, Paul B, Kang Y. The promise of chimeric antigen receptor (CAR) T cell therapy in multiple myeloma. *Cell Immunol*. 2019;345:103964. doi:10.1016/j.cellimm.2019.103964
76. Feng D, Sun J. Overview of anti-BCMA CAR-T immunotherapy for multiple myeloma and relapsed/refractory multiple myeloma. *Scand J Immunol*. 2020;92(2):e12910. doi:10.1111/sji.12910
77. Cohen AD. Myeloma: next generation immunotherapy. *Hematology Am Soc Hematol Educ Program*. 2019;2019(1):266-272. doi:10.1182/hematology.2019000068
78. Dimopoulos M, Quach H, Mateos MV, et al. Carfilzomib, dexamethasone, and daratumumab versus carfilzomib and dexamethasone for patients with relapsed or refractory multiple myeloma (CANDOR):

results from a randomised, multicentre, openlabel, phase 3 study. *Lancet*. 2020;396(10245):186-197. doi:10.1016/S0140- 6736(20)30734-0

79. Attal M, Richardson PG, Rajkumar SV, et al. Isatuximab plus pomalidomide and low-dose dexamethasone versus pomalidomide and low-dose dexamethasone in patients with relapsed and refractory multiple myeloma (ICARIA-MM): a randomised, multicentre, open-label, phase 3 study [published correction appears in *Lancet*. 2019 Dec 7;394(10214):2072]. *Lancet*. 2019;394(10214):2096-2107. doi:10.1016/S0140- 6736(19)32556-5

80. Dimopoulos MA, Dytfield D, Grosicki S, et al. Elotuzumab plus Pomalidomide and Dexamethasone for Multiple Myeloma. *N Engl J Med*. 2018;379(19):1811-1822. doi:10.1056/NEJMoa1805762

81. moreau eha 2020 abstract LB2603

82. Cornell RF, Kassim AA. Evolving paradigms in the treatment of relapsed/refractory multiple myeloma: increased options and increased complexity. *Bone Marrow Transplant*. 2016 Apr;51(4):479-91. doi: 10.1038/bmt.2015.307. Epub 2016 Jan 4. PMID: 26726946; PMCID: PMC4827007.

83. Gertz MA, Lacy MQ, Inwards DJ, Gastineau DA, Tefferi A, Chen MG, Witzig TE, Greipp PR, Litzow MR. Delayed stem cell transplantation for the management of relapsed or refractory multiple myeloma. *Bone Marrow Transplant*. 2000 Jul;26(1):45-50. doi: 10.1038/sj.bmt.1702445. PMID: 10918404.

84. Schneidawind C, Duerr-Stoerzer S, Faul C, Kanz L, Weisel K, Bethge W, Schneidawind D. Follow-up of patients with refractory or relapsed multiple myeloma after allogeneic hematopoietic cell transplantation. *Clin Transplant*. 2017 Jul;31(7). doi: 10.1111/ctr.12994. Epub 2017 May 26. PMID: 28470884.

85. Atanackovic D, Schilling G. Second autologous transplant as salvage therapy in multiple myeloma. *Br J Haematol* 2013; 163: 565–572. [[PubMed](#)] [[Google Scholar](#)]

86. Giralt S, Garderet L, Durie B, Cook G, Gahrton G, Bruno B et al. American Society of Blood and Marrow Transplantation, European Society of Blood and Marrow Transplantation, Blood and Marrow Transplant Clinical Trials Network, and International Myeloma Working Group Consensus Conference on salvage hematopoietic cell transplantation in patients with relapsed multiple myeloma. *Biol Blood Marrow Transplant* 2015; 21: 2039–2051. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

87. Cook G, Ashcroft AJ, Cairns DA, Williams CD, Brown JM, Cavenagh JD et al. The effect of salvage autologous stem-cell transplantation on overall survival in patients with relapsed multiple myeloma (final results from BSBMT/UKMF Myeloma X Relapse [Intensive]): a randomised, open-label, phase 3 trial. *Lancet Haematol* 2016; 3: e340–e351. [[PubMed](#)] [[Google Scholar](#)]