Intractable SARS-CoV-2 infection in a CAR-T-cell Therapy Recipient

Matthew K. Hensley¹, William G. Bain^{1,2,3}, Jana Jacobs⁴, Sham Nambulli^{5,6}, Urvi Parikh⁴, Kevin D. McCormick⁴, Anthony Cillo^{7,8}, Brittany Staines⁴, Amy Heaps⁴, Michele D. Sobolewski⁴, Linda J. Rennick^{5,6}, Georgios D. Kitsios^{1,2}, Barbara Methe², Ashwin Somasundaram^{7,9}, Tullia Bruno⁷, Carly Cardello⁷, Feng Shan⁷, Creg Workman⁷, Prabir Ray^{2,7}, Anuradha Ray^{2,7}, Janet Lee^{1,2,10}, Mark S. Ladinsky¹¹, Pamela J. Bjorkman¹¹, Dario A. Vignali⁷, W. Paul Duprex^{5,6}, Mounzer E. Agha⁹, John W. Mellors^{4,12}, Alison Morris^{1,2}, and Ghady Haidar^{4,12*}

¹Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Internal Medicine, UPMC, Pittsburgh, PA

²Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA

³Staff Physician, VA Pittsburgh Healthcare System, Pittsburgh, PA

⁴Division of Infectious Disease, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA

⁵Center for Vaccine Research, University of Pittsburgh School of Medicine, Pittsburgh, PA

⁶Department of Microbiology and Molecular Genetics, University of Pittsburgh School of Medicine, Pittsburgh, PA

⁷Department of Immunology, University of Pittsburgh School of Medicine, Pittsburgh, PA

⁸Tumor Microenvironment Center, UPMC Hillman Cancer Center, Pittsburgh, PA

⁹Division of Hematology, Oncology, Department of Internal Medicine, UPMC, Pittsburgh, PA

¹⁰Acute Lung Injury Center of Excellence, University of Pittsburgh, Pittsburgh, PA

¹¹Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA

¹²Division of Infectious Disease, Department of Internal Medicine, UPMC, Pittsburgh, PA

Keywords: COVID-19, Immunotherapy, Humoral Immunity, SARS-CoV-2 RNAemia, Severe Acute Respiratory Syndrome Coronavirus 2

Running Title: Persistent COVID-19 in a CAR-T-cell Therapy Recipient

***Correspondence:**

Ghady Haidar, MD

Division of Infectious Diseases, University of Pittsburgh and UPMC

3601 Fifth Ave, Falk Medical Building, Suite 5B

Pittsburgh, PA 15213

haidarg@upmc.edu

Office: 412-648-6601

Fax: 412-648-6399

Alternate corresponding author:

Matthew K Hensley, MD MPH

Division of Pulmonary & Critical Care Medicine, UPMC

5200 Centre Ave, Ste 610

Pittsburgh, PA 15232

hensleymk2@upmc.edu

Office: 412-621-1200

Funding Statement: Research reported in this publication was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number KL2TR001856 awarded to G.H., Career Development Award Number IK2 BX004886 from the United States Department of Veterans Affairs Biomedical Laboratory R&D (BLRD) Service awarded to W.B., NIH grant P50 8 P50 AI150464-13 (P.J.B.), George Mason University Fast

Grants (P.J.B.), Center for Vaccine Research/University of Pittsburgh, P01HL114453 (P.R, J.S.L.), R01 HL136143 (J.S.L.), and by the University of Pittsburgh Medical Center (UPMC) awarded to D.A.A.V. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Drs. Hensley and Bain contributed equally to this article. Drs. Jacobs and Nambulli contributed equally to this article. All authors have seen and approved the manuscript and contributed significantly to the work. The manuscript has not been previously published nor is it being considered for publication elsewhere.

Disclosures: GH receives research funds from Karius, Inc. JWM is a consultant to Gilead Sciences and Merck, and owns shares in Co-Crystal Pharmaceuticals, Infectious Disease Connect, and Abound Bio. GDK has received research funds from Karius, Inc. DAAV: cofounder and stock holder – Novasenta, Tizona and Potenza; stock holder – Oncorus and Werewolf; patents licensed and royalties - Astellas, BMS; scientific advisory board member - Tizona, Werewolf and F-Star; consultant - Astellas, BMS, Almirall, Incyte; research funding – BMS and Novasenta. The rest of the authors have no disclosures.

Acknowledgements: The authors would like to thank Dr. Rima Abdelmassih, Michelle Busch, Dr. Jennifer McComb, Dr. Bryan McVerry, Heather Michael, Dr. Stephanie Mitchell, John Ries, Caitlin Schaefer, Dr. Michael Shurin, Dr. Alan Wells, and Dr. Sarah Wheeler.

Word Count: 1997

Tables/Figures:3

References: 20

Abstract:

Background: Patients with B-cell aplasia, such as recipients of CAR-T-cell therapy for hematological malignancies, may be at risk for protracted SARS-CoV-2 replication and chronic shedding of infectious virus.

Methods: A man in his 70's who had undergone CAR-T-cell therapy presented with COVID-19 pneumonia and improved after receipt of convalescent plasma and remdesivir. He was discharged on room air but returned on day 41 after COVID-19 diagnosis with progressive radiographic abnormalities and persistent detection of SARS-CoV-2 RNA from nasopharyngeal swabs. He died due to progressive respiratory failure on day 74 of illness. Plasma and endotracheal aspirate (ETA) samples were collected for viral quantification, viral titration and culture, and immunological assays.

Findings: The patient had high-grade SARS-CoV-2 RNAemia in plasma collected on days 4, 9, 13, 16, and 71. RNAemia peaked at $>10^5$ copies/ml, declined after administration of remdesivir and convalescent plasma (10^4 copies/ml), but subsequently rebounded to $>10^5$ copies/ml before death. On day 72, ETA showed 2.78×10^{10} copies/mL of SARS-CoV-2 RNA, high-titer infectious virus (1.125×10^6 plaque-forming units/mL) that was successfully cultured, and numerous virus particles by electron microscopy. Serologic assays showed minimal to no antibody to SARS-CoV-2, and B-cells were not detected by immunophenotyping or by single-cell RNAseq.

Interpretation: This case provides new insights into the potential prolonged duration and extent of viral replication without appropriate humoral immune responses. This case also challenges the belief that persistent detection of SARS-CoV-2 RNA in respiratory secretions represents shedding of non-infectious RNA, which should prompt re-evaluation of infection prevention guidelines in immunosuppressed hosts.

Introduction:

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in patients with hematologic malignancies results in frequent coronavirus disease 2019 (COVID-19)-related acute respiratory distress syndrome (ARDS) and death^{1,2}. Chimeric antigen-modified T-cell therapy (CAR-T-cell) recipients are at risk for severe COVID-19 because of their unique immune defects. Such patients experience chronic B-cell aplasia and hypogammaglobulinemia due to “on-target, off-tumor” effects of CAR-T-cells³, which occurs when CAR-T-cells kill normal B-cells that express the CAR-T-cell target antigen. Although persistent detection of SARS-CoV-2 RNA from respiratory specimens of normal hosts is not thought to represent infectiousness^{4,5}, it is possible that CAR-T-cell recipients with B-cell aplasia can persistently shed infectious SARS-CoV-2. Similarly, SARS-CoV-2 RNAemia, which correlates with disease severity⁶, is thought to be uncommon and short-lived⁷ but may be more pronounced in CAR-T-cell recipients.

We present a case of prolonged SARS-CoV-2 infection in a patient who received CAR-T-cells targeting the B-cell maturation antigen (BCMA), which is part of the tumor necrosis factor superfamily and is involved with plasma cell survival and B-cell differentiation into plasma cells^{8,9}. The patient experienced persistent infection and SARS-CoV-2 RNAemia for more than two months and ultimately succumbed to the infection.

Case report:

The patient was a man in his 70's who underwent anti-BCMA CAR-T-cell therapy after lymphodepletion with fludarabine and cyclophosphamide. Pre-treatment nasopharyngeal (NP) swab reverse transcriptase polymerase chain reaction (RT-PCR) testing for SARS-CoV-2 (Cepheid Xpert Xpress¹⁰) was negative 17 and 2 days before CAR-T-cell therapy. He received tocilizumab for cytokine release syndrome, a known CAR-T-cell toxicity³, but did not require corticosteroids. He was discharged home then readmitted 12 days later with a two-day history of cough and shortness of breath. Temperature was 37.1°C, and oxygen saturation was 86% on room air. Laboratory evaluation showed a white blood cell (WBC) count of 3.5×10^9 cells/L.

(normal 3.8×10^9 - 10.6×10^9), with an absolute lymphocyte count (ALC) and absolute neutrophil count (ANC) of 0.7×10^9 cells/L (normal 0.8×10^9 - 3.65×10^9) and 2.6×10^9 cells/L (normal 2.24×10^9 - 7.68×10^9), respectively. NP RT-PCR testing was positive for SARS-CoV-2 RNA (day 0), with cycle thresholds of 20.1 and 21.5 for the nucleoprotein (N)1 and envelope (E) genes, respectively (**Figure 1**). Chest radiograph revealed bibasilar and mid-zone opacities. He was admitted to the intensive care unit (ICU) where he received five doses of remdesivir (days 5-10) and two units of convalescent plasma (day 2). He was discharged home in stable condition without supplemental oxygen on day 17. Asymptomatic SARS-CoV-2 NP RT-PCR testing remained positive on days 15, 26, and 37. Cycle thresholds were unavailable.

He was readmitted to the ICU with weakness and four days of dyspnea 41 days after his initial COVID-19 diagnosis. Temperature was 36.9°C , and oxygen saturation was 81% on room air. Laboratory evaluation showed a WBC count of 4.4×10^9 cells/L, ALC of 0.1×10^9 cells/L, and normal ANC. Lymphocyte subset testing showed 0% CD19+ B-cells. Chest computed tomography demonstrated diffuse bilateral ground glass opacities (**Figure 1**).

Given the radiographic findings and persistence of SARS-CoV-2 PCR NP swab positivity pre-admission, he received 10 days of dexamethasone and another course of convalescent plasma but was unable to receive additional remdesivir due to allocation issues at the time. He developed shock and ARDS, requiring vasopressor support then intubation on day 55. Bronchoalveolar lavage fluid from day 55 grew rare *Klebsiella pneumoniae*, which was not thought to be significantly contributing to his presentation but was treated with meropenem. A diagnostic workup for other infections, including blood and respiratory cultures, and non-culture diagnostics for fungi and viruses, was negative. SARS-CoV-2 NP RT-PCR testing remained positive on day 55, with cycle thresholds of 13.3 and 16 for the N1 and E genes, respectively. Serum from days 45, 56, and 71 was negative for SARS-CoV-2 IgG and IgA (Euroimmun assay^{11,12}). The family decided to focus on comfort measures because of non-resolving ARDS. The patient passed away on day 74 after COVID-19 diagnosis.

Methods:

The patient was enrolled in the University of Pittsburgh's Acute Lung Injury Registry and Biospecimen Repository (IRB# PRO10110387). Longitudinal characterizations of SARS-CoV-2 RNA and immune responses were performed on plasma and respiratory samples.

Virologic assays:

SARS-CoV-2 RNA quantification was performed using a sensitive (limit of detection 3 copies/reaction) quantitative RT-PCR (qRT-PCR) assay on plasma collected on days 4, 9, 13, 16, 67, and 71, and in endotracheal aspirate (ETA) fluid from day 72. Virus isolation from plasma (days 4, 65, and 71) and ETA fluid (day 72) was attempted on Vero E6 cells^{13,14}. To confirm that the isolated virus was SARS-CoV-2, indirect immunofluorescence (IIF) and sequencing of the spike (S) gene were performed. IIF was performed using anti-SARS CoV-2 S-antibodies, with phalloidin and DAPI counterstains for filamentous actin and cell nuclei, respectively. We performed a plaque assay¹³ to quantify viral titers in ETA samples for comparison with the qRT-PCR results.

Immune responses:

Total anti-SARS-CoV-2 IgG antibody titers directed against the S-protein receptor-binding domain (RBD) were determined on serially diluted plasma (days 4, 9, 13, 16, and 72) using an indirect enzyme-linked immunosorbent assay (ELISA)¹⁵. We focused on IgGs against the S-protein RBD because prior studies have suggested high antigenicity of the RBD¹⁶. We also performed a competitive ELISA assay with human ACE2 to assess neutralizing activity of anti-RBD IgG. Pseudovirus neutralization assays and plaque reduction neutralization assays on cultured SARS-CoV-2 were also performed^{13,14}. To characterize T-cell and B-cell responses, we performed single-cell RNAseq (sc-RNASeq) on peripheral blood mononuclear cells (PBMC) isolated from blood obtained on days 4 and 9.

Results:

SARS-CoV-2 RNA measurements:

High levels of SARS-CoV-2 RNA were detected in all plasma samples (**Figure 1**). RNAemia was greatest on day 4 (126,792 copies/mL) and subsequently showed a greater than 10-fold decrease by day 9 (8,100 copies/mL), after administration of convalescent plasma and remdesivir. RNAemia began to increase by days 13 (11,508 copies/mL) and 16 (18,000 copies/mL). Late in the course of infection, RNAemia remained relatively stable (14,720 copies/mL on day 67), but subsequently increased to 101,800 copies/mL by day 71 during administration of dexamethasone and before death. Over 20 billion copies of SARS-CoV-2 RNA (2.78×10^{10} copies/mL) were detected by qRT-PCR in an ETA sample from day 72.

Viral isolation and sequencing:

Viral isolation from plasma (days 4, 65, and 71) was unsuccessful despite the presence of SARS-CoV-2 RNA. Infectious SARS-CoV-2 was recovered from the day 72 ETA sample (**Figure 2**) with a titer of 1.125×10^6 plaque forming units (PFU)/mL. Numerous SARS-CoV-2 virions were detected by electron microscopy associated with respiratory epithelial cells in the ETA sample (**Figure 3**). Sanger sequencing of the S gene revealed three changes from the original Wuhan SARS-CoV-2 sequence: deletion of histidine at position 146 (H146)¹⁷, substitution of aspartic acid with glycine at position 614 (D614G), and a silent nucleotide change (V1094V). The S sequence was consistent with virus strains circulating in Pittsburgh and did not suggest superinfection or reinfection.

Immune responses:

Analysis of PBMCs by immunophenotyping and scRNA-Seq from days 4 and 9 demonstrated absence of B cells, consistent with anti-BCMA CAR-T-cell therapy. scRNA-Seq profiling was consistent with expansion of CAR-T-cells, and also showed evidence of interferon-stimulated genes in the monocyte lineage (e.g. *IFITM3*) consistent with a viral infection. No IgG antibody targeting the S-protein RBD was detected at any time point (**Supplementary table 1**). Additionally, no antibody that competed with human ACE2 binding to RBD was detected at any time point (**Supplementary table 1**). However, SARS-CoV-2 pseudovirus neutralization assays

demonstrated up to 50% inhibition of viral entry into cells at 1:10 dilution after day 13. SARS-CoV-2 plaque reduction neutralization assays also showed 50% neutralization at plasma dilutions of 1:4 to 1:16 on days 65 and 71 but not day 4, suggesting either the presence of low titers of neutralizing antibodies directed to non-RBD regions of S or detection of residual convalescent plasma (**Supplementary table 1**). Taken together, these findings demonstrate B-cell aplasia, low levels of neutralizing antibodies, and lack of anti-S (RBD) IgG.

Discussion

Here we report sustained SARS-CoV-2 RNAemia and active viral replication for greater than 2 months in a COVID-19 patient with B-cell aplasia related to anti-BCMA CAR-T-cells. The initial clinical improvement and reduction in plasma RNAemia observed between days 4 and 9 suggests a treatment response related to convalescent plasma and remdesivir, which he received on days 2 and 5-10, respectively. RNAemia was never completely suppressed, however, and we hypothesize that the absence of humoral responses to SARS-CoV-2 related to the expected effects of anti-BCMA CAR-T-cells resulted in uncontrolled viral replication after the initial hospitalization and overwhelming SARS-CoV-2 pneumonia (1.125×10^6 PFU/mL and 2.78×10^{10} RNA copies/mL in ETA) that likely contributed to death. These findings provide insights into SARS-CoV-2 replication and dissemination and the need to reassess infection control guidelines in immunocompromised patients with COVID-19.

RNAemia has been described in immunocompromised patients with respiratory viruses other than SARS-CoV-2. For instance, respiratory syncytial virus (RSV) RNAemia developed in 30% of hematopoietic cell transplant recipients 2 days after onset of RSV pneumonia and was a predictor of mortality¹⁸. Emerging data have shown that SARS-CoV-2 RNAemia may be a marker of disease severity in persons with COVID-19 pneumonia⁶, but a state of sustained RNAemia for >2 months has not yet been described. We were unable to isolate replication-competent virus from plasma, which may be due to technical issues, or may suggest that RNAemia is not caused by the presence of virions in the plasma, but rather due to “spillover” of

infected cells with SARS-CoV-2 RNA from the lung. Well-powered studies are needed to determine the prevalence of SARS-CoV-2 RNAemia, and to evaluate whether measurement of RNAemia should be adopted as a strategy to determine clinical response to SARS-CoV-2 therapies in clinical trials.

We successfully isolated virus from respiratory samples collected 72 days after onset of COVID-19, demonstrating the potential for infectivity late into the clinical course. In a prior study of patients with mild COVID-19, virus from respiratory samples was isolated only within the first week after symptom onset, with no virus isolated from samples obtained after 8 days⁵. Sequence analysis of isolated virus was consistent with circulating viruses and did not show evidence of superinfection. Importantly, our findings suggest that certain severely immunosuppressed patients with COVID-19 may require isolation longer than the 20-day period currently proposed by the Centers for Disease Control¹⁹. Whether cycle thresholds (which were consistently <25 in our patient) can be used to make decisions related to discontinuation of transmission-based precautions in such patients warrants further investigation.

Humoral immunity after SARS-CoV-2 infection in the general population appears to be sustained²⁰, but the role of antibodies in neutralization of virus and termination of infection is ill-defined. PBMC analysis confirmed B-cell aplasia, which is compatible with negative assay results for IgGs against the S-protein RBD, or IgAs and IgGs using a clinically approved assay. However, both viral and pseudoviral assays showed evidence of low-level neutralizing activity antibodies in the patient's serum. Based on the severe B-cell depletion, it is most likely that neutralization occurred because of receipt of convalescent plasma on day 58, although no neutralization was detected after the initial dose of convalescent plasma. Interestingly, scRNA-Seq profiling from days 4 and 9 showed upregulation of interferon-stimulated genes in monocytes, suggesting the presence of innate antiviral immunity, which was unable to control the infection. These findings suggest that effective B-cell responses may be required for viral clearance, although additional studies are needed.

In conclusion, our case highlights that immunocompromised patients with B-cell defects such as CAR-T-cell therapy recipients are at risk for prolonged and severe SARS-CoV-2 infections. Other patients with B-cell aplasia, such as recipients of anti-CD20 monoclonal antibodies, may also be at risk. Clinicians caring for such patients should be cautious not to attribute persistent detection of SARS-CoV-2 RNA from clinical samples of these patients to residual, non-infectious virus. Further work is needed to define the role of monitoring RNAemia in managing severely ill and immunocompromised persons with COVID-19, the duration of infectivity in these patients, and the immune phenotypes associated with recovery and fatal infection.

References

1. Wei J, Zhao J, Han M, Meng F, Zhou J. SARS-CoV-2 infection in immunocompromised patients: humoral versus cell-mediated immunity. *J Immunother Cancer* 2020;8(2).
2. Malard F, Genthon A, Brissot E, et al. COVID-19 outcomes in patients with hematologic disease. *Bone Marrow Transplant* 2020;
3. Haidar G, Garner W, Hill JA. Infections after anti-CD19 chimeric antigen receptor T-cell therapy for hematologic malignancies: timeline, prevention, and uncertainties. *Curr Opin Infect Dis* 2020;
4. Xiao AT, Tong YX, Zhang S. Profile of RT-PCR for SARS-CoV-2: a preliminary study from 56 COVID-19 patients. *Clin Infect Dis* 2020;
5. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020;581(7809):465–9.
6. Veyer D, Kernéis S, Poulet G, et al. Highly sensitive quantification of plasma SARS-CoV-2 RNA sheds light on its potential clinical value. *Clin Infect Dis* 2020;
7. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA* 2020;323(18):1843–4.
8. D'Agostino M, Raje N. Anti-BCMA CAR T-cell therapy in multiple myeloma: can we do better? *Leukemia* 2020;34(1):21–34.
9. Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-Cell Therapy bb2121 in Relapsed or Refractory Multiple Myeloma. *N Engl J Med* 2019;380(18):1726–37.
10. Xpert Xpress SARS-CoV-2 [Internet]. [cited 2020 Oct 26];Available from: <https://www.fda.gov/media/136314/download>
11. Van Elslande J, Houben E, Depypere M, et al. Diagnostic performance of seven rapid IgG/IgM antibody tests and the Euroimmun IgA/IgG ELISA in COVID-19 patients. *Clin Microbiol Infect* 2020;26(8):1082–7.

12. Anti-SARS-CoV-2 ELISA (IgG) [Internet]. [cited 2020 Oct 27];Available from:
<https://www.fda.gov/media/137609/download>
13. Hartman AL, Nambulli S, McMillen CM, et al. SARS-CoV-2 infection of African green monkeys results in mild respiratory disease discernible by PET/CT imaging and shedding of infectious virus from both respiratory and gastrointestinal tracts. *PLoS Pathog* 2020;16(9):e1008903.
14. Klimstra WB, Tilston-Lunel NL, Nambulli S, et al. SARS-CoV-2 growth, furin-cleavage-site adaptation and neutralization using serum from acutely infected hospitalized COVID-19 patients. *J Gen Virol* 2020;
15. Liu X, Drelich A, Li W, et al. Enhanced elicitation of potent neutralizing antibodies by the SARS-CoV-2 spike receptor binding domain Fc fusion protein in mice. *Vaccine* 2020;38(46):7205–12.
16. Yang J, Wang W, Chen Z, et al. A vaccine targeting the RBD of the S protein of SARS-CoV-2 induces protective immunity. *Nature* 2020;
17. Singh PK, Kulsum U, Rufai SB, Mudliar SR, Singh S. Mutations in SARS-CoV-2 Leading to Antigenic Variations in Spike Protein: A Challenge in Vaccine Development. *J Lab Physicians* 2020;12(2):154–60.
18. Waghmare A, Campbell AP, Xie H, et al. Respiratory syncytial virus lower respiratory disease in hematopoietic cell transplant recipients: viral RNA detection in blood, antiviral treatment, and clinical outcomes. *Clin Infect Dis* 2013;57(12):1731–41.
19. Centers for Disease Control. Duration of Isolation and Precautions for Adults with COVID-19 [Internet]. [cited 2020 Oct 16]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/duration-isolation.html>
20. Gudbjartsson DF, Norddahl GL, Melsted P, et al. Humoral Immune Response to SARS-CoV-2 in Iceland. *N Engl J Med* 2020;

Figure 1. Clinical timeline showing CAR-T infusion (day -25 to 0), first hospital admission (days 0-17), home stay (days 18-41), and second hospital admission (days 41-74). Day 0 denotes day of first positive nasopharyngeal (NP) swab PCR for SARS-CoV-2 RNA. The top panel shows the clinical course and timeline, with results of clinical SARS-CoV-2 NP swab RT-PCR testing and clinical IgG and IgA testing (Euroimmun assay). The bottom graph shows serial plasma SARS-CoV-2 RNA quantification obtained using a research qRT-PCR assay, with viral quantification from endotracheal aspirate fluid using research assays (qRT-PCR and viral titer). N1 and E are the SARS-CoV-2 genes detected by the clinical nasopharyngeal swab assay used. Inset shows CT (computed tomography) from day 41. Green squares: convalescent plasma (CP); red squares: remdesivir; light blue squares: dexamethasone.

A. 20x

B. 40x

Figure 2. Indirect immunofluorescence of SARS-CoV-2 from day 72 respiratory (endotracheal aspirate) sample isolated and grown in Vero E6 cells. Green: antibody directed against SARS CoV-2 spike protein (Sinobiologicals). Blue: DNA counterstained with DAPI. Red: filamentous actin counterstained with phalloidin. A syncytium is shown in the inset (B) at higher magnification. Scale bars: 100 μm.

Figure 3. Electron microscopy of respiratory (endotracheal aspirate) sample obtained on day 72. (A) Comparison of virus localization in a bronchial epithelial cell (top) and a transient secretory cell (bottom). Each image is shown as montaged 2D overviews of the whole cell in a 200 nm section (left), overlaid with colored dots to indicate positions of virions (center) and 3D tomographic reconstructions detailing virus populations within cytoplasmic compartments. (B) 2D overview of SARS-CoV-2–infected ciliated epithelial cell in a 200 nm semi-thick section. The apical side of the cell is characterized by numerous membrane-bound compartments that are filled with SARS-CoV-2 virions. (C) Montaged tomogram of the apical side of the cell shown in B; hundreds of virus particles are contained within smooth-walled cytoplasmic compartments. Inset in C: Detail of a single virion with spikes indicated by red dots. (D) Tomogram detail of a smooth-walled cytoplasmic compartment containing at least 30 SARS-CoV-2 virions within the shown 15 nm thick volume.