Acyclic subgraph is hard

Thinh D. Nguyen
Moscow State University
kosmofarmer@gmail.com

Abstract. We prove the hardness of yet another problem in graph theory, namely ACYCLIC SUBGRAPH. A reduction from INDEPENDENT SET shows that our claim holds.

Keywords: acyclic, edge, graph, independent, vertex

1. Introduction

In this section, we give the description of the problem and necessary notions from graph theory.

1.1. GRAPH THEORY NOTIONS

A graph $G = (V, E)$ is an undirected graph with the *vertex set* $V = V(G)$ and *edge set* $E = E(G)$. In a graph, every edge (u, v) ($u, v \in V$) is undirected. Digraph is a different notion, a digraph $D = (V, A)$ is a directed graph with the vertex set $V = V(D)$ and *arc set* $A = A(D)$. In a digraph, every arc (u, v) ($u, v \in V$) is directed, meaning that u is the starting vertex of the arc, and v is the ending vertex of the arc. A *cycle* in a digraph D is a sequence of vertices $(v_1, v_2, \dots, v_k)$ in which, for all $1 \leq i < k$, (v_i, v_{i+1}) is an arc of D , and also (v_k, v_1) is an arc of D . A digraph without any cycle is called a *DAG* (Directed Acyclic Graph). In an (undirected) graph, two vertices u, v are adjacent if (u, v) is an edge.

In a graph G , a set $V' \subset V$ is an independent set if V' has no pair of adjacent vertices. A well-known result states that the problem of finding the largest independent set INDEPENDENT SET is hard.

Definition 1.1. INDEPENDENT SET = $\{ \langle G, k \rangle \mid \text{the graph } G \text{ has an independent set } V' \subset V(G) \text{ containing } k \text{ vertices} \}$

1.2. ACYCLIC SUBGRAPH

Now, we are going to give the definitions that are pertinent to our soon-to-be-defined problem. The notion of induced subgraph of an undirected graph is well studied. But, in this article, we consider the notion of induced subgraph of a digraph. The concept is similar. An induced subgraph of a digraph $D = (V, A)$ is also determined by its vertex set $V' \subset V$. Its arc set is all the arc in $V' \times V' \cap A(D)$. Our problem is to find the largest DAG subgraph of a given digraph.

Definition 1.2. ACYCLIC SUBGRAPH = $\{ \langle D, k \rangle \mid \text{the digraph } D \text{ has an induced DAG subgraph on } V' \subset V(D) \text{ containing } k \text{ vertices, in other words, } D[V'] \text{ is a DAG and } |V'| = k \}$

We also utilize the notation of the induced subgraph by $V' \subset V$ in a digraph $D = (V, A)$ by $D[V']$.

2. Main claim and proof

Claim 2.1. INDEPENDENT SET $\leq_p$ ACYCLIC SUBGRAPH

Proof. For an instance of INDEPENDENT SET $G = (V, E)$ and an integer k , create a digraph $D = (V, A)$ as follows. The vertex set V is the same as in G . For each edge $(u, v) \in E(G)$, we create two arcs (u, v) and (v, u) in $A(D)$. Clearly, if G has an independent set V' of size k , the induced subgraph of V' in D is necessarily a DAG. Since a cycle in $D[V']$ would have some arc, and an arc in D corresponds to an edge in G . Conversely, if D has a induced DAG subgraph $D[V']$ with $|V'| = k$ then V' must form an independent set in G . Assume to the contrary that $|V'|$ contains an edge (u, v) in G , that edge necessarily corresponds to two arcs (u, v) and (v, u) in D . These two arcs form a cycle, namely (u, v) in D . $\square$

References

- [1] Michael R. Garey, David S. Johnson, **Computers and Intractability: A Guide to the Theory of *NP*-completeness**
- [2] The free encyclopedia Wikipedia, *<https://en.wikipedia.org/wiki/NP-completeness>*