

## Digital detox: The effect of smartphone abstinence on mood, anxiety, and craving

Wilcockson T.D.W.<sup>a,b,\*</sup>, Osborne A.M.<sup>c</sup>, Ellis D.A.<sup>b</sup>

<sup>a</sup> *Loughborough University, Loughborough, UK*

<sup>b</sup> *Lancaster University, Lancaster, UK*

<sup>c</sup> *Duke-NUS Medical School, Singapore*

\* Corresponding author at: Loughborough University, Loughborough, UK. *E-mail address:* [t.wilcockson@lboro.ac.uk](mailto:t.wilcockson@lboro.ac.uk) (T.D.W. Wilcockson).

Cite as: Wilcockson, T. D., Osborne, A. M. and Ellis, D. A. (2019). Digital detox: The effect of smartphone withdrawal on mood, anxiety, and craving. *Addictive Behaviors*, 99, 106013.

### HIGHLIGHTS

- *Investigates the impact of smartphone abstinence on mood, anxiety, and craving.*
- *Participants had to abstain from using their smartphone for 24 hours.*
- *Levels of craving increased following smartphone abstinence.*
- *Mood and anxiety were unaffected following smartphone abstinence.*

### ABSTRACT

Whether behavioural addictions should be conceptualised using a similar framework to substance-related addictions remains a topic of considerable debate. Previous literature has developed criteria, which allows any new behavioural addiction to be considered analogous to substance-related addictions. These imply that abstinence from a related object (e.g., smartphones for heavy smartphone users) would lead to mood fluctuations alongside increased levels of anxiety and craving. In a sample of smartphone users, we measured three variables (mood, anxiety, and craving) on four occasions, which included a 24-hour period of smartphone abstinence. Only craving was affected following a short period of abstinence. The results suggest that heavy smartphone usage does not fulfil the criteria required to be considered an addiction. This may have implications for other behavioural addictions.

## 1. Introduction

Behavioural addiction is defined by the DSM-V as an addictive disorder that does not involve the ingestion of a psychoactive substance (APA, 2013). However, Kardefelt-Winther et al. (2017) have recently argued that research concerning behavioural addictions has not yet clarified whether sufferers become functionally impaired, experience psychological distress, or demonstrate any separation from normative behaviour. Failure to meet these criteria may indicate that an addiction is not present. Symptoms associated with substance addiction include mood modification, tolerance, and withdrawal (Griffiths, 2005). Therefore, we would also expect to see these symptoms in behavioural addictions, however, their measurement is often problematic. For example, how would one quantify tolerance within internet addiction? Further, with behavioural addictions in digital domains, it is difficult to appreciate where a line might be drawn between typical, excessive and problematic usage (see Ellis et al., 2019). Problematic usage should impair normal functioning and cause distress. For example, abstinence from addiction-related behaviours (e.g., drinking for heavy drinkers), leads to changes in mood, anxiety, and craving (cf. Kardefelt-Winther et al., 2017). If abstinence results in changes across all three measures, then this might reveal analogous symptoms necessary for a new phenomenon to be considered a genuine behavioural addiction.

In recent years, a growing body of research has focused on the potential problems associated with excessive smartphone use (e.g., Pan et al., 2019; Kim et al., 2019; Lee et al., 2019; Wilcockson, Ellis, & Shaw, 2018). However, Billieux, Maurage, Lopez-Fernandez, Kuss, and Griffiths (2015) argue that very little evidence supports the notion that smartphone use can be considered a form of behavioural addiction. Related research has focused specifically on social media. For example, Stieger and Lewetz (2018) observed that social media abstinence led to an increase in craving for social media, but anxiety and mood were unaffected. Another study by Vanman, Baker, and Tobin (2018) however, observed that people who gave-up Facebook reported lower levels of wellbeing. In contrast, comparatively little research has considered the psychological changes that occur as people experience smartphone abstinence, which are primarily used to access to these services. Such research could support or refute the current literature base concerning the potential psychological consequences of smartphone addiction. Previously, Clayton, Leshner, and Almond (2015) reported that smartphone separation led to negative affect if a participant was prevented from answering their phone while it rang in another room. But this separation anxiety may not necessarily reflect addiction-like anxiety, which would be the result of prolonged functional impairment and distress and not simply event-based (cf. Kardefelt-Winther et al., 2017). To date, no study has examined smartphone abstinence over a 24-hour period. The aim of this project is therefore to **consider how mood, anxiety, and craving change when participants stop using their smartphone for 24-hours.**

Participants attended the lab on four occasions and completed a battery of tasks. The first session took place a week before the abstinence task, with the second session occurring immediately before abstinence. A third session took place immediately after a 24-hour smartphone abstinence, with the final session taking place the following day. We expected no differences between responses on sessions 1 and 4, however, we predicted that changes would likely occur immediately before and after the abstinence task (sessions 2 and 3). Specifically, before the abstinence task people may be concerned about giving-up their device for 24 h. Conversely, people are likely to be relieved after any period of abstinence is over.

## 2. Method

### 2.1. Participants

There were 45 participants who started the study (33% male; average age = 22.4), however, nine participants did not complete all four lab sessions (see Fig. 1). Participants were recruited from the Psychology subject-pool at Lancaster University and by advertising the study across campus using posters. Recruitment was blind to any current levels of smartphone usage however, previous research demonstrates that younger participants spend more time on their smartphone than older adults (Christensen et al., 2016; Ellis et al., 2019). They were reimbursed £15 for their time. Full ethical approval was obtained prior to the study and all participants provided written informed consent.


## 2.2. Materials

We used a number of paper-based measures to assess anxiety (Marteau & Bekker, 1992), mood (Mayer & Gaschke, 1988), craving for smartphones (modified desire for drinking questionnaire: Love, James, & Willner, 1998), and smartphone addiction (Chen et al., 2014). The STAI-6 (State-trait Anxiety Inventory) is a 6-item measure where participants can respond to each statement using a 4-point Likert scale e.g., “I feel calm”. Responses can range from “not at all” to “very much”. The Brief Mood Introspection Scale (BMIS) consists of two parts, [1] a 16-item questionnaire (e.g., happy, lively, sad) with a 4-point Likert response scale ranging from “definitely do not feel” to “definitely feel” and [2] an ‘overall mood’ question where participants indicate their current mood on a 21-point scale ranging from “very unpleasant” to “very pleasant”. To assess craving, we used a modified version of the Love et al. (1998) Desire for Alcohol Questionnaire with smartphone terminology replacing alcohol terminology. This is a 37-item questionnaire (e.g., “I could easily limit how much I use my phone”) with a 7-point Likert response scale ranging from “strongly agree” to “strongly disagree”. Finally, the Smartphone Addiction Inventory (SPAI) is a 26-item questionnaire (e.g., “I feel restless and irritable when my smartphone is unavailable”) with 4 responses ranging from “disagree” to “agree”. Cronbach’s alpha were  $> 0.75$  for all measures.

## 3. Procedure

In the first lab session, participants completed all questionnaires. They then returned to the lab one week later and had their phone placed in an evidence bag, which they were requested to not open/use. Selected questionnaires were also administered: mood, craving, and anxiety. 24 h after the abstinence task began, participants returned to the lab and completed the selected questionnaires again (session 3). After the abstinence task was completed, participants were asked to return to the lab a fourth and final time to complete the selected questionnaires.


During the abstinence period, participants were instructed to place their smartphone in a secure evidence bag. In case of an emergency or if they wished to withdraw from the study, it was possible to quickly tear the bag open and use their phone at any time. Note that no participants returned to the lab with opened or tampered evidence bags.



**Fig. 1.** Flow diagram of procedure and participation discontinuation at each stage. Session 1 occurred a week prior to the abstinence task, session 2 occurred immediately before abstinence. The Abstinence Task lasted 24 h with session 3 taking place immediately after. Session 4 occurred a further 24 h later. This diagram also reports average Smartphone Addiction Inventory (aSPAI) scores for participants who left the study. **Note, the average SPAI score for participants who discontinued at session 3 (67) was higher than the average SPAI score derived from all participants at the start of the study (55).** See supplementary materials for differences between participants who completed or discontinued based on session 1 scores (Table S1).

#### 4. Results

A number of measures were taken at different time periods. Therefore, for each measure, we initially calculated a repeated-measures ANOVA with 4 levels (session: 1, 2, 3, 4). If appropriate, comparisons were then conducted between different sessions. Additionally, a Bayes factor with default prior scales is computed for each analysis (Love et al., 2015; Morey & Rouder, 2014; Rouder, Morey, Speckman, & Province, 2012). Computing a Bayes factor provides us with the ability to interpret p-values > .05. Therefore, a  $BF_{10} < 0.33$  indicates some evidence for the null hypothesis.  $BF_{10} > 3$  provides strong evidence for the alternate hypothesis (e.g., Rouder et al., 2012).



**Fig. 2.** Average scores across sessions for [A] mood, [B] anxiety, and [C] craving. Error bars represent standard error of the mean. Individual responses for participants who completed all four lab sessions appear below. These illustrate no systematic changes in mood or anxiety, but consistent changes in craving for most participants. Note that a 24-hour period of smartphone abstinence occurred between sessions 2 and 3.

#### 4.1. Mood and anxiety

Overall, Fig. 2 suggests that mood was lower immediately before the abstinence task, but gradually increased **towards** the end of the study. A small reduction in anxiety is also apparent during the final session. However, ANOVAs did not reveal a significant main effect of session on mood [ $F(3,105) = 1.79$ ;  $p = .15$ ;  $BF_{10} = 0.29$ ] or anxiety [ $F(3,105) = 1.08$ ;  $p = .36$ ;  $BF_{10} = 0.13$ ].

#### 4.2. Craving

A significant main effect of session was observed on craving [ $F(3,105) = 14.15$ ;  $p < .0005$ ;  $BF_{10} > 100$ ]. Uncorrected comparisons revealed that most sessions differed significantly from each other: session 1 [ $M = 93.95$ ;  $SD = 45.58$ ] and session 4 [ $M = 66.86$ ;  $SD = 37.06$ ;  $t(35) = 4.67$ ;  $p < .0005$ ;  $BF_{10} > 100$ ]; session 2 [ $M = 84.80$ ;  $SD = 45.22$ ] and session 3 [ $M = 98.78$ ;  $SD = 44.21$ ;  $t(40) = 3.09$ ;  $p = .004$ ;  $BF_{10} = 9.68$ ]; session 2 and session 4 [ $t(35) = 3.93$ ;  $p < .0005$ ;  $BF_{10} = 74.19$ ]; session 3 and session 4 [ $t(35) = 8.16$ ;  $p < .0005$ ;  $BF_{10} > 100$ ]. However no significant differences were observed between session 1 and session 2 [ $t(40) = 1.97$ ;  $p = .06$ ;  $BF_{10} > 0.97$ ] or session 1 and session 3 [ $t(40) = 0.91$ ;  $p = .38$ ;  $BF_{10} > 0.25$ ].

#### 4.3. Problematic smartphone usage

During the first session, participants completed the problematic phone usage questionnaire (SPAI). We observed that this measure positively correlated with craving measures taken during session 1 [ $r(43) = 0.69$ ;  $p < .0005$ ;  $BF_{10} > 100$ ], session 2 [ $r(39) = 0.60$ ;  $p < .0005$ ;  $BF_{10} > 100$ ], session 3 [ $r(39) = 0.62$ ;  $p < .001$ ;  $BF_{10} > 100$ ], and session 4 [ $r(34) = 0.66$ ;  $p < .0005$ ;  $BF_{10} > 100$ ]. Therefore, while levels of craving varied between each session, it would appear that participants who believed they used their smartphone more also reported higher levels of craving. Mood and anxiety scores were not associated with the SPAI at any time point [all  $p$ 's  $> 0.1$ ].

## 5. Discussion

Whether or not behavioural addictions are akin to substance addictions remains a matter of considerable debate (Kardefelt-Winther et al., 2017). However, our results suggest that while smartphone abstinence can lead to craving, mood and anxiety remain unaffected. **We also note that while craving did increase during the abstinence period (between sessions 2 and 3), it remained similar to levels reported a week prior to taking part (session 1). Nevertheless, while these results may indicate** that smartphone users like to use their smartphones and crave them when they are unavailable, the lack of evidence for mood modification and increased anxiety suggests a key distinction between technology-related behaviours and substance abusers. Substance abusers during abstinence would demonstrate mood modification and increased anxiety. Therefore, this distinction suggests that behavioural addictions (e.g., technology usage) are unlikely to inhabit the same underlying processes as substance-related addictions (e.g., alcohol usage). This distinction is important from an addictions perspective as substance abusers continue to take substances in the absence of liking (see Robinson & Berridge, 1993). While liking is not necessarily the strongest motivator in substance abuse addiction, it may be the strongest driver in any technology-related behavioural addiction.

Although there was no significant effect of abstinence on mood, we note that some improvement in mood does occur between sessions 2 and 4. This suggests that once participants were reunited with their phone following abstinence, they reported improved mood compared to immediately before the abstinence period (session 2). While being reunited with their phone may have made people feel happier, this difference may also be the result of poorer mood when pre-empting abstinence. These factors combined could also magnify this effect. However, while this may provide some evidence to confirm mood modification, our Bayes result suggests that more evidence is required to support any **changes to mood** before or after any period of smartphone abstinence.

This study involved restricting the use of smartphones, but not all technology (e.g., laptops) completely. Our findings are therefore limited by the possibility that participants may have been using other digital devices. This may explain why anxiety and mood were not affected, but changes in craving scores contradict this interpretation somewhat. It would have also been ethically difficult to restrict all technology use completely. Further, our sample may not harbour problematic smartphone usage and have therefore not responded accordingly. However, as problematic smartphone usage scores increased so did craving. Problematic smartphone users in the current study may have simply discontinued (see Fig. 1). It is striking that **drop-outs at session 3 had slightly higher SPAI scores**. This may indicate that smartphone ‘addicts’ were unable to fully participate in the study and so discontinued, thus affecting our findings. However, we would caution this interpretation somewhat as these scales do not align favourably with objective behaviour (Ellis, 2019). Future research could focus on heavy users, based on objective behaviour, who may be more likely to demonstrate expected patterns of withdrawal.

In summary, our data suggest that normal emotional functioning is not impaired by smartphone abstinence, which is outlined as a key symptom of any addiction (see Kardefelt-Winther et al., 2017; Robinson & Berridge, 1993). Therefore, heavy smartphone usage may not meet the criteria for a behavioural-type addiction. It does appear that smartphone abstinence is associated with craving, but this alone does not necessarily reflect any form of addiction.

## Acknowledgements

This work was funded by a Faculty of Science and Technology research grant (PSA7866) from Lancaster University. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## References

American Psychiatric Association (APA) (2013). *Diagnostic and Statistical Manual of Mental Disorders, 5th edn*. Arlington, VA: Washington, DC: American Psychological Association Press.

Billieux, J., Maurage, P., Lopez-Fernandez, O., Kuss, D. J., & Griffiths, M. D. (2015). Can disordered mobile phone use be considered a behavioral addiction? An update on current evidence and a comprehensive model for future research. *Current Addiction Reports*, 2(2), 156–162.

Chen, S. H. (2014). Development and validation of the smartphone addiction inventory (SPAI). *PLoS One*, 9(6), e98312.

Christensen, M. A., Bettencourt, L., Kaye, L., Moturu, S. T., Nguyen, K. T., Olgin, J. E., ... Marcus, G. M. (2016). Direct measurements of smartphone screen-time: Relationships with demographics and sleep. *PLoS One*, 11(11), e0165331.

Clayton, R. B., Leshner, G., & Almond, A. (2015). The extended iSelf: The impact of iPhone separation on cognition, emotion, and physiology. *Journal of Computer-Mediated Communication*, 20(2), 119–135.

Ellis, D. A. (2019). Are smartphones really that bad? Improving the psychological measurement of technology-related behaviors. *Computers in Human Behavior*, 97, 60–66.

Ellis, D. A., Davidson, B. I., Shaw, H., & Geyer, K. (2019). Do smartphone usage scales predict behavior? *International Journal of Human-Computer Studies*, 130, 86–92.

Griffiths, M. (2005). A 'components' model of addiction within a biopsychosocial framework. *Journal of Substance Use*, 10(4), 191–197.

Kardefelt-Winther, D., Heeren, A., Schimmenti, A., van Rooij, A., Maurage, P., Carras, M., ... Billieux, J. (2017). How can we conceptualize behavioural addiction without pathologizing common behaviours? *Addiction*, 112(10), 1709–1715.

Kim, S. G., Park, J., Kim, H. T., Pan, Z., Lee, Y., & McIntyre, R. S. (2019). The relationship between smartphone addiction and symptoms of depression, anxiety, and attention-deficit/hyperactivity in South Korean adolescents. *Annals of General Psychiatry*, 18(1), 1.

Lee, S. J., Choi, M. J., Rho, M. J., Kim, D. J., & Choi, I. Y. (2019). Factors Affecting User Acceptance in Overuse of Smartphones in Mobile Health Services: An Empirical Study Testing a Modified Integrated Model in South Korea. *Frontiers in Psychiatry*, 10, 259.

Love, A., James, D., & Willner, P. (1998). A comparison of two alcohol craving questionnaires. *Addiction*, 93(7), 1091–1102.

Love, J., Selker, R., Marsman, M., Jamil, T., Dropmann, D., Verhagen, A. J., & Wagenmakers, E. J. (2015). *JASP (version 0.7) [computer software]*. Amsterdam, the netherlands: Jasp project.

Marteau, T. M., & Bekker, H. (1992). The development of a six-item short-form of the state scale of the Spielberger state—Trait anxiety inventory (STAI). *British Journal of Clinical Psychology*, 31(3), 301–306.

Mayer, J. D., & Gaschke, Y. N. (1988). The experience and meta-experience of mood. *Journal of Personality and Social Psychology*, 55(1), 102–111.

Morey, R. D., & Rouder, J. N. (2014). *BayesFactor (version 0.9. 9)*.

Pan, Y. C., Lin, H. H., Chiu, Y. C., Lin, S. H., & Lin, Y. H. (2019). Temporal Stability of Smartphone Use Data: Determining Fundamental Time Unit and Independent Cycle. *JMIR mHealth and uHealth*, 7(3).

Robinson, T. E., & Berridge, K. C. (1993). The neural basis of craving: An incentive-sensitization theory of addiction. *Brain Research Reviews*, 18, 247–291.

Rouder, J. N., Morey, R. D., Speckman, P. L., & Province, J. M. (2012). Default Bayes factors for ANOVA designs. *Journal of Mathematical Psychology*, 56(5), 356–374.

Stieger, S., & Lewetz, D. (2018). A week without using social media: Results from an ecological momentary intervention study using smartphones. *Cyberpsychology, Behavior, and Social Networking*, 21(10), 618–624.

Vanman, E. J., Baker, R., & Tobin, S. J. (2018). The burden of online friends: the effects of giving up Facebook on stress and well-being. *The Journal of Social Psychology, 158*(4), 496–507.

Wilcockson, T. D., Ellis, D. A., & Shaw, H. (2018). Determining typical smartphone usage: What data do we need? *Cyberpsychology, Behavior and Social Networking, 21*(6), 395–398.



Table S1. Mean and (SD) responses from participants who completed or discontinued. Each score is derived from Session 1 measurements.

	Remained in study	Discontinued participation
Mood	6.6 (1.5)	7.6 (1.2)
Craving	27.81 (16.1)	33.8 (16.2)
Anxiety	6.9 (2.7)	7.2 (1.9)