NEUROD4 is differentially expressed in the tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab (Herceptin) is a monoclonal antibody targeting the extracellular domain of the human epidermal growth factor receptor 2 (HER2)¹ utilized for the treatment of adjuvant and metastatic breast cancer² in the United States and worldwide. We mined published microarray and gene expression data^{3,4} to discover in an unbiased manner the most significant transcriptional changes associated with trastuzumab treatment. We identified NEUROD4 among the genes most differentially expressed in the primary tumors of patients with breast cancer treated with trastuzumab. The primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of NEUROD4 messenger RNA than did patients not treated with trastuzumab, and a single administration of trastuzumab was sufficient to result in differential expression of NEUROD4 in primary tumors of the breast, demonstrating increased expression of a transcription factor with neurogenic potential^{5,6} as a direct transcriptional consequence of treatment with trastuzumab.

Keywords: trastuzumab, breast cancer, NEUROD4, systems biology of breast cancer, targeted therapeutics in breast cancer.

The mechanism of action ascribed to trastuzumab function is thought to be binding of the F_{ab} domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2 (HER2)¹, but the transcriptional consequences resulting from signaling downstream of trastuzumab binding to HER2 is less well understood. Through unbiased, systems-level comparative transcriptome analysis of published microarray and gene expression data^{3,4}, we report here that NEUROD4 is a differentially expressed gene in the tumors of trastuzumab-treated patients: patients treated with trastuzumab expressed significantly higher levels of NEUROD4 than did those of patients not treated with trastuzumab.

Methods

We used datasets GSE130788³, and GSE114082⁴ for this differential gene expression analysis of the transcriptional response to trastuzumab treatment in patients with breast cancer. GSE130788 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with $n=11$ primary breast tumors from patients treated with trastuzumab and $n=36$ primary tumors from patients treated with trastuzumab, analysis performed using platform GPL6480. GSE114082 was generated using Illumina HumanHT-12 WG-DASL V4.0 R2 expression beadchip technology with $n=17$ primary breast tumors from patients before trastuzumab treatment and $n=17$ primary tumors from patients after trastuzumab treatment, analysis performed using platform GPL14951. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global or multiplexed differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether NEUROD4 gene expression was significantly between primary breast tumors of patients receiving trastuzumab using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

Results

We performed unbiased comparative transcriptome analysis using published microarray and gene expression data^{3,4} to identify the most significant transcriptional features of the primary breast tumors of patients treated with trastuzumab (Herceptin).

NEUROD4 is differentially expressed in the tumors of patients treated with trastuzumab.

We identified neuronal differentiation 4, encoded by NEUROD4 as among the genes whose expression changed most significantly in the tumors of breast cancer patients treated with trastuzumab in a multiplexed gene expression dataset³ (Table 1). When sorting each of the genes expressed in the primary tumors of breast cancer patients treated with trastuzumab based on significance of change as compared to that of patients not treated with trastuzumab, NEUROD4 ranked 70 out of 41000 total transcripts (Table 1), equating to 99.8% differential expression. NEUROD4 differential expression in the tumors of patients treated with trastuzumab trended towards statistical significance (Table 1; $p=3.89E-63$).

We turned to a second microarray dataset to validate our findings of NEUROD4 differential expression in the patients treated with trastuzumab. The tumors compared in this dataset were sampled from the same patient before and after a single, brief exposure to trastuzumab, not comparing tumors from different patients treated with a course of trastuzumab or not. Thus, while differentially expressed genes when comparing tumors based on trastuzumab treatment can result from a variety of sources, including gene expression changes resulting from tumor evolution over the course of treatment, changes identified

1 in this dataset, occurring after only a single treatment of trastuzumab, are more likely to be a direct
2 transcriptional result of trastuzumab binding events. We found again that NEUROD4 was among the
3 genes most differentially expressed in the tumors of patients administered a single treatment of
4 trastuzumab (Table 2). When sorting each of the genes expressed in the tumors of patients administered a
5 single treatment of trastuzumab, NEUROD4 ranked 1197 out of 29377 total transcripts (Table 2),
equating to 95.9% differential expression. Differential expression of NEUROD4 in the primary tumors of
breast cancer patients following treatment with trastuzumab was statistically significant (Table 2;
 $p=1.13E-04$).

**NEUROD4 is expressed at significantly higher levels in the tumors of breast cancer patients treated
with trastuzumab than in the tumors of those not treated with trastuzumab.**

8 We obtained exact mRNA expression levels for NEUROD4 from tumors of patients treated with
9 trastuzumab and from the tumors of patients not treated with trastuzumab to directly evaluate direction
10 and statistical significance of change in expression of NEUROD4 based on history of trastuzumab
11 treatment. NEUROD4 was expressed at higher levels in the tumors of patients treated with trastuzumab
as compared to those of patients not treated with trastuzumab, and this difference was deemed statistically
significant (Figure 1; $p<0.0001$).

12 Thus, we found using published multiplexed gene expression and microarray data^{3,4} that
13 NEUROD4 was among the genes most differentially expressed in the tumors of patients treated with
14 trastuzumab, that NEUROD4 was expressed at higher levels in the tumors of patients treated with
trastuzumab as compared to those not treated with trastuzumab, and that this transcriptional change was
likely a direct result of trastuzumab treatment as opposed to a gene expression change associated with
clonal selection and evolution of the tumor across the course of treatment.

Discussion

16 We provided evidence here that NEUROD4 is a differentially expressed gene in the tumors of
17 patients treated with trastuzumab, and expressed at significantly higher levels in the primary tumors of
18 breast cancer patients treated with trastuzumab as compared to patients not treated with trastuzumab.
19 Thus, trastuzumab, a treatment associated with a 34% rate of central nervous system metastases⁷,
administration in patients with breast cancer results in increased primary tumor expression of a basic
helix-loop-helix transcription factor with neurogenic potential^{5,6}.

References

1. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
2. Herceptin (Trastuzumab). Indications. <https://www.herceptin.com/hcp/indications.html>.
3. GSE130788. A Phase II Randomized Trial of Neoadjuvant Trastuzumab or Lapatinib or the Combination of Trastuzumab and Lapatinib, Followed by Docetaxel and Carboplatin with Trastuzumab and/or Lapatinib in Patients with HER2+ Breast Cancer. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788>. Hsiaowang Chen. UCLA, Department of Medicine. Santa Monica, CA.
4. Triulzi, T., Regondi, V., De Cecco, L., Cappelletti, M.R., Di Modica, M., Paolini, B., Lollini, P.L., Di Cosimo, S., Sfondrini, L., Generali, D. and Tagliabue, E., 2018. Early immune modulation by single-agent trastuzumab as a marker of trastuzumab benefit. *British journal of cancer*, 119(12), pp. 1487-1494.
5. Buchman, V.L., Ninkina, N.N., Bogdanov, Y.D., Bortvin, A.L., Akopian, H.N., Kiselev, S.L., Krylova, O.Y., Anokhin, K.V. and Georgiev, G.P., 1992. Differential splicing creates a diversity of transcripts from a neurospecific developmentally regulated gene encoding a protein with new zinc-finger motifs. *Nucleic acids research*, 20(21), pp.5579-5585.
6. Hardwick, L.J. and Philpott, A., 2015. Multi-site phosphorylation regulates NeuroD4 activity during primary neurogenesis: a conserved mechanism amongst proneural proteins. *Neural development*, 10(1), pp.1-18.
7. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.

Rank	ID	p-value	t	B	Gene	Gene name
70	A_32_P320247	3.89E-63	24.909751	133.319395	NEUROD4	neuronal differentiation 4

Table 1: NEUROD4 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene
1197	ILMN_1677608	1.13E-04	4.3349581	1.14905	NEUROD4

Table 2: A single exposure to trastuzumab is sufficient to drive differential expression of NEUROD4 in the primary tumors of patients with breast cancer.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: NEUROD4 is expressed at significantly higher levels in the primary tumors of patients treated with trastuzumab.

The mRNA expression level of NEUROD4 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between NEUROD4 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.