The BRD8 bromodomain protein is differentially expressed in brain metastatic breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a serious complication of breast cancer¹⁻³ with limited treatment options^{4,5}. We mined published microarray data^{6,7} to discover genes associated with brain metastasis in human breast cancer, and identified differential expression of the gene encoding the bromodomain protein BRD8 when comparing brain metastatic tissues to primary tumors of the breast. BRD8 was expressed at significantly lower levels in metastasis to the brain as compared to primary tumors, and primary tumor expression of BRD8 was significantly correlated with relapse-free survival. Understanding the transcriptional functions of BRD8 in the context of the Tip60/NuA4 histone acetyltransferase complex in human breast cancer may shed light on mechanisms that contribute to migration of tumor cells from the breast and subsequent colonization of the brain.

Keywords: breast cancer, BRD8, brain metastasis, central nervous system metastasis, systems biology of breast cancer, targeted therapeutics in breast cancer.

Patients with metastatic breast cancer are provided limited treatment options^{4,5}. One study reported that the rate of metastasis to the central nervous system in patients treated with trastuzumab, a monoclonal antibody targeting the human epidermal growth factor receptor 2 (HER2), was 34%². The severity of disease and limited treatment options demand an enhanced understanding of the fundamental transcriptional nature of metastasis to the brain in human breast cancer. We mined published microarray data^{6,7} generated using the primary tumors and brain metastatic tissues of patients with human breast cancer to identify genes whose expression was most quantitatively different in metastasis to the brain. These analyses revealed significant differential expression of the bromodomain protein BRD8 in brain metastatic tumor tissues of patients with human breast cancer.

Methods

We used datasets GSE43837⁶ and GSE125989⁷ for this differential gene expression analysis of brain metastasis in human breast cancer in conjunction with GEO2R. GSE43837 was generated using Affymetrix Human X3P Array technology with $n=19$ primary breast tumors and $n=19$ brain metastases from patients with breast cancer, analysis performed using platform GPL1352. GSE125989 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=16$ primary tumors from patients with breast cancer and $n=16$ brain metastases from patients with brain metastatic breast cancer. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether BRD8 gene expression was significantly between primary breast tumors and brain metastases in humans with breast cancer using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455). For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁸ for correlation of BRD8 mRNA expression levels with relapse (recurrence)-free survival (RFS) in $n=3951$ breast cancer patients.

Results

We performed comparative transcriptome analysis of primary and metastatic tumor tissues from patients with breast cancer using published microarray data^{6,7} to describe the transcriptional landscape of brain metastasis in human breast cancer and for the identification of potential therapeutic targets.

BRD8 is differentially expressed in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

We identified the gene encoding the bromodomain protein BRD8 as among the genes whose expression was most significantly different in the brain metastatic tissues of patients with human breast cancer as compared to primary tumors of the breast. When sorting each of the genes expressed in brain metastases based on significance of change in expression as compared to primary tumors of the breast, BRD8 ranked 46 out of 61359 total transcripts (Table 1), equating to 99.93% differential expression. Differential expression of BRD8 in metastases to the brain was statistically significant (Table 1; $p=0.00016466$).

We mined a second microarray dataset to validate differential expression of BRD8 in brain metastatic breast cancer. We again identified BRD8 as among the genes whose expression was most different in the brain metastatic tissues of patients with human breast cancer as compared to primary tumors of the breast. When sorting each of the genes expressed in brain metastases based on significance of change in expression as compared to primary tumors of the breast, in this dataset, BRD8 ranked 451 out of 22277 total transcripts (Table 2), equating to 98.0% differential expression. Differential expression

of BRD8 in metastases to the brain was again statistically significant (Table 2; $p=6.77E-03$).

BRD8 is expressed at lower levels in metastases to the brain as compared to primary tumors of the breast.

We obtained exact mRNA expression values for BRD8 from primary tumors of the breast and from brain metastases of patients with breast cancer to evaluate magnitude and direction of change in BRD8 mRNA expression in metastasis to the brain. BRD8 was expressed at lower levels in metastasis to the brain as compared to primary tumors of the breast (Figure 1), and this difference approached the level of statistical significance (Figure 1; $p=0.0571$). We calculated a mean fold change of 0.44 ± 0.79 in BRD8 mRNA expression in brain metastases relative to primary tumors of the breast.

BRD8 expression associates with survival outcomes in breast cancer.

We performed Kaplan-Meier survival analysis to determine whether BRD8 mRNA expression levels correlated with survival outcomes in patients with breast cancer. In 3951 breast cancer patients, we observed a statistically significant correlation between BRD8 expression and relapse-free survival (RFS) (Figure 2; log rank p -value: 0.00088 for relapse-free survival, hazard ratio: 0.83 (0.75 - 0.93) (Fig. 2)). Lower primary tumor BRD8 expression, like which we observed in metastases to the brain, was a negative prognostic indicator in patients with breast cancer. Median RFS was 39.26 months for patients with low tumor expression of BRD8 while median RFS was 61.2 months for patients with high tumor expression of BRD8 (Table 3).

Thus, through analysis of published microarray datasets, we found that BRD8 was among the genes whose expression was most significantly different in the brain metastases of patients with metastatic breast cancer as compared to primary tumors of the breast, that BRD8 expression was significantly lower in brain metastatic tissues as compared to primary tumors of the breast, and that low primary tumor expression of BRD8 significantly correlated with worse relapse-free survival.

Discussion

BRD8 is a subunit of the human NuA4 histone acetyltransferase (HAT) complex which include TRRAP, p400/Domino, EPC1, Tip60, DMAP1, ING3, RUVBL1, RUVBL2, BAF53a, Actin, MRG15 and hEaf6 and GAS41⁹. The Tip60/NuA4 HAT complex functions in acetylation of histone H4 and histone H2A.Z, the amino-terminal tail of histone H2, as well as non-histone substrates including the cytosolic PEPCK PCK1⁹⁻¹¹.

Depletion of BRD8 results in induction of p21, cell cycle arrest in G1/S, and activation of the DNA damage response (DDR)¹⁰; this may be cell-type specific behavior as it is unclear how this reconciles with a specific down-regulation of BRD8 in metastasis to the brain. Expression of BRD8 is higher in multiple colorectal cancer cell lines (e.g., DLD1, HCA7, HCT116) as compared to cancer cell lines of non-colorectal origin including HeLa, BcPC3, and SkBR3, and depletion of Brd8 *in vitro* resulted in induction of some degree of cell death through apoptosis¹². In endometrial stromal sarcoma, investigators have described fusion of BRD8 to PHF1¹³.

We used published microarray data here to demonstrate that BRD8 is among the genes whose expression is most significantly different in metastasis to the brain as compared to primary tumors of the breast, that BRD8 expression is markedly decreased in brain metastatic tissues as compared to primary tumors of the breast and that decreased primary tumor BRD8 expression is significantly correlated with inferior relapse-free survival. More thorough understanding of BRD8 transcriptional functions in the breast, in the brain, in primary tumors of the breast and in brain metastases may provide insights into mechanisms that contribute to spread of breast cancers to the central nervous system.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Robson, M., Im, S.A., Senkus, E., Xu, B., Domchek, S.M., Masuda, N., Delaloge, S., Li, W., Tung, N., Armstrong, A. and Wu, W., 2017. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *New England Journal of Medicine*, 377(6), pp.523-533.
5. Turner, N.C., Ro, J., André, F., Loi, S., Verma, S., Iwata, H., Harbeck, N., Loibl, S., Huang Bartlett, C., Zhang, K. and Giorgetti, C., 2015. Palbociclib in hormone-receptor-positive advanced breast cancer. *New England Journal of Medicine*, 373(3), pp.209-219.
6. McMullin, R.P., Wittner, B.S., Yang, C., Denton-Schneider, B.R., Hicks, D., Singavarapu, R., Moulis, S., Lee, J., Akbari, M.R., Narod, S.A. and Aldape, K.D., 2014. A BRCA1 deficient-like signature is enriched in breast cancer brain metastases and predicts DNA damage-induced poly (ADP-ribose) polymerase inhibitor sensitivity. *Breast Cancer Research*, 16(2), pp.1-10.
7. Iwamoto, T., Niikura, N., Ogiya, R., Yasojima, H., Watanabe, K.I., Kanbayashi, C., Tsuneizumi, M., Matsui, A., Fujisawa, T., Iwasa, T. and Shien, T., 2019. Distinct gene expression profiles between primary breast cancers and brain metastases from pair-matched samples. *Scientific reports*, 9(1), pp. 1-8.
8. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.
9. Doyon, Y., Selleck, W., Lane, W.S., Tan, S. and Côté, J., 2004. Structural and functional conservation of the NuA4 histone acetyltransferase complex from yeast to humans. *Molecular and cellular biology*, 24(5), pp.1884-1896.
10. Lashgari, A., Fauteux, M., Maréchal, A. and Gaudreau, L., 2018. Cellular Depletion of BRD8 causes p53-dependent apoptosis and induces a DNA damage response in non-stressed cells. *Scientific reports*, 8(1), pp.1-12.
11. Lin, Y.Y., Lu, J.Y., Zhang, J., Walter, W., Dang, W., Wan, J., Tao, S.C., Qian, J., Zhao, Y., Boeke, J.D. and Berger, S.L., 2009. Protein acetylation microarray reveals that NuA4 controls key metabolic target regulating gluconeogenesis. *Cell*, 136(6), pp.1073-1084.
12. Yamada, H. and Rao, C., 2008. Brd8 is a chemosensitizing target for spindle poisons in colorectal tumors.
13. Micci, F., Brunetti, M., Dal Cin, P., Nucci, M.R., Gorunova, L., Heim, S. and Panagopoulos, I., 2017. Fusion of the genes BRD8 and PHF1 in endometrial stromal sarcoma. *Genes, Chromosomes and Cancer*, 56(12), pp.841-845.

Rank	ID	p-value	t	B	FC	Gene	Gene name
46	Hs.144662.0.S1_3p_at	0.00016466	-4.1300122	0.60687	0.44 ± 0.79	BRD8	bromodomain containing 8

Table 1: Differential expression of BRD8 in brain metastatic tissues as compared to primary tumors in patients with breast cancer.

Rank of differential expression, probe ID, *p*-value with respect to differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, fold change (FC) of BRD8 expression between the brain metastases of patients with breast cancer and primary tumors of the breast, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
451	202227_s_at	6.77E-03	2.8808051	-2.3985	BRD8	bromodomain containing 8

Table 2: Differential expression of BRD8 in brain metastatic tissues as compared to primary tumors in patients with breast cancer.

Rank of differential expression, probe ID, *p*-value with respect to differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: BRD8 is expressed at lower levels in brain metastases as compared to primary tumors of the breast in metastatic breast cancer.

The mRNA expression level of BRD8 in primary tumors of the breast (left) and in the brain metastases of brain metastatic breast cancer patients (right) is graphically represented here with mean BRD8 levels marked and the result of a statistical test evaluating difference in mRNA expression level between brain metastases and primary tumors of the breast, a p -value, listed above. The Y-axis is set to natural logarithm (ln).

Figure 2: Expression of BRD8 (SMAP) in primary tumors of the breast significantly correlates with relapse-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of relapse-free survival (RFS) for $n=3951$ total patients stratified into two groups, based on low or high expression of BRD8, also known as SMAP, in primary tumors of patients with breast cancer. The log rank p -value denoting statistical significance of difference in relapse-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on BRD8 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low expression cohort (months)	High expression cohort (months)
39.26	61.2

Table 3: Median relapse-free survival is greater in breast cancer patients whose primary tumors express high levels of BRD8.

The median relapse-free survival of $n=3951$ breast cancer patients based on stratification into low or high tumor expression of BRD8 is listed in this chart.