Riemann Hypothesis proof using Balazard, Saias and Yor criterion

Shekhar Suman
Email: shekharsuman068@gmail.com

11 June 2021

Abstract

In this manuscript, we define a conformal map from the punctured unit disc onto the punctured semi plane. Later, we define a function $f(z) = (s-1)\zeta(s)$. We apply a method similar to Jensen's formula using Blaschke products and Cauchy's residue theorem to a cross-cut contour. Finally using the Balazard Saias and Yor logarithmic integral of Riemann zeta function as a sum over its non trivial zeros, we get

$$\int_{-\infty}^{\infty} \frac{\log |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = 0$$

Keywords: Blaschke product, Jensen's formula, Schwarz reflection principle, Cauchy's residue theorem, Lebesgue's dominated convergence theorem, Harmonic function, Critical strip, Critical line, Riemann zeta function, Riemann Hypothesis.

Mathematics Subject Classification: 11M26, 11M06

1 Introduction

The Riemann zeta function, $\zeta(s)$ is defined as the analytic continuation of the Dirichlet series

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s}$$

which converges in the half plane $\Re(s) > 1$. The Riemann zeta function is a meromorphic function on the whole complex s-plane, which is holomorphic everywhere except for a simple pole at $s = 1$ with residue 1. All the non trivial zeros of the Riemann zeta function lie in the critical strip $0 < \Re(s) < 1$. The Riemann Hypothesis states that all the non trivial zeros of the Riemann zeta function lies on the critical line $\Re(s) = \frac{1}{2}$.

Levinson [6], in 1974 proved that more than one third of zeros of Riemann zeta function are on the critical line. Balazard et al.[1] in 1999 proved an equivalent of the Riemann Hypothesis. Shaoji Feng [7], in 2012 proved that atleast 41.28 % of the zeros of Riemann zeta function are on the critical line. Pratt et al.[8] in 2020 proved that more than five-twelfths of the zeros are on the critical line.

2 Main Result

Let, $\sum_{\Re(\rho) > \frac{1}{2}}$ be the sum over the hypothetical zeros with real part greater than $\frac{1}{2}$ of the Riemann zeta function, $\zeta(s)$. In the sum, zeros of multiplicity m are counted m times. Balazard et al.[1, p.1] proved that,

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\log |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = \sum_{\Re(\rho) > \frac{1}{2}} \log \left| \frac{\rho}{1 - \rho} \right| \quad (1)$$

and the Riemann Hypothesis is true if and only if [1, p.1],

$$\int_{-\infty}^{\infty} \frac{\log |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = 0 \quad (2)$$

The goal of this paper is to prove the following result.

Theorem 1: If $\zeta(s)$ denotes the Riemann zeta function then

$$\int_{-\infty}^{\infty} \frac{\log |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = 0$$

We start the proof of Theorem 1 as follows: Let, $z \in \mathbb{D}$ where $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$. Considering the positive real axis as the branch cut or $0 < \arg z < 2\pi$ write,

$$s = s(z) = \frac{1}{2} + \frac{1 + \sqrt{z}}{2(1 - \sqrt{z})} = \frac{1}{1 - \sqrt{z}}$$

Define a semi plane as $\mathbb{P} = \{s \in \mathbb{C} \mid \Re(s) > \frac{1}{2}\}$.

Lemma 1.1: Map $s : \mathbb{D} \rightarrow \mathbb{P}$ is injective and onto.

Proof. For proving the map injective, let, $s(z) = s(z')$ where $z, z' \in \mathbb{D}$. Write $z = re^{i\theta}$ and $z' = r'e^{i\theta'}$ so we get, $\sqrt{r}e^{i\theta/2} = \sqrt{r'}e^{i\theta'/2}$ and taking modulus we have $\sqrt{r} = \sqrt{r'}$ or $r = r'$ and hence $e^{i\theta/2} = e^{i\theta'/2}$. Which gives $\cos(\frac{\theta - \theta'}{2}) = 1$ and $\sin(\frac{\theta - \theta'}{2}) = 0$. Since, $\theta, \theta' \in (0, 2\pi)$ so we get $\theta = \theta'$ and hence $z = z'$.

For onto, let, $s_0 \in \mathbb{P}$ then there exists $z_0 \in \mathbb{D}$ such that $\sqrt{z_0} = \left(\frac{s_0 - 1}{s_0}\right)$ and $s(z_0) = s_0$. □

Denote $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ (where $\mathbb{R}_{\geq 0} = \{x \in \mathbb{R} \mid x \geq 0\}$) as a slit disk having all points that are in disc $\mathbb{D}$ except the non negative reals which means $\mathbb{D} \setminus [0, 1)$. Denote $\overline{\mathbb{D}} = \{z \in \mathbb{C} \mid |z| \leq 1\}$.

Now we consider the function,

$$f(z) = (s - 1)\zeta(s)$$

where $s = \frac{1}{1 - \sqrt{z}}$ then,

$$f(z) = \left(\frac{\sqrt{z}}{1 - \sqrt{z}}\right) \zeta\left(\frac{1}{1 - \sqrt{z}}\right) \quad (3)$$

Lemma 1.2: $f(z) = \left(\frac{\sqrt{z}}{1 - \sqrt{z}}\right) \zeta\left(\frac{1}{1 - \sqrt{z}}\right)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$.

Proof. Any $z \in \overline{\mathbb{D}} \setminus \mathbb{R}_{\geq 0}$ can be written uniquely as $z = re^{i\theta}$, where $r > 0$ and $\theta \in (0, 2\pi)$. Next, we define a function $h : \overline{\mathbb{D}} \setminus \mathbb{R}_{\geq 0} \rightarrow \mathbb{C}$ as follows:

$$\forall (r, \theta) \in \mathbb{R}_{>0} \times (0, 2\pi) : \quad h(re^{i\theta}) = \sqrt{r}e^{i\theta/2} \quad (4)$$

$$= \sqrt{r} \cos\left(\frac{\theta}{2}\right) + i \left[\sqrt{r} \sin\left(\frac{\theta}{2}\right) \right] \quad (5)$$

$$= u(r, \theta) + i \cdot v(r, \theta). \quad (6)$$

Now, clearly the functions u and v satisfy the polar version of the Cauchy-Riemann Equations:

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta} \quad \text{and} \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}.$$

So, by Goursat's theorem, function h is holomorphic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$.

Since, $s(z) = \frac{1}{1 - \sqrt{z}}$ and $h(z) = \sqrt{z}$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$, also $q(z) = \frac{1}{1 - z}$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ and hence $s(z) = q(h(z))$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ and since $k(z) = (z - 1)\zeta(z)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$, so the composition $k(s(z)) = f(z)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$. □

Denote a set as $\mathbb{R}_{\geq 1} = \{x \in \mathbb{R} \mid x \geq 1\}$.

Lemma 1.3: Map $s : \mathbb{D} \setminus \mathbb{R}_{\geq 0} \rightarrow \mathbb{P} \setminus \mathbb{R}_{\geq 1}$ is conformal which takes the disc $\mathbb{D}$ to the plane $\mathbb{P}$.

Proof. Since $s(z)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ so we have,

$$s'(z) = \frac{-1}{2\sqrt{z}(1-\sqrt{z})^2}$$

which is non zero and does not tends to infinity for all $z \in \mathbb{D} \setminus \mathbb{R}_{\geq 0}$ and hence conformal.

Also $s = \frac{1}{1-\sqrt{z}}$, hence $z = \left(\frac{s-1}{s}\right)^2$ so that $|z| < 1$ if and only if $\Re(s) > \frac{1}{2}$.

Next we prove that if $0 < \arg(z) < 2\pi$ then $-\frac{\pi}{2} < \arg(s) < \frac{\pi}{2}$

We have shown earlier that $(\arg(s)) \subseteq (-\frac{\pi}{2}, \frac{\pi}{2})$

Next by Appendix (a) for any $\phi \in (-\frac{\pi}{2}, \frac{\pi}{2})$ there exists $z \in \mathbb{C}$ such that $|z| < 1$ and $\arg(z) = \phi$ □

Now we proceed to next lemma

Lemma 1.4:

$$\frac{1}{2\pi} \int_0^{2\pi} \log_e |f(e^{i\theta})| d\theta = \log_e |f(0)| + \sum_{|\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{|\alpha_n|}$$

Proof. Let, $R > 1$ and $U = \mathbb{D}_R \setminus \mathbb{R}_{\geq 0}$ where $\mathbb{D}_R = \{z \in \mathbb{C} \mid |z| < R\}$ be a simply connected [14, p.108] open subset of complex plane. Consider a simple closed contour C contained in U consisting of two concentric circles, a bigger circle C_1 of radius 1 unit and a smaller circle C_ϵ of radius ϵ where $0 < \epsilon < 1$ and having an infinitesimally small cross cut to join C_1 and C_ϵ . Let, this cross-cut be L_1 and L_2 . Then,

$$C = C_1 + L_1 - C_\epsilon + L_2, \text{ where } 0 < \epsilon < 1$$

Since f is analytic in domain $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$, so its zeros inside the disc $\mathbb{D}$ (since f as defined in equation (3) has no zeros in $\mathbb{R}_{\geq 0}$ because $\zeta(s)$ has no zeros on $\mathbb{R}_{\geq 1}$, so the zeros of f on $\mathbb{D}$ and $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ are the same) are at-most countable say, α_n . If these zeros α_n inside the unit disc satisfy the Blaschke condition [4, p.19],

$$\sum_{|\alpha_n| < 1, f(\alpha_n)=0} (1 - |\alpha_n|) < \infty \quad (7)$$

then we define a Blaschke product,

$$B(z) = \prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left(\frac{\alpha_n - z}{1 - \overline{\alpha_n}z} \right) \frac{|\alpha_n|}{\alpha_n}$$

where in the above product, zeros of multiplicity m are counted m times. Define a function,

$$g(z) := \frac{1}{f(z)} \prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left(\frac{\alpha_n - z}{1 - \overline{\alpha_n}z} \right) \frac{|\alpha_n|}{\alpha_n} \quad (8)$$

where in the above product, zeros of multiplicity m are counted m times. Since $f(z)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ and when inequality (7) holds, then $B(z)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ and $|B(z)| < 1$ for $|z| < 1$ and has no zeros except at the points α_n [4, p.19], so $g(z)$ is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$. Also, $g(z) \neq 0$ for all $z \in \mathbb{D} \setminus \mathbb{R}_{\geq 0}$ (see Appendix (b)).

Let, $R > 1$ and $U = \mathbb{D}_R \setminus \mathbb{R}_{\geq 0}$ where $\mathbb{D}_R = \{z \in \mathbb{C} \mid |z| < R\}$ be a simply connected [14, p.108] open subset of complex plane containing the closed curve C . By Cauchy's residue theorem since $\frac{\log_e g(z)}{z}$ is analytic on and inside the simple closed contour C , and there are no poles on and inside C , so,

$$\begin{aligned} \int_C \frac{\log_e g(z)}{z} dz &= 0 \\ \Rightarrow \int_{C_1} \frac{\log_e g(z)}{z} dz - \int_{C_\epsilon} \frac{\log_e g(z)}{z} dz + \int_{L_1} \frac{\log_e g(z)}{z} dz + \int_{L_2} \frac{\log_e g(z)}{z} dz &= 0 \end{aligned} \quad (9)$$

On C_1 , we have $z = e^{i\theta}$ and on C_ϵ we have $z = \epsilon.e^{i\theta}$ so we have

$$i. \int_0^{2\pi} \log_e g(e^{i\theta}) d\theta - i. \int_0^{2\pi} \log_e g(\epsilon.e^{i\theta}) d\theta + \int_{L_1} \frac{\log_e g(z)}{z} dz + \int_{L_2} \frac{\log_e g(z)}{z} dz = 0$$

As $\epsilon \rightarrow 0^+$ we have an overlap in L_1 and L_2 , but they both are in opposite sense and hence, $\lim_{\epsilon \rightarrow 0^+} \left(\int_{L_1} \frac{\log_e g(z)}{z} dz + \int_{L_2} \frac{\log_e g(z)}{z} dz \right) = 0$, so we get,

$$\int_0^{2\pi} \log_e g(e^{i\theta}) d\theta = \lim_{\epsilon \rightarrow 0^+} \int_0^{2\pi} \log_e g(\epsilon.e^{i\theta}) d\theta$$

Taking real parts on both sides we get,

$$\int_0^{2\pi} \log_e |g(e^{i\theta})| d\theta = \lim_{\epsilon \rightarrow 0^+} \int_0^{2\pi} \log_e |g(\epsilon.e^{i\theta})| d\theta$$

By Appendix (c), using Lebesgue's dominated convergence theorem, we have,

$$\int_0^{2\pi} \log_e |g(e^{i\theta})| d\theta = 2\pi \log_e |g(0)| \quad (10)$$

On, $|z| = 1$ we have,

$$\left| \frac{\alpha_n - z}{1 - \overline{\alpha_n} z} \right| = 1$$

So equation (8) gives,

$$|g(e^{i\theta})| = \frac{1}{|f(e^{i\theta})|} \quad (11)$$

Also from equation (8) we have,

$$|g(0)| = \frac{1}{|f(0)|} \prod_{|\alpha_n| < 1, f(\alpha_n) = 0} |\alpha_n| \quad (12)$$

Using equation (11) and (12) in equation (10), we have

$$\frac{1}{2\pi} \int_0^{2\pi} \log_e |f(e^{i\theta})| d\theta = \log_e |f(0)| + \sum_{|\alpha_n| < 1, f(\alpha_n) = 0} \log_e \frac{1}{|\alpha_n|} \quad (13)$$

where in the sum, zeros of multiplicity m are counted m times. □

Now we proceed to a useful lemma.

Lemma 1.5:

$$\int_0^{2\pi} \log_e |f(e^{i\theta})| d\theta = \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt$$

Proof. Let,

$$I = \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt$$

Since by contour integration or by substitution, $t = \frac{\tan \theta}{2}$ we have [13, p.550],

$$\int_{-\infty}^{\infty} \frac{\log_e |\frac{1}{2} + it|}{\frac{1}{4} + t^2} dt = 0$$

so we can write

$$I = \int_{-\infty}^{\infty} \frac{\log_e |(\frac{1}{2} + it)\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt$$

Substituting $t = \frac{\tan \theta}{2}$ we get,

$$I = 2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log_e \left| \left(\frac{e^{i\theta}}{e^{i\theta} + e^{-i\theta}} \right) \zeta \left(\frac{e^{i\theta}}{e^{i\theta} + e^{-i\theta}} \right) \right| d\theta$$

Using Schwarz reflection principle $\overline{\zeta(s)} = \zeta(\bar{s})$ we get,

$$I = 2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log_e \left| \left(\frac{1}{e^{2i\theta} + 1} \right) \zeta \left(\frac{1}{e^{2i\theta} + 1} \right) \right| d\theta$$

Substituting $\phi = 2\theta$ we get,

$$\begin{aligned} I &= \int_{-\pi}^{\pi} \log_e \left| \left(\frac{1}{1 + e^{i\phi}} \right) \zeta \left(\frac{1}{1 + e^{i\phi}} \right) \right| d\phi \\ \Rightarrow I &= \int_{-\pi}^{\pi} \log_e \left| \left(\frac{1}{2} - \frac{i}{2} \tan \left(\frac{\phi}{2} \right) \right) \zeta \left(\frac{1}{2} - \frac{i}{2} \tan \left(\frac{\phi}{2} \right) \right) \right| d\phi \end{aligned}$$

Substitute $\phi = \frac{\alpha}{2}$ so we get,

$$I = \frac{1}{2} \int_{-2\pi}^{2\pi} \log_e \left| \left(\frac{1}{2} - \frac{i}{2} \tan \left(\frac{\alpha}{4} \right) \right) \zeta \left(\frac{1}{2} - \frac{i}{2} \tan \left(\frac{\alpha}{4} \right) \right) \right| d\alpha$$

By Schwarz reflection principle since the integrand is an even function so we have,

$$I = \int_0^{2\pi} \log_e \left| \left(\frac{1}{2} - \frac{i}{2} \tan \left(\frac{\alpha}{4} \right) \right) \zeta \left(\frac{1}{2} - \frac{i}{2} \tan \left(\frac{\alpha}{4} \right) \right) \right| d\alpha$$

Since, $\int_0^{2\pi} f(\alpha) d\alpha = \int_0^{2\pi} f(2\pi - \alpha) d\alpha$ so we have,

$$I = \int_0^{2\pi} \log_e \left| \left(\frac{1}{2} - \frac{i}{2} \cot \left(\frac{\alpha}{4} \right) \right) \zeta \left(\frac{1}{2} - \frac{i}{2} \cot \left(\frac{\alpha}{4} \right) \right) \right| d\alpha$$

Schwarz reflection principle gives,

$$I = \int_0^{2\pi} \log_e \left| \left(\frac{1}{2} + \frac{i}{2} \cot \left(\frac{\alpha}{4} \right) \right) \zeta \left(\frac{1}{2} + \frac{i}{2} \cot \left(\frac{\alpha}{4} \right) \right) \right| d\alpha$$

Observe that,

$$\begin{aligned} \frac{1}{1 - e^{\frac{i\alpha}{2}}} &= \frac{1}{2} + \frac{i}{2} \cot \left(\frac{\alpha}{4} \right) \quad \text{and} \quad |e^{\frac{i\alpha}{2}}| = 1 \\ \Rightarrow I &= \int_0^{2\pi} \log_e \left| \left(\frac{e^{\frac{i\alpha}{2}}}{1 - e^{\frac{i\alpha}{2}}} \right) \zeta \left(\frac{1}{1 - e^{\frac{i\alpha}{2}}} \right) \right| d\alpha \\ \Rightarrow \int_0^{2\pi} \log_e |f(e^{i\theta})| d\theta &= \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt \end{aligned} \tag{14}$$

□

We proceed to another lemma

Lemma 1.6:

$$\log_e |f(0)| = 0$$

Proof. Let, $\epsilon > 0$ be given. Consider for $0 < \arg z < 2\pi$ and $z \in \mathbb{D}$, We have $|z| \leq r < 1$ and since $s(z) = \frac{1}{1-\sqrt{z}}$ thus, $s(0) = 1$ so,

$$|s(z) - s(0)| = \left| \frac{\sqrt{z}}{1 - \sqrt{z}} \right| \leq \frac{|z|^{1/2}}{|1 - |z|^{1/2}|} \leq \frac{|z|^{1/2}}{1 - \sqrt{r}}$$

Choose $\delta = \min((1 - \sqrt{r})^2 \epsilon^2, r)$ then we have

$|s(z) - s(0)| < \epsilon$ whenever $|z| < \delta$. So, $s(z)$ is continuous at 0.

Define $p(z) := (z - 1)\zeta(z)$. Since $f(z) = (s - 1)\zeta(s)$ where $s = \frac{1}{1-\sqrt{z}}$ so, $p(s(z)) = f(z)$. Since $s(z)$ is continuous at 0 and $p(z)$ is continuous at $s(0) = 1$, so we have the composition $p(s(z)) = f(z)$ is continuous at 0. Hence,

$$f(0) = \lim_{z \rightarrow 0} f(z) = \lim_{s \rightarrow 1} (s - 1)\zeta(s) = 1$$

So,

$$\log_e |f(0)| = 0 \tag{15}$$

□

Since,

$$f(z) = \left(\frac{\sqrt{z}}{1 - \sqrt{z}} \right) \zeta \left(\frac{1}{1 - \sqrt{z}} \right)$$

Since, $f(0) \neq 0$, so $f(\alpha) = 0$ corresponds to $\zeta \left(\frac{1}{1 - \sqrt{\alpha}} \right) = 0$.

Let, ρ denote non trivial zeros of Riemann zeta function then,

$$\rho = \frac{1}{1 - \sqrt{\alpha}}$$

Since the non trivial zeros of Riemann zeta function are countable so,

$$\rho_n = \frac{1}{1 - \sqrt{\alpha_n}}$$

We proceed to another lemma

Lemma 1.7:

$$\sum_{|\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{|\alpha_n|} = 2 \sum_{\Re(\rho_n) > \frac{1}{2}, \zeta(\rho_n)=0} \log_e \left| \frac{\rho_n}{1 - \rho_n} \right|$$

Proof. We have $f(\alpha_n) = 0$ if and only if $\zeta(\rho_n) = 0$. Also there does not exist any non trivial zero ρ_n of $\zeta(s)$ such that $\rho_n(\in \mathbb{R}) \geq 1$, so there does not exist $\alpha_n \in \mathbb{D}$ such that $\alpha_n(\in \mathbb{R}) \geq 0$ and $f(\alpha_n) = 0$. Since, $\rho_n = \frac{1}{1 - \sqrt{\alpha_n}}$ so we have $\sqrt{|\alpha_n|} = \left| \frac{1 - \rho_n}{\rho_n} \right|$. By Lemma 1.3, the map $s : \mathbb{D} \setminus \mathbb{R}_{\geq 0} \rightarrow \mathbb{P} \setminus \mathbb{R}_{\geq 1}$ defined as $s(z) = \frac{1}{1 - \sqrt{z}}$ is conformal so we have,

$$\begin{aligned} \sum_{|\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{|\alpha_n|} &= \sum_{0 < |\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{|\alpha_n|} = 2 \sum_{0 < |\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{\sqrt{|\alpha_n|}} \\ &\Rightarrow \sum_{|\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{|\alpha_n|} = 2 \sum_{\Re(\rho_n) > \frac{1}{2}, \rho_n \neq 1, \zeta(\rho_n)=0} \log_e \left| \frac{\rho_n}{1 - \rho_n} \right| \end{aligned}$$

Since, $\rho_n \neq 1$ as $0 < \Re(\rho_n) < 1$, so we get,

$$\sum_{|\alpha_n| < 1, f(\alpha_n)=0} \log_e \frac{1}{|\alpha_n|} = 2 \sum_{\Re(\rho_n) > \frac{1}{2}, \zeta(\rho_n)=0} \log_e \left| \frac{\rho_n}{1 - \rho_n} \right| \tag{16}$$

□

Next we prove that equation (7) is satisfied for $|\alpha_n| = \left| \frac{1-\rho_n}{\rho_n} \right|^2$

Lemma 1.8: For $|\alpha_n| = \left| \frac{1-\rho_n}{\rho_n} \right|^2$,

$$\sum_{|\alpha_n| < 1, f(\alpha_n)=0} (1 - |\alpha_n|) < \infty$$

Proof. For, $|\alpha_n| = \left| \frac{1-\rho_n}{\rho_n} \right|^2$, since $0 < \Re(\rho_n) < 1$,

$$\sum_{|\alpha_n| < 1, f(\alpha_n)=0} (1 - |\alpha_n|) = \sum_{\Re(\rho_n) > \frac{1}{2}, \zeta(\rho_n)=0} \frac{2\Re(\rho_n) - 1}{|\rho_n|^2} < \sum_{\Re(\rho_n) > \frac{1}{2}, \zeta(\rho_n)=0} \frac{1}{|\rho_n|^2} \leq \sum_{0 < \Re(\rho_n) < 1, \zeta(\rho_n)=0} \frac{1}{|\rho_n|^2}$$

The sum on the right hand side is convergent [12, p.35]. $\square$

Using equation (14), (15) and (16) in equation (13), we get,

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = 2 \sum_{\Re(\rho_n) > \frac{1}{2}, \zeta(\rho_n)=0} \log_e \left| \frac{\rho_n}{1 - \rho_n} \right| \quad (17)$$

Since the non trivial zeros of zeta function are countable so, equation (1) can be written as [13, p.549]

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\log |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = \sum_{\Re(\rho_n) > \frac{1}{2}} \log \left| \frac{\rho_n}{1 - \rho_n} \right| \quad (18)$$

Combining equation (17) and (18) we get,

$$\begin{aligned} \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt \\ &\Rightarrow \int_{-\infty}^{\infty} \frac{\log_e |\zeta(\frac{1}{2} + it)|}{\frac{1}{4} + t^2} dt = 0 \end{aligned}$$

This proves equation (2) and completes the proof of Theorem 1. Hence the Riemann Hypothesis is true.

3 Appendix

Appendix (a): We prove that for $s(z) = \frac{1}{1-\sqrt{z}}$, for any $\phi \in (-\frac{\pi}{2}, \frac{\pi}{2})$ there exists $z \in \mathbb{C}$ such that $|z| < 1$ and $\arg(z) = \phi$.

Proof. Let us define a function $sign : (-\infty, \infty) \rightarrow \mathbb{R}$ as,

$$sign(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ -1 & \text{if } x < 0 \end{cases}$$

For any $\phi \in (-\frac{\pi}{2}, \frac{\pi}{2})$ there exists

$$z = -\sin^2(\phi) \cos(2\phi) + i \, sign(sin(\phi)) |\sin(\phi)| \sin(\phi) \sin(2\phi)$$

such that $|z| = \sin^2(\phi) < 1$

$$\Rightarrow z = \sin^2(\phi)(\sin^2(\phi) - \cos^2(\phi)) + i \, sign(sin(\phi)) |\sin(\phi)| \sin(\phi) \sin(2\phi)$$

$$\Rightarrow z = \sin^4(\phi) - \sin^2(\phi) \cos^2(\phi) + 2i \, sign(sin(\phi)) \sin^2(\phi) |\sin(\phi)| \cos(\phi)$$

$$\Rightarrow z = (\sin^2(\phi) + i \, sign(sin(\phi)) |\sin(\phi)| \cos(\phi))^2$$

there exists $\sqrt{z} = \sin^2(\phi) + i \, sign(sin(\phi)) |\sin(\phi)| \cos(\phi)$ such that

$$1 - \sqrt{z} = 1 - \sin^2(\phi) - i \, sign(sin(\phi)) |\sin(\phi)| \cos(\phi)$$

$$\Rightarrow 1 - \sqrt{z} = \cos^2(\phi) - i \, sign(sin(\phi)) |\sin(\phi)| \cos(\phi)$$

$$\Rightarrow 1 - \sqrt{z} = \cos^2(\phi) - i \sin(\phi) \cos(\phi)$$

$$\Rightarrow 1 - \sqrt{z} = \cos(\phi)(\cos(\phi) - i \sin(\phi))$$

$$\Rightarrow 1 - \sqrt{z} = \cos(\phi)e^{-i\phi}$$

Since, $\phi \in (-\frac{\pi}{2}, \frac{\pi}{2})$ so we have $\cos(\phi) > 0$ so we get $\arg(1 - \sqrt{z}) = -\phi$.

Hence, $\arg(s) = -\arg(1 - \sqrt{z}) = \phi$ □

Now we proceed to next appendix.

Appendix (b): $g(z) = \frac{1}{f(z)} \prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left(\frac{z - \alpha_n}{\overline{\alpha_n}z - 1} \right) \frac{|\alpha_n|}{\alpha_n}$ then $g(z) \neq 0$ for all $z \in \overline{\mathbb{D}} \setminus \mathbb{R}_{\geq 0}$.

Proof. If $z \neq \alpha_n$ for all n then since $B(z)$ has zeros only at α_n so, $B(z) \neq 0$ and $f(z) \neq 0$.

Assume on the contrary that there exists a zero of $g(z)$ (which is of the form α_m for some m) of multiplicity k in $\overline{\mathbb{D}} \setminus \mathbb{R}_{\geq 0}$. Since f is analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ and has a zero at α_m of multiplicity k , so by definition of zeros of an analytic function there is function h analytic on and inside of $\mathbb{D} \setminus \mathbb{R}_{\geq 0}$ such that $f(z) = (z - \alpha_m)^k h(z)$ where $h(\alpha_m) \neq 0$

$$g(z) = \frac{1}{h(z)(z - \alpha_m)^k} \prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left(\frac{z - \alpha_n}{\overline{\alpha_n}z - 1} \right) \frac{|\alpha_n|}{\alpha_n}$$

$$\Rightarrow g(z) = \frac{1}{h(z)} \left(\prod_{n \neq m, |\alpha_n| < 1, f(\alpha_n)=0} \left(\frac{z - \alpha_n}{\overline{\alpha_n}z - 1} \right) \frac{|\alpha_n|}{\alpha_n} \right) \left(\frac{1}{\overline{\alpha_m}z - 1} \frac{|\alpha_m|}{\alpha_m} \right)$$

which gives $g(\alpha_m) \neq 0$. □

Now we proceed to last appendix.

Appendix (c): By Lebesgue's dominated convergence theorem we show that,

$$\lim_{\epsilon \rightarrow 0^+} \int_0^{2\pi} \log_e |g(\epsilon.e^{i\theta})| d\theta = 2\pi \log_e |g(0)|$$

Proof. Define $k_\epsilon(\theta) = \log_e |g(\epsilon.e^{i\theta})|$. If we define $\log_e^+ x = \begin{cases} \log x & \text{if } x \geq 1 \\ 0 & \text{if } 0 \leq x < 1 \end{cases}$ and

$\log_e^- x = \max\{-\log x, 0\}$. We may write [13, p.75],

$$|k_\epsilon(\theta)| = |\log_e |g(\epsilon.e^{i\theta})|| = 2\log_e^+ |g(\epsilon.e^{i\theta})| - \log_e |g(\epsilon.e^{i\theta})|$$

Since, $\log |x| = \log^+ |x| - \log^- |x|$ so we get

$$\Rightarrow |k_\epsilon(\theta)| = \log_e^+ |g(\epsilon.e^{i\theta})| + \log_e^- |g(\epsilon.e^{i\theta})| = |l_\epsilon(\theta)| + |m_\epsilon(\theta)|$$

where $|l_\epsilon(\theta)| = \log_e^+ |g(\epsilon.e^{i\theta})|$ and $|m_\epsilon(\theta)| = \log_e^- |g(\epsilon.e^{i\theta})|$.

Using equation (8) we have,

$$|l_\epsilon(\theta)| = -\log_e^+ |f(\epsilon.e^{i\theta})| + \log_e^+ \left(\prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left| \frac{\alpha_n - \epsilon.e^{i\theta}}{1 - \overline{\alpha_n}\epsilon.e^{i\theta}} \right| \right)$$

$$\Rightarrow \int_0^{2\pi} |l_\epsilon(\theta)| d\theta \leq \int_0^{2\pi} \log_e^+ \left(\prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left| \frac{\alpha_n - \epsilon.e^{i\theta}}{1 - \overline{\alpha_n}\epsilon.e^{i\theta}} \right| \right) d\theta = \int_0^{2\pi} \log_e^+ |B(\epsilon.e^{i\theta})| d\theta$$

Now since $|B(z)| < 1$ for $|z| < 1$ [4, p.19] and $\epsilon < 1$, so by definition of $\log_e^+ x = 0$ if $0 \leq x < 1$ and $\log_e^+$ is monotonically increasing, the above equation gives,

$$\int_0^{2\pi} |l_\epsilon(\theta)| d\theta \leq 0$$

Similarly using equation (8) we have,

$$|m_\epsilon(\theta)| \leq \log_e^- \left(\prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left| \frac{\alpha_n - \epsilon.e^{i\theta}}{1 - \overline{\alpha_n}\epsilon.e^{i\theta}} \right| \right)$$

Now since $|B(z)| < 1$ for $|z| < 1$ [4, p.19] so we have

$$\begin{aligned} |m_\epsilon(\theta)| &\leq -\log_e \left(\prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left| \frac{\alpha_n - \epsilon.e^{i\theta}}{1 - \overline{\alpha_n}\epsilon.e^{i\theta}} \right| \right) \\ \Rightarrow \int_0^{2\pi} |m_\epsilon(\theta)| d\theta &\leq -\int_0^{2\pi} \log_e \left(\prod_{|\alpha_n| < 1, f(\alpha_n)=0} \left| \frac{\alpha_n - \epsilon.e^{i\theta}}{1 - \overline{\alpha_n}\epsilon.e^{i\theta}} \right| \right) d\theta \end{aligned}$$

Since $B(z)$ is analytic in $|z| < 1$ and is harmonic so by mean value theorem of harmonic functions and using Lemma 1.8, we have,

$$\int_0^{2\pi} |m_\epsilon(\theta)| d\theta \leq -2\pi \log_e \left(\prod_{|\alpha_n| < 1, f(\alpha_n)=0} |\alpha_n| \right) = 2\pi \sum_{|\alpha_n| < 1, f(\alpha_n)=0} \log \frac{1}{|\alpha_n|} = 4\pi \sum_{\Re(\rho_n) > \frac{1}{2}, \zeta(\rho_n)=0} \log \left| \frac{\rho_n}{1 - \rho_n} \right|$$

The last sum is finite due to Balazard et al. [1].

$$\Rightarrow \int_0^{2\pi} |m_\epsilon(\theta)| d\theta < \infty \Rightarrow \int_0^{2\pi} |k_\epsilon(\theta)| d\theta < \infty$$

Hence, by Lebesgue's dominated convergence theorem the result follows. $\square$

4 References

1. Balazard, M. Saias, E., and Yor, M. "Notes sur la fonction ζ de Riemann, 2", Advances in Mathematics , Volume 143 (1999) , 284-287.
2. Titchmarsh, E.C. The theory of functions, 2nd edition , Oxford science publication , 1939.
3. Hoffman, K. Banach Spaces of Analytic functions , Dover, New York (1988).
4. Peter Duren, Theory of H^p spaces, Vol. 38 , 1st edition, Academic Press (1970).
5. John B. Garnett, Bounded Analytic Functions, Vol.236 (2007).
6. Norman Levinson, More than one third of zeros of Riemann zeta function are on $\sigma = 1/2$, Advances in Mathematics (1974).
7. Shaoji Feng , Zeros of Riemann zeta function on the critical line, Journal of Number Theory (2012).
8. Kyle Pratt, Nicolas Robles, Alexandru Zaharescu , More than five-twelfths of zeros of ζ are on critical line , Research in the Mathematical Sciences, Springer (2020).
9. Titchmarsh, E.C. The theory of the Riemann zeta function , 2nd edition, revised by D.R. Heath Brown, Oxford university press (1986).
10. Rudin, W., Real and complex analysis, McGraw-Hill (1987),
11. Paul Koosis, Introduction to Hp Spaces, Cambridge University Press (1998).
12. Kevin Broughan, Equivalents of Riemann Hypothesis I , Arithmetic equivalents, Cambridge University press (2017).
13. K. Ilgar Eroglu and Iossif V. Ostrovskii, On an Application of the Hardy Classes to the Riemann Zeta-Function , Turk J Math (2001).
14. Joseph Bak, Donald J. Newman, Complex Analysis, Third edition, Springer (2010).