Compatibility of concurrent aerobic and strength training for skeletal muscle size and function: an updated systematic review and meta-analysis

Moritz Schumann^{1*}, Joshua F Feuerbacher¹, Marvin Sünkeler¹, Nils Freitag^{1,2}, Bent R Rønnestad³, Kenji Doma⁴ & Tommy R Lundberg^{5, 6}

1: Department of Molecular and Cellular Sports Medicine, German Sport University, Cologne

2: Olympic Training Centre Berlin, Germany

3: Section for Health and Exercise Physiology, Department of Public Health and Sport Sciences, Inland Norway University of Applied Sciences, Elverum, Norway

4: Sport and Exercise Science, College of Healthcare Sciences, James Cook University, Townsville, Australia.

5: Division of Clinical Physiology, Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden

6: Unit of Clinical Physiology, Karolinska University Hospital, Stockholm, Sweden

Short title: Compatibility of concurrent aerobic and strength training

*Correspondence to:

Moritz Schumann, PhD
Institute of Cardiovascular Research and Sports Medicine
Department of Molecular and Cellular Sports Medicine
German Sport University
Am Sportpark Müngersdorf 6
50933 Cologne, Germany
Tel: +49 221 4982 4821
E-Mail: m.schumann@dshs-koeln.de

Final version in press in Sports Medicine

Cite as:

Schumann, M., Feuerbacher, J. F., Sünkeler, M., Freitag, N., Rønnestad, B., Doma, K., & Lundberg, T. (2021, May 20). An updated systematic review and meta-analysis on the compatibility of concurrent aerobic and strength training for skeletal muscle size and function. <https://doi.org/10.31236/osf.io/e7tvr>

All authors have read and approved this version of the manuscript. This work is a PREPRINT.

DOI: <https://doi.org/10.31236/osf.io/e7tvr>

Abstract

Background

Both athletes and recreational exercisers often perform relatively high volumes of aerobic and strength training simultaneously. However, the compatibility of these two distinct training modes remains unclear.

Objective

This systematic review assessed the compatibility of concurrent aerobic and strength training compared with strength training alone, in terms of adaptations in muscle function (maximal and explosive strength) and muscle mass. Subgroup analyses were conducted to examine the influence of training modality, training type, exercise order, training frequency, age, and training status.

Design

A systematic literature search was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). PROSPERO: CRD42020203777.

Data sources

PubMed/MEDLINE, ISI Web of Science, Embase, CINAHL, SPORTDiscus and Scopus were systematically searched (12th of August 2020, updated on the 15th of March 2021).

Eligibility criteria

Population: Healthy adults of any sex and age; Intervention: supervised, concurrent aerobic and strength training of at least 4 weeks; Comparison: identical strength training prescription, with no aerobic training; Outcome: maximal strength, explosive strength, and muscle hypertrophy.

Results

A total of 43 studies were included. The estimated standardised mean differences (SMD) based on the random-effects model were -0.06 (95% CI: -0.20, 0.09, $p=0.446$), -0.28 (95% CI: -0.48, -0.08, $p=0.007$) and -0.01 (95% CI: -0.16, 0.18, $p=0.919$) for maximal strength, explosive strength, and muscle hypertrophy, respectively. Attenuation of explosive strength was more pronounced when concurrent training was performed within the same session ($p=0.043$) compared to when sessions were separated by at least 3 h ($p>0.05$). No significant effects were found for the other moderators, i.e. type of aerobic training (cycling vs. running), frequency of concurrent training (> 5 weekly sessions vs. < 5 weekly sessions), training status (untrained vs. active) and mean age (< 40 years vs. > 40 years).

Summary/Conclusion

Concurrent aerobic and strength training does not compromise muscle hypertrophy and maximal strength development. However, explosive strength gains may be attenuated, especially when aerobic and strength training are performed in the same session. These results appeared to be independent of the type of aerobic training, frequency of concurrent training, training status, and age.

Key points

- While concurrent aerobic and strength training is recommended to improve physical fitness and health, the compatibility of these two distinct training modes remains unclear.
- In this meta-analysis, we report that concurrent training does not interfere with adaptations in maximal strength and muscle hypertrophy, regardless of the type of aerobic training (cycling vs. running), frequency of concurrent training (> 5 weekly sessions vs. < 5 weekly sessions), training status (untrained vs. active), mean age (< 40 years vs. > 40 years), and training modality (same session vs. same day vs. different day training).
- However, concurrent training may attenuate gains in explosive strength, which is exacerbated when aerobic and strength training are performed within the same training session.

Declarations

Funding

No funding sources were used in the preparation of this article.

Conflicts of interest/Competing interests

Moritz Schumann, Joshua F. Feuerbacher, Marvin Sünkeler, Nils Freitag, Bent R. Rønnestad, Kenji Doma and Tommy Lundberg declare that they have no conflicts of interest relevant to the content of this review.

Availability of data and material: N/A

Code availability: The code will be available upon reasonable request.

Acknowledgements: The authors thank Dr. James Steele (Solent University, UK) for his valuable input concerning the data analysis.

Authors contribution: Design of the study: MS, JFF, MSü, NF, KD, BRR, TL; Literature search: MS, JFF, MSü; Data screening and extraction: JFF, MSü; Statistical analyses: MS, JFF, MSü, NF, TL; Manuscript preparation and editing: MS, JFF, TL; All authors have read and agreed to the submitted version. All authors read and approved the final manuscript.

Ethics approval: N/A

Consent to participate: N/A

Consent for publication: N/A

1. Introduction

Performing aerobic and strength training concurrently is an integrative part of physical training aimed at improving both athletic performance and health. The recommendation to perform both aerobic and strength training is important because these activities to some extent induce distinct adaptations and health benefits [1, 2]. For example, aerobic training promotes increased aerobic capacity (i.e. central adaptations) and metabolic changes in skeletal muscle, such as increased mitochondrial density and capillarisation [3]. Conversely, regular strength training results in muscle hypertrophy and increased strength and power [4], but may also improve bone mineral density [5]. The role of skeletal muscle in health maintenance has received increased attention in the last decade, with muscle tissue being understood as a secretory organ that releases several hundred of myokines related to the function of other organs, such as the brain, adipose tissue, bone, liver, gut, pancreas, vascular bed and skin [6]. In addition, the role of muscle power has recently been highlighted as being strongly associated with a lower risk of fall-related injuries in older adults [7, 8], further underlining the importance of both muscle mass and muscle function as indicators of physical health and independence in daily life.

Aside from the health perspective, many sports require the athlete to simultaneously incorporate divergent training modalities, including aerobic and strength training, into their training regimen. Considering that both athletes and recreational exercisers often perform relatively high volumes (and/or frequencies) of aerobic training alongside resistance-type training, it is pertinent to revisit the compatibility of aerobic and strength training. Aerobic training has been shown to interfere with the development of maximal strength when the overall training volume is high [9]. In contrast, no interference in maximal strength was observed when training volume was reduced to two weekly aerobic and strength training sessions, respectively [10–12]. Importantly, however, even low volumes of concurrent aerobic training have been shown to decrease gains in rapid force production [10, 13], which could translate into reduced muscle-power related benefits. Identifying additional moderators hypothesised in the literature to potentially influence neuromuscular adaptations to concurrent aerobic and strength training (such as type of aerobic training, concurrent training modality, age and training status) could further aid in fine-tuning exercise guidelines for health and/or fitness performance.

To date, scarce attempts have been made to quantitatively synthesise the literature concerning concurrent aerobic and strength training. The first meta-analysis conducted a decade ago by Wilson and colleagues showed that peak power was attenuated with concurrent training compared with strength training alone, while the development of muscle hypertrophy and maximal strength were not compromised [14]. A more recent meta-analysis aimed to compare the effect of concurrent aerobic and strength training with strength training alone on the development of maximal strength in untrained, moderately trained and trained individuals [15]. The results suggested that concurrent training may have a negative effect on lower body strength development in trained individuals, but not in moderately trained or untrained individuals. While this study updated information on the effect of training status on maximal strength development, several other important outcome variables related to muscle mass and function have not been examined in a meta-analysis since 2012. Therefore, the aim of the current study was to systematically assess the compatibility of concurrent aerobic and strength training on adaptations in maximal strength, explosive strength, and muscle hypertrophy by means of pooled analyses. Subgroup analysis was also conducted to examine the influence of aerobic training type, training modality, exercise order, concurrent training frequency, age, and training status. An updated literature synthesis on this topic is relevant to physicians, physiotherapists, exercise scientists and sports practitioners when designing programmes aimed at developing both aerobic and strength qualities for health purposes, rehabilitation, and/or fitness performance.

2. Methods

2.1 Systematic literature search

A systematic literature search was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) and was registered with the International Database of Prospectively Registered Systematic Reviews in Health and Social Care (PROSPERO: CRD42020203777). The PubMed/MEDLINE, ISI Web of Science, Embase, CINAHL, SPORTDiscus and Scopus databases were systematically searched using a search string specifically adapted to the search requirements of each database (Online supplemental Table S1).

The search was conducted on the 12th of August 2020 and updated on the 15th of March 2021. The literature search process was performed independently by two researchers and included saving the online search, removing duplicates and screening titles, abstracts, and full texts. Potential conflicts were resolved by consulting with a third author. In addition, a grey literature search was performed by screening Google Scholar and the reference lists of previously identified eligible full texts. A flowchart of the search process and study selection is shown in Figure 1.

Figure 1. Flowchart of the search process and the study selection.

2.2 Eligibility criteria

Inclusion criteria were defined based on the Population, Intervention, Control and Outcomes (PICO) criteria [16]. The population included healthy adults with no restrictions in terms of sex and age. The intervention had to consist of supervised, combined aerobic and strength training of at least 4 weeks. As a comparator, eligible studies had to include a group receiving the identical strength training prescription, with no aerobic training. Outcomes of interest included maximal strength, explosive strength, and muscle hypertrophy. The exercise tests included had to be specific to the training performed. For maximal strength, both isometric and isoinertial measurements were accepted. For explosive strength, any form of jump test, isometric rate of force development (RFD), or dynamic power measurements were considered eligible. For muscle hypertrophy, objective measurements of whole muscle cross-sectional area or muscle thickness (e.g. ultrasound, computed tomography (CT) or magnetic resonance imaging (MRI)) were required. In addition, segmental lean mass as determined by dual-energy X-ray absorptiometry (DXA) was accepted if values were reported separately for segments that were engaged in training.

Exclusion criteria included language other than English or German, abstracts and dissertations, cross-sectional studies assessing only acute exercise responses, and observational studies.

2.3 Data extraction

Data extraction was performed independently by two authors. The following data were extracted from each included study: (1) general characteristics (e.g. author(s), year of publication and aim of the study); (2) participant information (e.g. sample size, training status, and age); (3) intervention data for all groups (e.g. intervention duration, type of intervention); (4) specific outcomes (e.g. measures of maximal and explosive strength and hypertrophy). If the mean and standard deviation of each group were not specified, the authors of the primary studies were contacted to request baseline and post-intervention data. In case data were

presented within a graph and no additional data were provided upon request, mean and standard deviation were extracted using WebPlotDigitizer (Pacifica, California, USA, version: 4.4) [17].

2.4 Data synthesis and analyses

Standardised mean differences (SMD) were calculated, and an inverse variance-weighted random-effects model was fitted to the effect sizes (ES). Additionally, log variability ratios were calculated, and an inverse variance-weighted random-effects model was fitted to the ES. Meta-analyses were performed using R (3.6.2), RStudio (1.2.5033) and the metafor package (version 2.4.0) [18]. Effect sizes were calculated for *pre-test post-test control group designs* using the previously recommended raw score standardisation [19, 20]. Furthermore, exact sampling variance of effect sizes was computed according to recommendations [19].

Heterogeneity (i.e. τ^2) was estimated using the restricted maximum-likelihood estimator (REML) [21]. In order to complete the heterogeneity analyses, the Q-test for heterogeneity [22] and the I^2 statistic [23] were also calculated. Studentised residuals and Cook's distances were examined to assess whether studies might be outliers and/or overly influential [24]. Studies with a studentised residual greater than the $100 \times (1 - 0.05 / (2 \times k))^{\text{th}}$ percentile of a standard normal distribution were declared potential outliers (i.e. using a Bonferroni correction with two-sided $\alpha = 0.05$ for k studies included in the meta-analyses). Studies with a Cook's distance larger than the median plus six times the interquartile range of the Cook's distances were considered overly influential. In the event that a study was identified as a potential outlier or overly influential, a sensitivity analysis was performed. A trim-and-fill-contour funnel plot was created to estimate the number of studies that may be missing from the meta-analysis (Online Supplemental Figure S1). Rank correlation test [25] and regression test [26] using the standard error of observed outcomes as predictor were used to check for funnel plot asymmetry.

Effect sizes from studies with more than two intervention or control groups were combined according to the recommendations of the Cochrane Handbook [27], except for subgroup analysis when different interventions from individual studies were included in separate subgroups. In the case of multiple measurements for the same outcome, only one measurement was included in the analysis, based on the following hierarchy. For maximal strength, 1) dynamic bilateral leg press, 2) squat, 3) unilateral isometric torque (knee extension) and 4) bilateral dynamic knee extension. For explosive strength, 1) jump height and 2) other measures of rapid force production as well as squat jump power and leg press power at 50% of maximal strength. For muscle hypertrophy, 1) whole muscle cross-sectional area of the quadriceps femoris muscles (i.e. panoramic ultrasound, CT, MRI), 2) muscle thickness of the vastus lateralis, 3) segmental DXA of the lower extremities. Thus, each study was included in the final analyses with only one parameter to avoid inflating the weighting of individual studies.

Due to a lack of systematic reporting, subgroup analyses were only performed for aerobic training type (i.e. cycling vs. running), concurrent training frequency (i.e. low frequency of 4.1 ± 0.3 vs. high frequency of 6.1 ± 1.6 weekly sessions, based on 2.0 ± 0.3 vs. 3.1 ± 0.6 weekly sessions in the comparison training group), training status (i.e. untrained vs. active), mean age of the study population (18–40 years vs. >40 years) and training modality (i.e. concurrent training on different days vs. concurrent training on the same day vs. concurrent training in the same session). For studies comparing concurrent training in the same session, when a sufficient number of studies were available, training order was also compared (i.e. aerobic exercise before strength exercise vs. strength before aerobic exercise). Studies were divided into subgroups based on the description in the manuscript. This was particularly true for training status, with studies classified as “untrained” if participants were clearly described as “sedentary”, “previously untrained”, or “inactive”. Conversely, all other studies were classified as “active” (i.e. “recreationally active”, “trained”, “well-trained” etc.). Specific rationale for the exclusion of individual studies can be found at Online Supplemental Table S2.

2.5 Assessment of methodological quality

Risk of bias for the included studies was assessed using the Physiotherapy Evidence Database (PEDro) scale independently by two reviewers. The PEDro scale has previously been assessed as a valid measure of the methodological quality of randomised trials [28]. Studies scoring >6 were classified as “high-quality”, studies scoring 4–5 were classified as “medium-quality”, and studies scoring <4 were classified as “low-quality”. The following sources of bias were considered: Selection (sequence generation and allocation concealment), performance (blinding of participants/personnel), detection (blinding outcome assessors), attrition (incomplete outcome data), reporting (selective reporting), and other potential biases (e.g. recall bias). The risk of bias scores for the included studies are presented in Online Supplemental Table S3. The mean score for scale criteria 2–11 of the PEDro scale was 4.3/10, i.e., medium quality.

3. Results

3.1 Study characteristics

The database search identified 15,729 potentially eligible articles. After further screening and eligibility assessment, a total of 43 studies were included in the final analysis (Figure 1). The characteristics of the studies, participants, and training interventions are summarised in Supplemental Table S1. The meta-analysis included a total of 1,090 participants of which 590 participants performed supervised combined aerobic and strength training and 500 participants performed strength training alone. Of the included studies, cycling was the most common type of aerobic training (24 studies), followed by running (16 studies). In addition, the combination of running and cycling [9], rowing [29], and continuous repeated leg extensions [30] were each evaluated by one study.

3.2 Maximal strength

The final analysis included 37 studies [9–11, 29–62] with 525 participants performing combined aerobic and strength training and 442 participants performing strength training alone. The observed SMD ranged from -1.37 to 1.99, and the estimated average SMD based on the random-effects model was -0.06 (95% CI: -0.20, 0.09, $p = 0.446$), indicating no interference effect of aerobic training (Figure 2). The estimated log variability ratio based on the random-effects model was 0.05 (95% CI: -0.05, 0.15, $p = 0.311$). According to the Q-test, there was no significant heterogeneity in the true outcomes ($Q(36) = 32.591$, $p = 0.632$, $\hat{\tau}^2 = 0.000$, $I^2 = 0.00\%$). An examination of the studentised residuals showed that there was no evidence of outliers within this model and none of the studies were overly influential.

Subgroup analyses showed no statistical differences ($p > 0.05$) (Online Supplemental Figures S2–S7).

Figure 2. Forest plot of studies comparing differences in maximal strength. SMD = standardised mean difference; CI = confidence interval.

3.3 Explosive strength

The final analyses included 18 studies [11, 31, 34, 38, 39, 42, 49, 51–54, 56, 58–60, 62–64], with 270 participants performing combined aerobic and strength training and 208 participants performing strength training alone. The observed SMD ranged from -1.60 to 0.22., and the estimated mean SMD based on the random-effects model was -0.28 (95% CI: -0.48, -

0.08, $p = 0.007$), indicating an interference effect of aerobic training (Figure 3). The estimated log variability ratio based on the random-effects model was 0.04 (95% CI: -0.09, 0.18, $p = 0.533$). According to the Q-test there was no significant heterogeneity in the true outcomes ($Q(17) = 26.675$, $p = 0.068$, $\hat{\tau}^2 = 0.068$, $I^2 = 35.81\%$). The studentised residuals pointed out Mikkola et al. [31] as a potential outlier that may be overly influential. Sensitivity analyses revealed that excluding this study reduced the amount of observed heterogeneity to $I^2 = 0.00\%$ ($Q(16) = 13.860$, $p = 0.061$, $\hat{\tau}^2 = 0.061$).

Subgroup analyses showed no statistical differences ($p > 0.05$) (Online Supplemental Figures S8-S11). When studies were grouped by type of aerobic training, the SMD was significantly in favour of strength training for cycling -0.44 (95% CI: -0.86, -0.01, $p = 0.043$), but not for running (Supplemental Figure S8). However, after removing the overly influential study by Mikkola et al. [31], this effect was no longer observed (SMD: -0.27, 95% CI: -0.58, 0.04, $p = 0.086$). A similar effect was also seen for low concurrent training frequency, with an initial SMD of -0.45 (95% CI: -0.87, -0.02, $p = 0.039$) in favour of the strength training group (Online Supplemental Figure S9). After removing the study by Mikkola et al. [31] this was reduced to -0.25 (95% CI: -0.50, 0.01, $p = 0.059$). Conversely, when studies were grouped by training modality, a significant interference effect was observed for studies that performed concurrent training within the same session (≤ 20 minutes between aerobic and strength training) (SMD: -0.31, 95% CI: -0.62, -0.01, $p = 0.043$), but not when concurrent training was separated by at least 3 hours (Online Supplemental Figure S11).

Figure 3. Forest plot of studies comparing differences in explosive strength. SMD = standardised mean difference; CI = confidence interval.

3.4 Muscle hypertrophy

The final analyses included 15 studies [10, 11, 33, 45–47, 49, 54, 55, 59, 62, 65–68] with 201 participants performing combined aerobic and strength training and 188 participants performing strength training alone. The observed SMD in each trial ranged from -0.67 to 0.28, and the estimated mean SMD based on the random-effects model was -0.01 (95% CI: -0.16, 0.18, $p = 0.919$), indicating no interference effect of aerobic training (Figure 4). The estimated log variability ratio based on the random-effects model was 0.04 (95% CI: -0.11, 0.19, $p = 0.567$). According to the Q-test, there was no significant heterogeneity in the true outcomes ($Q(14) = 4.687$, $p = 0.990$, $\hat{\tau}^2 = 0.000$, $I^2 = 0.00\%$). An examination of the studentised residuals showed that there was no potential outlier within this model. According to the Cook's distances, no study could be considered overly influential. Subgroup analyses revealed no statistical differences ($p > 0.05$) (Online Supplemental Figures S12-S14).

Figure 4. Forest plot of studies comparing differences in muscle hypertrophy. SMD = standardised mean difference; CI = confidence interval.

4. Discussion

The aim of this study was to provide systematic and evidence-based appraisal on whether aerobic training interferes with adaptations to strength training in terms of muscle function (maximal and explosive strength) and whole-muscle hypertrophy. In addition, the impact of important mediating covariates such as type of aerobic training, training modality, exercise order, concurrent training frequency, age, and training status were assessed. The main finding was that concurrent aerobic and strength training did not interfere with the development of maximal strength and muscle hypertrophy compared with strength training alone. However, the

development of explosive strength was negatively affected by concurrent training. Our subgroup analysis showed that this negative affect was exacerbated when concurrent training was performed within the same session, compared with when aerobic and strength training were separated by at least 3 hours. No significant effects were found for other moderators, such as type of aerobic training (cycling vs. running), frequency of concurrent training (> 5 weekly sessions vs. < 5 weekly sessions), training status (untrained vs. active), and mean age (< 40 years vs. > 40 years).

An important goal of this meta-analysis was to provide evidence that can be translated into optimised and fine-tuned exercise recommendations for fitness and health purposes. While our results are generally consistent with those reported by Wilson et al. a decade ago [14], these authors considered anaerobic power measures such as Wingate performance as indicators of explosive strength. Since we intentionally included only direct measures of explosive strength (i.e. jump performance, isometric rate of force development and dynamic leg press power), our findings reinforce that concurrent aerobic and strength training can compromise strength qualities that require rapid neural activation.

The mechanism for compromised explosive but not maximal strength is interesting and requires further research. Our findings are supported by an early study showing that muscle hypertrophy and maximal strength were unaffected by concurrent training, while rate of force development was blunted, likely due to interference with rapid voluntary neural activation [10]. More specifically, while the maximal neural activation was not compromised, the increase in the integrated electromyographic signal (iEMG) during the first 500 ms was attenuated in the group performing both aerobic and strength training. Since the rate of recruitment and maximal discharge of motor neurons largely determine the maximal rate of force development [69], it appears that the rate of recruitment and discharge of motor units is particularly sensitive to the interference effect of aerobic training. It could be speculated that residual fatigue induced by aerobic training affects the corticospinal inputs received by the motor neurons before force is generated, which would subsequently compromise rapid force generation. The latter could potentially reduce the quality, but not the quantity, of strength training sessions performed concurrently with aerobic training, thereby potentially reducing the development of explosive strength, but not maximal strength or muscle hypertrophy. This, in turn, could have implications for programme design, as it is apparent that concurrently improving both cardiorespiratory fitness and rapid force production through rather generic exercise recommendations presents a physiological challenge.

Consistent with this, our subgroup analysis indicated that the magnitude of interference in explosive strength development was dependent on the programming of the exercise sessions, with significant interference observed when aerobic and strength training were performed within the same training session. Previous studies have indicated that neuromuscular interference may be more pronounced when strength training is immediately preceded by aerobic training both in young [70] and older individuals [71]. However, our pooled analysis does not provide evidence for an order-specific effect, but rather highlights that combining aerobic and strength training in close proximity attenuates adaptations in explosive strength regardless of exercise order. Apart from limitations in rapid neural drive [10], previous studies have also suggested that adaptations in pennation angle and fascicle length [54] or patella tendon cross-sectional area [72] could be possible mechanistic explanations for these findings.

The moderators, including frequency of concurrent training, type of training, age, and training status, did not significantly influence adaptations in maximal and explosive strength, nor muscle hypertrophy. Similarly, no significant effects were observed in our analysis of log variability, indicating that there were no within-group differences in variability after concurrent training compared with strength training alone. Our results differ from the recently published meta-

analysis that focused exclusively on the effect of training status on maximal strength during concurrent training [15]. In this study, the one-repetition maximum for leg press and squat was negatively affected by concurrent training in trained individuals, but not in moderately trained or untrained individuals compared to strength training alone. Moreover, their subgroup analysis suggested that the negative effect observed in trained individuals occurred only when aerobic and strength training were performed within the same training session. However, due to the lack of consistent reporting, we chose not to divide the active participants into moderately or well-trained athletes, which may have diluted potential significant effects. Furthermore, albeit the exact calculations of Petré et al. 2021 were not published [15], their analysis appears to differ from our approach. Apart from the smaller number of studies included (27 vs. 37 studies), studies consisting of multiple intervention groups with only one comparator were included multiple times in the same analysis, potentially inflating power [73]. Although the results did not reach statistical significance, our subgroup analysis for training status showed a similar direction for the SMD in trained versus untrained participants as reported by Petré et al. [15].

In previous concurrent training research, numerous studies have focused on the possible interference mechanisms related to muscle hypertrophy [74]. The rationale for these studies stems from rodent and cellular models indicating possible inhibition of mechanistic targeted of rapamycin (mTOR) signalling through activation of AMP-activated protein kinase (AMPK) following aerobic exercise [75–78]. However, subsequent human studies failed to confirm these findings when examining physiological mechanisms such as metabolic stress and AMPK activation [67, 79] or protein synthesis [80] following concurrent exercise. Based on our systematic review, this is not surprising as none of the identified studies reported a significant interference effect on muscle hypertrophy. While Wilson et al. concluded from their subgroup analysis that there was a negative relationship between the effect size for hypertrophy and both aerobic training frequency and duration [14], our results do not confirm these observations. There are several possible explanations for this inconsistency, apart from the obvious fact that our analysis was conducted almost a decade later and therefore included more studies. First, the inclusion criteria differed since Wilson et al. included fibre hypertrophy as an outcome parameter and also included studies without a strength training control group. Second, we conducted our analysis based on an inverse variance-weighted random-effects model in a pre-test post-test control group design [18], whereas Wilson et al. estimated the effect sizes of each individual group, resulting in a total of 72 effect sizes for muscle hypertrophy. The reported aerobic training duration and intensity were then correlated with effect sizes, potentially leading to significant positive correlations.

While the current meta-analysis provides updated and novel information, there are some limitations that should be acknowledged. First, it should be noted that the majority of the included studies were only classified as medium quality (mean PEDro score 4.3 ± 0.9), while a total of $n = 7$ studies were of low quality. However, it is important to note that all items related to blinding may not be possible to achieve in exercise trials. In addition, poor reporting quality may have biased the outcome of this ranking. Thus, more importance can possibly be given to the studentised residuals and the Cooks distance [24]. Furthermore, meta-analyses are generally limited to the information provided within the included individual studies. Even though authors were contacted with requests for additional information, the response rate was low. Therefore, to avoid speculation, we decided to include only clearly defined moderators. For example, aerobic exercise intensity was not included because the included studies did not provide consistent information. However, it is possible that aerobic exercise intensity may impact the compatibility of aerobic and strength training. A previous meta-analysis examined the effects of concurrent high-intensity interval training (HIIT) and strength training and reported that lower body strength development was compromised by concurrent training compared to strength training alone, even though the authors noted that a possible negative effect on lower

body strength may be ameliorated by the inclusion of running-based HIIT and longer intermodal rest periods [81]. This was further supported by a recent narrative review reporting that HIIT can minimise the risk of neuromuscular interference and that this effect is even more pronounced when HIIT is replaced with sprint-interval training [82]. However, it should be acknowledged that based on previous research, the overall health benefits of concurrent training, apart from muscle function and size, appear to be greater than those obtained with isolated training of either aerobic or strength training [83, 84], and that the overall risk of interference effects is rather low. Therefore, most individuals, including recreational athletes, can enjoy complementary benefits from incorporating both aerobic and strength training into their training programme.

5. Conclusion

In summary, this updated meta-analysis shows that concurrent aerobic and strength training does not interfere with the development of maximal strength and muscle hypertrophy compared with strength training alone. This appears to be independent of the type of aerobic training (cycling vs. running), frequency of concurrent training (> 5 weekly sessions vs. < 5 weekly sessions), training status (untrained vs. active), and mean age (< 40 years vs. > 40 years). However, the evidence of reduced development of explosive strength with concurrent training, particularly when aerobic and strength training are performed in the same session, suggests that practitioners who prioritize explosive strength may benefit from separating aerobic and strength training to achieve optimal adaptations.

6. References

- 1 Li R, Xia J, Zhang XI, et al. Associations of Muscle Mass and Strength with All-Cause Mortality among US Older Adults. *Med Sci Sports Exerc* 2018;50(3):458–67.
- 2 Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA* 2009;301(19):2024–35.
- 3 Hawley JA. Adaptations of skeletal muscle to prolonged, intense endurance training. *Clin Exp Pharmacol Physiol* 2002;29(3):218–22.
- 4 Tesch PA. Skeletal muscle adaptations consequent to long-term heavy resistance exercise. *Med Sci Sports Exerc* 1988;20(5 Suppl):S132-4.
- 5 Daly RM, Dalla Via J, Fyfe JJ, et al. Effects of exercise frequency and training volume on bone changes following a multi-component exercise intervention in middle aged and older men: Secondary analysis of an 18-month randomized controlled trial. *Bone* 2021;148:115944. doi:10.1016/j.bone.2021.115944 [published Online First: 6 April 2021].
- 6 Severinsen MCK, Pedersen BK. Muscle-Organ Crosstalk: The Emerging Roles of Myokines. *Endocr Rev* 2020;41(4).
- 7 Casas-Herrero A, Cadore EL, Zambom-Ferraresi F, et al. Functional Capacity, Muscle Fat Infiltration, Power Output, and Cognitive Impairment in Institutionalized Frail Oldest Old. *Rejuvenation Research* 2013;16(5):396–403.
- 8 Sherrington C, Fairhall NJ, Wallbank GK, et al. Exercise for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2019;1:CD012424. doi:10.1002/14651858.CD012424.pub2 [published Online First: 31 January 2019].
- 9 Hickson RC. Interference of strength development by simultaneously training for strength and endurance. *Eur J Appl Physiol Occup Physiol* 1980;45(2-3):255–63.
- 10 Häkkinen K, Alen M, Kraemer WJ, et al. Neuromuscular adaptations during concurrent strength and endurance training versus strength training. *Eur J Appl Physiol* 2003;89(1):42–52. doi:10.1007/s00421-002-0751-9 [published Online First: 14 December 2002].
- 11 McCarthy JP, Agre JC, Graf BK, et al. Compatibility of adaptive responses with combining strength and endurance training. *Med Sci Sports Exerc* 1995;27(3):429–36.
- 12 McCarthy JP, Pozniak MA, Agre JC. Neuromuscular adaptations to concurrent strength and endurance training. *Med Sci Sports Exerc* 2002;34(3):511–19.
- 13 Chtara M, Chaouachi A, Levin GT, et al. Effect of concurrent endurance and circuit resistance training sequence on muscular strength and power development. *J Strength Cond Res* 2008;22(4):1037–45.

- 14 Wilson JM, Marin PJ, Rhea MR, et al. Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J Strength Cond Res* 2012;26(8):2293–307.
- 15 Petré H, Hemmingsson E, Rosdahl H, et al. Development of Maximal Dynamic Strength During Concurrent Resistance and Endurance Training in Untrained, Moderately Trained, and Trained Individuals: A Systematic Review and Meta-analysis. *Sports Med* 2021. doi:10.1007/s40279-021-01426-9 [published Online First: 22 March 2021].
- 16 Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLoS Medicine* 2009;6(7):e1000100. <http://dx.doi.org/10.1371/journal.pmed.1000100>.
- 17 Drevon D, Fursa SR, Malcolm AL. Intercoder Reliability and Validity of WebPlotDigitizer in Extracting Graphed Data. *Behav Modif* 2017;41(2):323–39. doi:10.1177/0145445516673998 [published Online First: 22 October 2016].
- 18 Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software* 2010;36(3):1–48. <https://lirias.kuleuven.be/1059637?limo=0>.
- 19 Morris SB. Estimating Effect Sizes From Pretest-Posttest-Control Group Designs. *Organizational Research Methods* 2008;11(2):364–86.
- 20 Becker BJ. Synthesizing standardized mean-change measures. *British Journal of Mathematical and Statistical Psychology* 1988;41(2):257–78.
- 21 Viechtbauer W. Bias and Efficiency of Meta-Analytic Variance Estimators in the Random-Effects Model. *Journal of Educational and Behavioral Statistics* 2005;30(3):261–93.
- 22 Cochran WG. The Combination of Estimates from Different Experiments. *Biometrics* 1954;10(1):101.
- 23 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539–58.
- 24 Viechtbauer W, Cheung MW-L. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods* 2010;1(2):112–25. doi:10.1002/jrsm.11 [published Online First: 4 October 2010].
- 25 Begg CB, Mazumdar M. Operating Characteristics of a Rank Correlation Test for Publication Bias. *Biometrics* 1994;50(4):1088.
- 26 Rothstein HR, Sutton AJ, Borenstein M. Publication bias in meta-analysis: Prevention, assessment and adjustments. Chichester, West Sussex: Wiley 2005.
- 27 Higgins JP, Green S. Cochrane handbook for systematic reviews of interventions version 5.1.0.: Cochrane Collab. 2011.
- 28 Elkins MR, Herbert RD, Moseley AM, et al. Rating the quality of trials in systematic reviews of physical therapy interventions. *Cardiopulm Phys Ther J* 2010;21(3):20–26.

- 29 Gravelle BL, Blessing DL. Physiological Adaptation in Women Concurrently Training for Strength and Endurance. *J Strength Cond Res* 2000;14(1):5–13.
- 30 Jones TW, Howatson G, Russell M, et al. Performance and neuromuscular adaptations following differing ratios of concurrent strength and endurance training. *J Strength Cond Res* 2013;27(12):3342–51.
- 31 Mikkola J, Rusko H, Izquierdo M, et al. Neuromuscular and cardiovascular adaptations during concurrent strength and endurance training in untrained men. *Int J Sports Med* 2012;33(9):702–10.
<http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L366368740>.
- 32 Gettman LR, Ward P, Hagan RD. A comparison of combined running and weight training with circuit weight training. *Med Sci Sports Exerc* 1982;14(3):229–34.
- 33 Sale DG, MacDougall JD, Jacobs I, et al. Interaction between concurrent strength and endurance training. *Journal of Applied Physiology* 1990;68(1):260–70.
<http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L20075241>.
- 34 Hennessy LC, Watson AWS. The Interference Effects of Training for Strength and Endurance Simultaneously. *The Journal of Strength and Conditioning Research* 1994;8(1):12.
- 35 Kraemer WJ, Patton JF, Gordon SE, et al. Compatibility of high-intensity strength and endurance training on hormonal and skeletal muscle adaptations. *J Appl Physiol (1985)* 1995;78(3):976–89.
<http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L25098185> <http://dx.doi.org/10.1152/jappl.1995.78.3.976>.
- 36 Dolezal BA, Potteiger JA. Concurrent resistance and endurance training influence basal metabolic rate in nondieting individuals. *J Appl Physiol (1985)* 1998;85(2):695–700.
- 37 Bell GJ, Syrotuik D, Martin TP, et al. Effect of concurrent strength and endurance training on skeletal muscle properties and hormone concentrations in humans. *Eur J Appl Physiol* 2000;81(5):418–27.
- 38 Balabinis CP, Psarakis CH, Moukas M, et al. Early phase changes by concurrent endurance and strength training. *J Strength Cond Res* 2003;17(2):393–401.
- 39 Glowacki SP, Martin SE, Maurer A, et al. Effects of resistance, endurance, and concurrent exercise on training outcomes in men. *Med Sci Sports Exerc* 2004;36(12):2119–27.
- 40 Haykowsky M, McGavock J, Muhll IV, et al. Effect of Exercise Training on Peak Aerobic Power, Left Ventricular Morphology, and Muscle Strength in Healthy Older Women. *Journals of Gerontology Series A: Biological Sciences & Medical Sciences* 2005;60(3):307–11.
- 41 Cadore EL, Pinto RS, Lhullier FLR, et al. Physiological effects of concurrent training in elderly men. *Int J Sports Med* 2010;31(10):689–97.

- 42 Hendrickson NR, Sharp MA, Alemany JA, et al. Combined resistance and endurance training improves physical capacity and performance on tactical occupational tasks. *Eur J Appl Physiol* 2010;109(6):1197–208.
- 43 Silva RF, Cadore EL, Kothe G, et al. Concurrent training with different aerobic exercises. *Int J Sports Med* 2012;33(8):627–34. <http://www.thieme-connect.de/DOI/DOI?10.1055/s-0031-1299698>.
- 44 Libardi CA, Souza GV de, Cavaglieri CR, et al. Effect of resistance, endurance, and concurrent training on TNF- α , IL-6, and CRP. *Med Sci Sports Exerc* 2012;44(1):50–56.
- 45 Lundberg TR, Fernandez-Gonzalo R, Gustafsson T, et al. Aerobic exercise does not compromise muscle hypertrophy response to short-term resistance training. *J Appl Physiol (1985)* 2013;114(1):81–89.
- 46 Souza EO de, Tricoli V, Roschel H, et al. Molecular adaptations to concurrent training. *Int J Sports Med* 2013;34(3):207–13.
- 47 Cantrell GS, Schilling BK, Paquette MR, et al. Maximal strength, power, and aerobic endurance adaptations to concurrent strength and sprint interval training. *Eur J Appl Physiol* 2014;114(4):763–71.
- 48 Lundberg TR, Fernandez-Gonzalo R, Norrbom J, et al. Truncated splice variant PGC-1 α 4 is not associated with exercise-induced human muscle hypertrophy. *Acta Physiologica* 2014;212(2):142–51.
- 49 Fyfe JJ, Bartlett JD, Hanson ED, et al. Endurance training intensity does not mediate interference to maximal lower-body strength gain during short-term concurrent training. *Frontiers in Physiology* 2016;7(NOV).
- 50 Kazior Z, Willis SJ, Moberg M, et al. Endurance exercise enhances the effect of strength training on muscle fiber size and protein expression of akt and mTOR. *PLoS One* 2016;11(2).
- 51 Robineau J, Babault N, Piscione J, et al. Specific Training Effects of Concurrent Aerobic and Strength Exercises Depend on Recovery Duration. *J Strength Cond Res* 2016;30(3):672–83.
- 52 Laird RH, Elmer DJ, Barberio MD, et al. Evaluation of Performance Improvements After Either Resistance Training or Sprint Interval-Based Concurrent Training. *J Strength Cond Res* 2016;30(11):3057–65.
- 53 Terzis G, Spengos K, Methenitis S, et al. Early phase interference between low-intensity running and power training in moderately trained females. *Eur J Appl Physiol* 2016;116(5):1063–73.
- 54 Tsitkanou S, Spengos K, Stasinaki A-N, et al. Effects of high-intensity interval cycling performed after resistance training on muscle strength and hypertrophy. *Scand J Med Sci Sports* 2017;27(11):1317–27. doi:10.1111/sms.12751 [published Online First: 23 September 2016].
- 55 Osuka Y, Fujita S, Kitano N, et al. Effects of Aerobic and Resistance Training Combined with Fortified Milk on Muscle Mass, Muscle Strength, and Physical Performance in Older

- Adults: A Randomized Controlled Trial. *The journal of nutrition, health & aging* 2017;21(10):1349–57.
- 56 Robineau J, Lacombe M, Piscione J, et al. Concurrent Training in Rugby Sevens: Effects of High-Intensity Interval Exercises. *International Journal of Sports Physiology & Performance* 2017;12(3):336–44.
 - 57 Panissa VLG, Fukuda DH, Oliveira FP de, et al. Maximum Strength Development and Volume-Load during Concurrent High Intensity Intermittent Training Plus Strength or Strength-Only Training. *Journal of sports science & medicine* 2018;17(4):623–32.
 - 58 Spiliopoulou P, Zaras N, Methenitis S, et al. Effect of Concurrent Power Training and High-Intensity Interval Cycling on Muscle Morphology and Performance. *J Strength Cond Res* 2019.
 - 59 Lee MJ-C, Ballantyne JK, Chagolla J, et al. Order of same-day concurrent training influences some indices of power development, but not strength, lean mass, or aerobic fitness in healthy, moderately-active men after 9 weeks of training. *PLoS One* 2020;15(5):e0233134.
 - 60 Karavirta L, Häkkinen A, Sillanpää E, et al. Effects of combined endurance and strength training on muscle strength, power and hypertrophy in 40-67-year-old men. *Scandinavian Journal of Medicine and Science in Sports* 2011;21(3):402–11.
 - 61 Karavirta L. Cardiorespiratory, Neuromuscular and Cardiac Autonomic Adaptations to Combined Endurance and Strength Training in Ageing Men and Women 162 Studies in sports, physical education and health.
 - 62 Shamim B, Devlin BL, Timmins RG, et al. Adaptations to Concurrent Training in Combination with High Protein Availability: A Comparative Trial in Healthy, Recreationally Active Men. *Sports Med* 2018;48(12):2869–83.
 - 63 Jones TW, Howatson G, Russell M, et al. Performance and Endocrine Responses to Differing Ratios of Concurrent Strength and Endurance Training. *J Strength Cond Res* 2016;30(3):693–702.
 - 64 Kraemer WJ, Vescovi JD, Volek JS, et al. Effects of concurrent resistance and aerobic training on load-bearing performance and the Army physical fitness test. *Military Medicine* 2004;169(12):994–99.
 - 65 Sillanpää E, Häkkinen A, Nyman K, et al. Body composition and fitness during strength and/or endurance training in older men. *Med Sci Sports Exerc* 2008;40(5):950–58.
 - 66 Sillanpää E, Häkkinen A, Laaksonen DE, et al. Serum basal hormone concentrations, nutrition and physical fitness during strength and/or endurance training in 3964-year-old women. *Int J Sports Med* 2010;31(2):110–17.
 - 67 Lundberg TR, Fernandez-Gonzalo R, Tesch PA. Exercise-induced AMPK activation does not interfere with muscle hypertrophy in response to resistance training in men. *J Appl Physiol (1985)* 2014;116(6):611–20. doi:10.1152/jappphysiol.01082.2013 [published Online First: 9 January 2014].
 - 68 Timmins RG, Shamim B, Tofari PJ, et al. Differences in Lower Limb Strength and Structure After 12 Weeks of Resistance, Endurance, and Concurrent Training.

- International journal of sports physiology and performance* 2020;1–8. <https://www.embase.com/search/results?subaction=viewrecord&id=L631343334&from=export> <http://dx.doi.org/10.1123/ijsp.2019-0788>.
- 69 Del Vecchio A, Negro F, Holobar A, et al. You are as fast as your motor neurons: speed of recruitment and maximal discharge of motor neurons determine the maximal rate of force development in humans. *The Journal of physiology* 2019;597(9):2445–56. doi:10.1113/JP277396 [published Online First: 1 March 2019].
 - 70 Eklund D, Pulverenti T, Bankers S, et al. Neuromuscular adaptations to different modes of combined strength and endurance training. *Int J Sports Med* 2015;36(2):120–29. doi:10.1055/s-0034-1385883 [published Online First: 26 September 2014].
 - 71 Cadore EL, Izquierdo M, Alberton CL, et al. Strength prior to endurance intra-session exercise sequence optimizes neuromuscular and cardiovascular gains in elderly men. *Experimental gerontology* 2012;47(2). <https://pubmed.ncbi.nlm.nih.gov/22178632/>.
 - 72 Rønnestad BR, Hansen EA, Raastad T. Strength training affects tendon cross-sectional area and freely chosen cadence differently in noncyclists and well-trained cyclists. *J Strength Cond Res* 2012;26(1):158–66.
 - 73 Cheung MW-L. A Guide to Conducting a Meta-Analysis with Non-Independent Effect Sizes. *Neuropsychol Rev* 2019;29(4):387–96. doi:10.1007/s11065-019-09415-6 [published Online First: 24 August 2019].
 - 74 Fyfe JJ, Bishop DJ, Stepto NK. Interference between concurrent resistance and endurance exercise: molecular bases and the role of individual training variables. *Sports Med* 2014;44(6):743–62.
 - 75 Coffey VG, Hawley JA. Concurrent exercise training: do opposites distract? *The Journal of physiology* 2017;595(9). <https://pubmed.ncbi.nlm.nih.gov/27506998/>.
 - 76 Hawley JA. Molecular responses to strength and endurance training: are they incompatible? *Appl Physiol Nutr Metab* 2009;34(3):355–61.
 - 77 Atherton PJ, Babraj J, Smith K, et al. Selective activation of AMPK-PGC-1 α or PKB-TSC2-mTOR signaling can explain specific adaptive responses to endurance or resistance training-like electrical muscle stimulation. *FASEB J* 2005;19(7):786–88. doi:10.1096/fj.04-2179fje [published Online First: 16 February 2005].
 - 78 Thomson DM, Fick CA, Gordon SE. AMPK activation attenuates S6K1, 4E-BP1, and eEF2 signaling responses to high-frequency electrically stimulated skeletal muscle contractions. *J Appl Physiol (1985)* 2008;104(3):625–32. doi:10.1152/jappphysiol.00915.2007 [published Online First: 10 January 2008].
 - 79 Apró W, Moberg M, Hamilton DL, et al. Resistance exercise-induced S6K1 kinase activity is not inhibited in human skeletal muscle despite prior activation of AMPK by high-intensity interval cycling. *American journal of physiology. Endocrinology and metabolism* 2015;308(6).
 - 80 Carrithers JA, Carroll CC, Coker RH, et al. Concurrent exercise and muscle protein synthesis: implications for exercise countermeasures in space. *Aviat Space Environ Med* 2007;78(5):457–62.

- 81 Sabag A, Najafi A, Michael S, et al. The compatibility of concurrent high intensity interval training and resistance training for muscular strength and hypertrophy: a systematic review and meta-analysis. *J Sports Sci* 2018;36(21):2472–83. doi:10.1080/02640414.2018.1464636 [published Online First: 16 April 2018].
- 82 Vechin FC, Conceição MS, Telles GD, et al. Interference Phenomenon with Concurrent Strength and High-Intensity Interval Training-Based Aerobic Training: An Updated Model. *Sports Med* 2021;51(4):599–605. doi:10.1007/s40279-020-01421-6 [published Online First: 6 January 2021].
- 83 Schroeder EC, Franke WD, Sharp RL, et al. Comparative effectiveness of aerobic, resistance, and combined training on cardiovascular disease risk factors: A randomized controlled trial. *PLoS One* 2019;14(1):e0210292. doi:10.1371/journal.pone.0210292 [published Online First: 7 January 2019].
- 84 Timmons JF, Minnock D, Hone M, et al. Comparison of time-matched aerobic, resistance, or concurrent exercise training in older adults. *Scand J Med Sci Sports* 2018;28(11). <https://pubmed.ncbi.nlm.nih.gov/29947107/>.