ANK1 is differentially expressed in the brains of patients with Parkinson's Disease.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY 11730

Parkinson's Disease (PD) is characterized by loss of dopaminergic neurons in the substantia nigra of the basal ganglia¹. We mined published microarray datasets^{2,3} to identify genes whose expression was most different in the substantia nigra of patients with PD as compared to that of non-affected patients. We identified significant changes in expression of the gene encoding ANK1 in the substantia nigra of patients with PD.

Keyword: Parkinson's Disease, motor disorders, substantia nigra, dopaminergic neurons, ANK1, ankyrin.

1   Parkinson's Disease is a neurodegenerative disorder with no cure. Patients with
2 Parkinson's Disease suffer from motor symptoms including rigidity, tremors, bradykinesia
3 (slowness of movement), and postural instability¹. With disease progression, non-motor
4 neuropsychiatric problems can emerge and worsen, including difficulties with cognition,
5 problems with executive functioning, and problems with mood and behavior¹. 40% of patients
6 with PD develop depression⁴ and approximately 50% of patients with PD will develop psychotic
7 symptoms - hallucinations⁵. Thus, motor and non-motor symptoms are a serious problem in
8 patient with PD. While treatment with dopamine precursor levodopa can provide significant
9 relief, such treatment is marked by "on" and "off" periods of effectiveness and the risk of
10 development of dyskinesias^{6,7}. Newer dopamine agonists such as pramipexole and ropinirole are
11 of limited efficacy⁸ and associated with the risk of impulse control disorders⁹. The development
12 of targeted treatments for patients with Parkinson's Disease, to halt disease progression in the
13 substantia nigra and control motor and neuropsychiatric symptoms in PD patients is required.

14
15
16
17
18   We mined published microarray datasets to identify the most striking changes in gene
19 expression in the substantia nigra of patients with Parkinson's Disease. We identified ANK1 as
20 among the most differentially expressed genes in the brains of patients with PD. ANK1 may be
21 of relevance to the biology of PD.

22 **Methods**

23
24
25   We used datasets GSE49036 and GSE8397 for this differential gene expression analysis
26 of the substantia nigra in Parkinson's Disease in conjunction with GEO2R. GSE49036 was
27
28

generated with Affymetrix Human Genome U133 Plus 2.0 Array technology with $n=8$ for substantia nigra from control subjects and $n=15$ for substantia nigra from patients with Parkinson's Disease. GSE8397 was generated with Affymetrix Human Genome U133A Array technology with $n=11$ for substantia nigra from control subjects and $n=26$ for substantia nigra from patients with Parkinson's Disease. The Benjamini and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether ANK1 expression was significantly different when comparing substantia nigra from control subjects and from patients with Parkinson's Disease using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses (Version 8.4.0)(455).

Results

We mined published microarray datasets^{2,3} to identify differentially expressed genes in patients with Parkinson's Disease.

ANK1 is differentially expressed in the Parkinson's Disease substantia nigra

We identified ANK1 as among the genes whose expression changed most significantly genome-wide when comparing the transcriptional profiles of the substantia nigra from control, non-affected subjects and from patients with Parkinson's Disease (PD)². When sorting each of

1 the transcripts identified by microarray based on degree of change in expression between control
2 subjects and patients with PD, ANK1 ranked 47 out of 54675 total transcripts (Table 1).

3
4 Differential expression of ANK1 in the substantia nigra of patients with PD was statistically
5 significant (Table 1; $p=0.00004525$).

6
7 We mined a second microarray dataset³ to determine if differential expression of ANK1
8 in patients with PD could be observed. ANK1 was again among the genes most differentially
9 expressed when comparing global transcriptional profiles of the substantia nigra in patients with
10 PD with that of control subjects (Table 2). When sorting each of the transcripts identified by
11 microarray based on degree of change in expression between control subjects and patients with
12 PD, ANK1 ranked 3 out of 22283 total transcripts (Table 2). Differential expression of ANK1 in
13 the substantia nigra of patients with PD in this second dataset was statistically significant (Table
14 2; $p=1.77E-09$).

15 ANK1 is expressed at significantly lower levels in patients with Parkinson's Disease.

16
17
18
19 We obtained exact mRNA expression levels for the differentially expressed ANK1
20 transcript, from control subjects and from patients with PD, to directly compare ANK1
21 expression levels. We also performed a statistical test to determine if this difference in
22 expression was statistically significant. ANK1 was expressed at lower levels in the substantia
23 nigra of patients with PD, and this difference was statistically significant (Figure 1; $p=0.0006$).
24
25 We calculated a mean fold change of 0.9056 ± 0.0424 in ANK1 expression when comparing
26 substantia nigra from patients with PD to that of control subjects (Table 1).
27
28

1 In the second microarray dataset queried, we again observed increased expression of
2 ANK1 in the substantia nigra of patients with PD; this difference was statistically significant
3 (Figure 2; $p < 0.0001$). We calculated a mean fold change of 0.7350 ± 0.1000 in ANK1
4 expression when comparing substantia nigra from patients with PD to that of control subjects
5 (Table 2).
6

7
8 Thus, we identified ANK1 as a differentially expressed gene in the substantia nigra of
9 patients with Parkinson's Disease and found that ANK1 was expressed at significantly lower
10 levels in the substantia nigra of patients with PD.
11

12 Discussion

13
14 We mined published microarray data to discover genes associated with Parkinson's
15 Disease in tissues classically associated with the disease, the substantia nigra of the basal
16 ganglia. We identified ANK1 as among the most differentially expressed genes in the substantia
17 nigra of patients with Parkinson's Disease.
18

19 ANK1 encodes for ankyrin¹⁰, a structural protein linking integral membrane proteins at
20 the plasma membrane to the cytoskeleton; the ankyrin repeat is a 33 amino acid motif with an L-
21 shaped structure and β -hairpin-helix-loop-helix that mediate interaction with multiple
22 cytoskeletal proteins¹¹. Ankyrin can interact with and has unique domains for interaction with
23 band-3 and spectrin or fodrin (non-erythroid spectrin)^{12,13}. Other integral membrane protein
24 tethered to the cytoskeleton through ankyrin is the Na,K-ATPase, where ankyrin, the Na,K-
25 ATPase and spectral can form a complex¹⁴. In erythrocytes, α - and β -spectrin heterodimers are
26
27
28

linked to the

plasma membrane by interactions with multiple proteins including band-3 and ankyrin¹⁵.

Ankyrin is important for integrity of the erythrocyte membrane; approximately 50% of cases of the inherited hemolytic anemia hereditary spherocytosis are caused by perturbed expression of ankyrin^{16, 17}. Interestingly, mice homozygous for the nb mutation on Chromosome 8, nb/nb mice, are deficient in ankyrin, and have severe hemolytic anemia¹⁸. Of significance, nb/nb mice develop a psychomotor disorder in adulthood at 6 months and in Purkinje cells of the cerebellum are significantly reduced concomitant with the psychomotor disorder¹⁸.

ANK1 is prominently implicated in at least one human disease of the brain. In Alzheimer's Disease, ANK1 expression is significantly increased in brain microglia, and the ANK1 locus is differentially methylated in association with Alzheimer's Disease pathology in the entorhinal cortex¹⁹. This region of ANK1 was differentially hyper-methylated in the superior temporal gyrus and the cerebral cortex but not in the cerebellum, where AD pathology is generally lacking²⁰. A separate study of 708 autopsied brains found significant association of CpG methylation at the ANK1 locus with the risk of development of Alzheimer's Disease²¹. Importantly, a study using bisulfite pyrosequencing to measure DNA methylation at 8 CpG sites within an 118 basepair region of the ANK1 gene found ANK1 hypermethylation in the entorhinal cortex of patients with Parkinson's Disease, as well as in patients with Alzheimer's Disease and Huntington's Disease²². Outside of the brain, SNPs at the ANK1 locus are also associated with risk of development of TIID in the Japanese population²³.

1   If validated in separate datasets generated with tissues from patients with Parkinson's
2 Disease, the effect of ANK1 depletion and over-expression *in vivo* in the substantia nigra
3 can be evaluated through conditional deletion or expression in wild-type mice or in mouse
4 models of Parkinson's Disease²⁴⁻²⁷. ANK1 may be relevant to the genesis, maintenance or
5 progression of Parkinson's Disease.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 =

References

1. Dauer, W. and Przedborski, S., 2003. Parkinson's disease: mechanisms and models. *Neuron*, 39(6), pp.889-909.
2. Dijkstra, A.A., Ingrassia, A., de Menezes, R.X., van Kesteren, R.E., Rozemuller, A.J., Heutink, P. and van de Berg, W.D., 2015. Evidence for immune response, axonal dysfunction and reduced endocytosis in the substantia nigra in early stage Parkinson's disease. *PloS one*, 10(6), p.e0128651.
3. Moran, L.B., Duke, D.C., Deprez, M., Dexter, D.T., Pearce, R.K.B. and Graeber, M.B., 2006. Whole genome expression profiling of the medial and lateral substantia nigra in Parkinson's disease. *Neurogenetics*, 7(1), pp.1-11.
4. Cummings, J.L., 1992. Depression and Parkinson's disease: a review. *The American journal of psychiatry*.
5. Fénelon, G. and Alves, G., 2010. Epidemiology of psychosis in Parkinson's disease. *Journal of the neurological sciences*, 289(1-2), pp.12-17.
6. Parkinson Study Group, 2004. Levodopa and the progression of Parkinson's disease. *New England Journal of Medicine*, 351(24), pp.2498-2508.
7. Lees, A.J., 1989. The on-off phenomenon. *Journal of Neurology, Neurosurgery & Psychiatry*, 52(Suppl), pp.29-37.
8. Stowe, R., Ives, N., Clarke, C.E., Ferreira, J., Hawker, R.J., Shah, L., Wheatley, K. and Gray, R., 2008. Dopamine agonist therapy in early Parkinson's disease. *Cochrane Database of Systematic Reviews*, (2).
9. Weintraub, D., Siderowf, A.D., Potenza, M.N., Goveas, J., Morales, K.H., Duda, J.E., Moberg, P.J. and Stern, M.B., 2006. Association of dopamine agonist use with impulse control disorders in Parkinson disease. *Archives of neurology*, 63(7), pp.969-973.
10. Polymeropoulos, M.H., Rath, D.S., Xiao, H. and Merril, C.R., 1991. Dinucleotide repeat polymorphism at the human ankyrin gene (ANK1). *Nucleic acids research*, 19(4), p.969.
11. Sedgwick, S.G. and Smerdon, S.J., 1999. The ankyrin repeat: a diversity of interactions on a common structural framework. *Trends in biochemical sciences*, 24(8), pp.311-316.

12. Bennett, V. and Stenbuck, P.J., 1979. The membrane attachment protein for spectrin is associated with band 3 in human erythrocyte membranes. *Nature*, 280(5722), pp.468-473.
13. Drenckhahn, D., Schluter, K., Allen, D.P. and Bennett, V., 1985. Colocalization of band 3 with ankyrin and spectrin at the basal membrane of intercalated cells in the rat kidney. *Science*, 230(4731), pp.1287-1289.
14. Morrow, J.S., Cianci, C.D., Ardito, T., Mann, A.S. and Kashgarian, M., 1989. Ankyrin links fodrin to the alpha subunit of Na, K-ATPase in Madin-Darby canine kidney cells and in intact renal tubule cells. *Journal of Cell Biology*, 108(2), pp.455-465.
15. Devarajan, P., Scaramuzzino, D.A. and Morrow, J.S., 1994. Ankyrin binds to two distinct cytoplasmic domains of Na, K-ATPase alpha subunit. *Proceedings of the National Academy of Sciences*, 91(8), pp.2965-2969.
16. del Giudice, E.M., Francese, M., Nobili, B., Morlé, L., Cutillo, S., Delaunay, J. and Perrotta, S., 1998. High frequency of de novo mutations in ankyrin gene (ANK1) in children with hereditary spherocytosis. *The Journal of pediatrics*, 132(1), pp.117-120.
17. Randon, J., MIRAGLIA DEL GIUDICE, E., Bozon, M., PERROTTA, S., De Vivo, M., Iolascon, A., Delaunay, J. and Morle, L., 1997. Frequent de novo mutations of the ANK1 gene mimic a recessive mode of transmission in hereditary spherocytosis: three new ANK1 variants: ankyrins Bari, Napoli II and Anzio. *British journal of haematology*, 96(3), pp.500-506.
18. Peters, L.L., Birkenmeier, C.S., Bronson, R.T., White, R.A., Lux, S.E., Otto, E., Bennett, V., Higgins, A. and Barker, J.E., 1991. Purkinje cell degeneration associated with erythroid ankyrin deficiency in nb/nb mice. *The Journal of cell biology*, 114(6), pp.1233-1241.
19. Mastroeni, D., Sekar, S., Nolz, J., Delvaux, E., Lunnon, K., Mill, J., Liang, W.S. and Coleman, P.D., 2017. ANK1 is up-regulated in laser captured microglia in Alzheimer's brain: the importance of addressing cellular heterogeneity. *PloS one*, 12(7), p.e0177814.
20. Lunnon, K., Smith, R., Hannon, E., De Jager, P.L., Srivastava, G., Volta, M., Troakes, C., Al-Sarraj, S., Burrage, J., Macdonald, R. and Condliffe, D., 2014. Methylomic profiling implicates cortical deregulation of ANK1 in Alzheimer's disease. *Nature neuroscience*, 17(9), pp.1164-1170.

21. De Jager, P.L., Srivastava, G., Lunnon, K., Burgess, J., Schalkwyk, L.C., Yu, L., Eaton, M.L., Keenan, B.T., Ernst, J., McCabe, C. and Tang, A., 2014. Alzheimer's disease: early alterations in brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. *Nature neuroscience*, 17(9), pp.1156-1163.
22. Smith, A.R., Smith, R.G., Burrage, J., Troakes, C., Al-Sarraj, S., Kalaria, R.N., Sloan, C., Robinson, A.C., Mill, J. and Lunnon, K., 2019. A cross-brain regions study of ANK1 DNA methylation in different neurodegenerative diseases. *Neurobiology of aging*, 74, pp.70-76.
23. Imamura, M., Maeda, S., Yamauchi, T., Hara, K., Yasuda, K., Morizono, T., Takahashi, A., Horikoshi, M., Nakamura, M., Fujita, H. and Tsunoda, T., 2012. A single-nucleotide polymorphism in ANK1 is associated with susceptibility to type 2 diabetes in Japanese populations. *Human molecular genetics*, 21(13), pp.3042-3049.
24. Fredriksson, A. and Archer, T., 1994. MPTP-induced behavioural and biochemical deficits: a parametric analysis. *Journal of neural transmission-Parkinson's disease and dementia section*, 7(2), pp.123-132.
25. Yavich, L., Oksman, M., Tanila, H., Kerokoski, P., Hiltunen, M., van Groen, T., Puoliväli, J., Männistö, P.T., García-Horsman, A., MacDonald, E. and Beyreuther, K., 2005. Locomotor activity and evoked dopamine release are reduced in mice overexpressing A30P-mutated human α -synuclein. *Neurobiology of disease*, 20(2), pp.303-313.
26. Li, Y., Liu, W., Oo, T.F., Wang, L., Tang, Y., Jackson-Lewis, V., Zhou, C., Geghman, K., Bogdanov, M., Przedborski, S. and Beal, M.F., 2009. Mutant LRRK2 R1441G BAC transgenic mice recapitulate cardinal features of Parkinson's disease. *Nature neuroscience*, 12(7), pp.826-828.
27. Brooks, S.P. and Dunnett, S.B., 2009. Tests to assess motor phenotype in mice: a user's guide. *Nature Reviews Neuroscience*, 10(7), pp.519-529.

Rank	ID	p-value	t	B	FC	Gene	Gene name
47	240363_at	0.00004525	-4.9399796	2.05896	0.9056 ± 0.0424	ANK1	ankyrin 1

Table 1: ANK1 is differentially expressed in the Parkinson's Disease substantia nigra.

The rank of differential expression, the probe/transcript ID, the *p*-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the fold change of ANK1 in PD substantia nigra when compared to control substantia nigra, the gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	FC	Gene	Gene name
3	207087_x_at	1.77E-09	-7.9426279	11.471113	0.7350 ± 0.1000	ANK1	ankyrin 1

Table 2: ANK1 is differentially expressed in the Parkinson's Disease substantia nigra.

The rank of differential expression, the probe/transcript ID, the *p*-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the fold change of ANK1 in PD substantia nigra when compared to control substantia nigra, the gene and gene name are listed in this chart.

Figure 1: ANK1 is expressed at significantly lower levels in the substantia nigra of patients with Parkinson's Disease.

The mRNA expression of ANK1 in normal substantia nigra from control subjects (left) and in the substantia nigra of patients with PD (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in ANK1 expression between the normal substantia nigra and substantia nigra from patients with PD, a *p*-value, listed above.

Figure 2: ANK1 is expressed at significantly lower levels in the substantia nigra of patients with Parkinson's Disease.

The mRNA expression of ANK1 in normal substantia nigra from control subjects (left) and in the substantia nigra of patients with PD (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in ANK1 expression between control substantia nigra and substantia nigra from patients with PD, a *p*-value, listed above.