Over-expression of ribosomal protein S6 kinase A1 in human endometrial cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Gynecologic cancers including cancers of the endometrium are a clinical problem¹⁻⁴. We mined published microarray data^{5,6} to discover genes associated with endometrial cancers by comparing transcriptomes of the normal endometrium and endometrial tumors from humans. We identified ribosomal protein S6 kinase A1, encoded by RPS6KA1, as among the most differentially expressed genes, transcriptome-wide, in cancers of the endometrium. RPS6KA1 was expressed at significantly higher levels in endometrial tumor tissues as compared to the endometrium. Importantly, in human endometrial cancer, primary tumor expression of RPS6KA1 was correlated with recurrence-free survival in white patients with low mutational burden. RPS6KA1 may be a molecule of interest in understanding the etiology or progression of human endometrial cancer.

Keywords: endometrial cancer, gynecologic cancers, endometrium, RPS6KA1, ribosomal protein S6 kinase A1, systems biology of endometrial cancer, targeted therapeutics in endometrial cancer.

Endometrial cancer is the most common gynecologic cancer in the developed world¹. Over the last three decades, the incidence of endometrial cancer has increased 21%⁴ and the death rate has increased 100%³. We harnessed the power of independently published microarray datasets^{5,6} to determine in an unbiased fashion and at the systems-level genes most differentially expressed in endometrial tumors. We report here the differential and increased expression of the ribosomal protein S6 kinase A1 (RPS6KA1) in human endometrial cancer.

Methods

We utilized datasets GSE63678⁵ and GSE115810⁶ for this global differential gene expression analysis of human endometrial cancer in conjunction with GEO2R. GSE63678 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=5$ control endometrial tissues (including $n=4$ uterine myomas and $n=1$ benign cyst) and $n=7$ endometrial cancers (including $n=2$ endometrial adenocarcinomas, $n=3$ mixed endometrioid adenocarcinomas, and $n=2$ adenocarcinomas with squamous differentiation); analysis was performed using platform GPL571. GSE115810 was generated using Affymetrix Human Genome U133A Array technology with $n=3$ control endometrial tissues and $n=24$ endometrial cancers; analysis was performed using platform GPL96. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether RPS6KA1 gene expression was significantly different between control endometrial tissue and endometrial tumor tissue in humans using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter tool⁷ for correlation of RPS6KA1 mRNA expression levels with recurrence-free survival in $n=543$ endometrial cancer patients.

Results

We harnessed the power of blind comparative transcriptome analysis using published microarray data^{5,6} to discover in an unbiased fashion genes associated with endometrial cancer in humans.

RPS6KA1 is differentially expressed in endometrial cancer.

We identified ribosomal protein S6 kinase A1, encoded by RPS6KA1, as among the genes most differentially expressed in cancers of the endometrium when compared to benign endometrial tissues (Chart 1). When sorting each of the genes expressed in endometrial tumor tissue based on significance of change in expression as compared to benign endometrial tissue, RPS6KA1 ranked 99 out of 22273 transcripts, equating to 99.6% differential expression (Chart 1). Differential expression of RPS6KA1 in human endometrial cancers was statistically significant (Chart 1; $p=8.96E-05$).

We queried a second microarray data to validate differential expression of RPS6KA1 in endometrial cancer. Again, we observed differential expression of RPS6KA1 when comparing endometrial tumor tissue to benign endometrial tissue (Chart 2). When sorting each of the genes expressed in endometrial tumor tissue based on significance of change in expression as compared to benign endometrial tissue, RPS6KA1 ranked 1316 out of 22283 transcripts, equating to 94.1% differential expression (Chart 2). Differential expression of RPS6KA1 in human endometrial cancers was statistically significant (Chart 2; $p=0.0179678$).

RPS6KA1 is expressed at significantly higher levels in endometrial cancers as compared to benign endometrial tissue.

We obtained exact mRNA expression levels for RPS6KA1 in endometrial tumor tissues and from benign endometrial tissue to evaluate direction and statistical significance of change in expression of RPS6KA1 in human endometrial cancer. RPS6KA1 was expressed at higher levels in endometrial tissue as compared to normal endometrial tissue, and this difference was statistically significant (Figure 1; $p<0.0001$). We calculated a mean fold change of 1.27 in RPS6KA1 mRNA levels in human endometrial cancer, as RPS6KA1 was expressed at 6.47 ± 0.76 arbitrary units (A.U.) in control endometrial tissue but at 8.20 ± 0.35 A.U. in endometrial tumor tissue.

RPS6KA1 expression is correlated with patient survival outcomes in endometrial cancer.

We performed Kaplan-Meier survival analysis to evaluate correlation between RPS6KA1 primary tumor expression and survival outcomes in 543 patients with endometrial cancer. We observed a correlation between primary tumor expression of RPS6KA1 and recurrence-free survival in patients with endometrial cancer, in white patients with low mutational burden (Figure 2). RPS6KA1 primary tumor mRNA levels were a positive prognostic indicator in white endometrial cancer patients with low mutational burden. White patients with low mutational burden whose primary tumors expressed low levels of RPS6KA1 possessed median RFS of 78.4 months, while white patients with low mutational burden whose tumors expressed high levels of RPS6KA1 possessed median RFS of 112.17 months. This difference in RFS based on RPS6KA1 tumor expression in white patients with endometrial cancer with low mutational burden approached statistical significance (Figure 2, Chart 3; logrank p -value: 0.057; hazard ratio: 0.52 (0.26-1.03)). RPS6KA1 primary endometrial tumor expression was not correlated with recurrence-free survival in white patients with high mutational burden (Figure 2, Chart 3; logrank p -value: 0.16; hazard ratio: 0.56 (0.25-1.27)), nor in black patients with high ((Figure 2, Chart 3; logrank p -value: 0.28; hazard ratio: 0.4 (0.07-2.22)) or low mutational burden (Figure 2, Chart 3; logrank p -value: 0.17; hazard ratio: 0.46 (0.15-1.42)).

Thus, by mining published microarray data^{5,6} in an unbiased and systematic fashion, we identified ribosomal protein S6 kinase A1, encoded by RPS6KA1, as among the genes whose

1 expression was most different, transcriptome-wide, in the endometrial tumor tissue of patients
2 with endometrial cancer when compared to benign endometrial tissue; we observed
3 significantly increased expression of RPS6KA1 in endometrial tumor tissue as compared to
4 benign endometrial tissue. Further, we found a correlation between RPS6KA1 expression and
5 patient survival outcomes in human endometrial cancer, as recurrence-free survival was superior
6 in patients whose tumors expressed higher levels of RPS6KA1 as compared to patients whose
tumors expressed lower levels of RPS6KA1, in white patients with low mutational burden, but
not in white patients with high mutational burden, nor in black patients with high or low
mutational burden.

7 **Discussion**

8
9 We provided evidence here that ribosomal protein S6 kinase A1 is among the genes most
10 differentially expressed in human endometrial cancer, that mRNA for RPS6KA1 is present at
11 significantly increased quantity in endometrial tumor tissue as compared to benign endometrium,
12 and that RPS6KA1 primary tumor expression is correlated with recurrence-free survival in white
13 endometrial cancer patients with low mutational burden. These data suggest RPS6KA1 may be
14 of importance to fundamental biological processes that underlie the initiation, progression or
15 maintenance of human endometrial cancer.
16
17
18
19
20
21
22
23
24
25
26
27
28

References

1. Amant, F., Moerman, P., Neven, P., Timmerman, D., Van Limbergen, E. and Vergote, I., 2005. Endometrial cancer. *The Lancet*, 366(9484), pp.491-505.
2. Sorosky, J.I., 2008. Endometrial cancer. *Obstetrics & Gynecology*, 111(2), pp.436-447.
3. Morice, P., Leary, A., Creutzberg, C., Abu-Rustum, N. and Darai, E., 2016. Endometrial cancer. *The Lancet*, 387(10023), pp.1094-1108.
4. Sorosky, J.I., 2012. Endometrial cancer. *Obstetrics & Gynecology*, 120(2 Part 1), pp.383-397.
5. Pappa, K.I., Polyzos, A., Jacob-Hirsch, J., Amariglio, N., Vlachos, G.D., Loutradis, D. and Anagnou, N.P., 2015. Profiling of discrete gynecological cancers reveals novel transcriptional modules and common features shared by other cancer types and embryonic stem cells. *PLoS One*, 10(11), p.e0142229.
6. Hermyt, E., Zmarzły, N., Grabarek, B., Kruszniewska-Rajs, C., Gola, J., Jęda-Golonka, A., Szczepanek, K., Mazurek, U. and Witek, A., 2019. Interplay between miRNAs and Genes Associated with Cell Proliferation in Endometrial Cancer. *International journal of molecular sciences*, 20(23), p.6011.
7. Nagy, A., Munkacsy, G. and Gyorffy, B., 2020. Pancancer survival analysis of cancer hallmark genes. *bioRxiv*.

Rank: 99
Probe ID: 203379_at
p-value: 8.96E-05
t: -5.7306728
B: 1.74552
Gene: RPS6KA1
Gene name: ribosomal protein S6 kinase A1

Chart 1: RPS6KA1 is differentially expressed in endometrial cancer when comparing primary endometrial tumors to benign endometrial tissue.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 1316
Probe ID: 203379_at
p-value: 0.0179678
t: -2.513721
B: -3.13642
Gene: RPS6KA1
Gene name: ribosomal protein S6 kinase A1

Chart 2: RPS6KA1 is differentially expressed in endometrial cancer when comparing primary endometrial tumors to benign endometrial tissue.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: RPS6KA1 is expressed at significantly higher levels in the endometrial tumors of patients with endometrial cancer when compared to benign endometrium.

The mRNA expression level of RPS6KA1 in benign endometrial tissue (left) and in primary tumors of the endometrium (right) is graphically depicted; the result of a statistical test evaluating significance of difference in RPS6KA1 expression between benign endometrial tissue and primary tumors of the endometrial tissue is $p < 0.0001$.

Figure 2: Correlation between RPS6KA1 primary tumor expression and recurrence-free survival in endometrial cancer, in white patients with low mutational burden.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=543$ total endometrial cancer patients stratified into two groups, based on low or high expression of RPS6KA1 in patient primary tumors, in black patients with high mutational burden (top left), black patients with low mutational burden (top right), white patients with high mutational burden (bottom left), and white patients with low mutational burden (bottom right). The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on RPS6KA1 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low RPS6KA1 expression: 78.4 months

High RPS6KA1 expression: 112.17 months

Chart 3: Median recurrence-free survival is superior in endometrial cancer patients with high primary tumor expression of RPS6KA1, in white patients with low mutational burden.

The median RFS (recurrence-free survival) of white endometrial cancer patients with low mutational burden, with low primary tumor expression of RPS6KA1 and high primary tumor expression of RPS6KA1 is listed in this chart.