FBXL8 is differentially expressed and transcriptionally induced in MERS coronavirus infection.

Shahan Mamoor, MS¹

¹Thomas Jefferson School of Law

San Diego, CA 92101

mamoorsk@tjssl.edu

The coronavirus COVID19 pandemic is an emerging biosafety threat to the nation and the world (1). There are no treatments approved for coronavirus infection in humans (2) and there is a lack of information available regarding the basic transcriptional behavior of human cells and mammalian tissues following coronavirus infection. We mined two independent datasets (3, 4), public (3) and published (4) containing transcriptome data from infection models of the Middle East respiratory syndrome (MERS) coronavirus to discover genes that are differentially expressed in human cells after infection with MERS-CoV coronaviruses and assess their differential expression after infection with other coronavirus types. We identified the F-box and Leucine Rich Region (LRR) protein 8 (FBXL8) as a differentially expressed gene following infection of primary human cells and a human cell line with wild-type MERS-CoV and MERS-CoV London, respectively. FBXL8 induction after viral infection appears to be unique to MERS-CoV among coronaviruses studied.

Keywords: FBXL8, coronavirus, MERS coronavirus, systems biology of viral infection.

Viruses are classified according to a system known as the “Baltimore” classification of viruses (5) wherein the characteristics of the viral genome, including whether it is plus or minus strand, also known as positive-sense or negative-sense, whether the genome is single-stranded or double-stranded, whether their genome is composed of RNA or DNA, and whether or not they use a reverse transcriptase to replicate are used to group viruses. Coronaviruses are plus-strand RNA viruses that contain an envelope surrounding their viral particle (6). Their genome is largest of all RNA viruses, ranging from 27 to 33 kb in size (7). They obtain their name from the crown-like appearance of the viral particle imparted by the structure of the large-surface glycoprotein (7). The coronaviridae family includes seven viruses capable of infecting humans, including the severe acute respiratory distress syndrome, or SARS coronavirus (8), the Middle East respiratory syndrome coronavirus, or MERS coronavirus (9), the human coronaviruses (HCoV) 229E, OC43, HKU and NL63 (10-12), and the novel human coronavirus now known as COVID19 (13, 14). As of March 19, 2020, the World Health Organization reported 209,839 cases of COVID19 and 8778 deaths from COVID19 infection world-wide (15). There are no FDA-approved treatments for human coronavirus infection.

We used a systems-level approach to identify the genes whose expression changes most significantly following infection of human cells with Middle East Respiratory Syndrome Coronavirus (MERS-CoV) using two independent datasets (3, 4). From these datasets were mined transcriptome data following coronavirus infection in human cell culture and in primary human endothelial cells. Across both of these datasets, we identified the F-box and Leucine Rich Region (LRR) protein 8, FBXL8 as among the genes most differentially expressed following MERS coronavirus infection. FBXL8 represents a transcriptional target of the host cell gene expression program following infection with MERS coronaviruses in human cells.

Methods

We used datasets GSE100509 (3) and GSE56677 (4) for this systems-level differential gene expression analysis of MERS coronavirus infections in conjunction with GEO2R.

1 The Benjamin and Hochberg method of p-value adjustment was used for ranking of
2 differential expression but raw p-values were used for assessment of statistical significance of
3 global differential expression. Log-transformation of data was auto-detected, and the NCBI
4 generated category of platform annotation was used.

6 A statistical test was performed to evaluate the significance of difference in FBXL8
7 mRNA expression levels in CALU3 2B4 cells with MERS-CoV infection as compared to CALU3
8 2B4 cells at baseline (0 hours) using a one-way ANOVA with Tukey's multiple comparisons test.

9 A statistical test was performed to evaluate the significance of difference in FBXL8 mRNA
10 expression levels in human primary microvascular endothelial cells with MERS-CoV as
11 compared to baseline infection at 0 hours using a one-way ANOVA with Tukey's multiple
12 comparisons test. Only p-values less than 0.05 were considered statistically significant. We
13 used PRISM for all statistical analyses (Version 8.4.0)(455).
14
15
16

17 **Results**

18 We mined two independent microarray datasets, public (3) and published (4) containing
19 transcriptome data from models of coronavirus infection in primary human cells and cell
20 culture. Across experimental models, we identified FBXL8 as differentially expressed following
21 coronavirus infection.

22 FBXL8 is differentially expressed in primary human microvascular endothelial cells when
23 comparing cells infected with wild-type Middle East respiratory syndrome coronavirus (MERS-
24 CoV), icMERS-CoV EMC2012 and uninfected cells.

25 We identified FBXL8 as differentially expressed following infection of primary human
26 microvascular endothelial cells with wild-type Middle East respiratory syndrome coronavirus
27 (MERS-CoV), icMERS-CoV EMC2012 when compared to non-infected cells (Table 1) (3). When
28 sorting all of the transcripts expressed in human microvascular endothelial cells measured by
microarray based on change in expression with and without infection, FBXL8 ranked first (1)

1 out of 34127 transcripts. Differential expression of FBXL8 in primary human microvascular
2 endothelial cells following infection with MERS-CoV was statistically significant (Table 1;
3 $p=3.42E-40$).

4
5 FBXL8 is differentially expressed in the human cell line HuH-7 when comparing cells with
6 human coronavirus 229E and uninfected cells.

7 We also identified FBXL8 as differentially expressed in the human cell line CALU3 2B4
8 when comparing cells infected with MERS-CoV London from at 3 hours, 7 hours, 12 hours, 18
9 hours and 24 hours when compared to cells infected at baseline (0 hours) (Table 2) (4). When
10 sorting all of the transcripts expressed in CALU3 2B4 cells measured by microarray based on
11 change in expression with and without infection with MERS-CoV London, FBXL8 ranked 3 out
12 of 28653 transcripts. Differential expression of FBXL8 in CALU3 2B4 cells following infection
13 with MERS-CoV London was statistically significant (Table 2; $p=1.22E-16$).

14
15 FBXL8 is transcriptionally repressed following infection of primary human microvascular
16 endothelial cells with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV),
17 icMERS-CoV EMC2012.

18 We extracted exact mRNA expression values for FBXL8 from primary human
19 microvascular endothelial cells infection with wild-type Middle East respiratory syndrome
20 coronavirus (MERS-CoV), icMERS-CoV, and from uninfected primary human microvascular
21 endothelial cells in order to compare expression levels of FBXL8 between these two groups
22 rather than relative to the rest of the transcriptome as assessed in differential gene expression
23 analysis. This dataset contained transcriptome information from infection of primary human
24 microvascular endothelial cells at 12 hours, 24 hours, 36 hours and 48 hours post-infection at
25 compared to baseline (0 hours). We also performed a statistical test to evaluate whether the
26 difference in expression of FBXL8 in primary human microvascular endothelial cells infection
27 with and without MERS-CoV infection was statistically significant. FBXL8 was expressed at
28 significantly higher levels at 12 hours, 24 hour, 36 hours, and 48 hours post-infection with
MERS-CoV as compared to cells infected at baseline (0 hours) (Figure 1; $p<0.0001$). FBXL8

expression peaked at 24 hours, then decreased from 24 hours to 36 hours and from 36 hours to 48 hours (Figure 1; $p < 0.0001$ between 24 and 36 hours, and between 36 and 48 hours).

FBXL8 is transcriptionally induced in the human cell line HuH-7 following infection with human coronavirus 229E.

We also extracted exact mRNA expression values for FBXL8 from CALU3 2B4 cells following infection with MERS-CoV London and from CALU3 2B4 cells infected with MERS-CoV London at baseline in order to compare expression levels of FBXL8 between these two groups. We also performed a statistical test to evaluate whether the difference in expression of FBXL8 in CALU3 2B4 cells following infection with MERS-CoV London as compared to cells infected with MERS-CoV London at baseline was statistically significant. Consistent with the induction of FBXL8 after infection of primary human microvascular endothelial cells with wild-type MERS-CoV, FBXL8 was expressed at significantly higher levels post-infection MERS-CoV London at 7, 12, 18 and 24 hours (Figure 2; $p < 0.0001$) but not at 3 hours (Figure 2; $p = 0.8455$).

Thus, we found that FBXL8 was among the genes whose expression changed most significantly after infection of human cells with MERS-CoV wild-type and MERS-CoV London, and that FBXL8 was transcriptionally induced in primary human microvascular endothelial cells and in a human lung adenocarcinoma cell line following infection with wild-type MERS-Cov and MERS-CoV London, respectively. FBXL8 was not significantly differentially expressed in the lungs of mice infected SARS coronavirus, in the blood and lungs of ferrets infected with SARS coronavirus, nor in a human cell line (Huh-7) infected with human coronavirus 229E (data not shown). Differential expression of FBXL8 after coronavirus infection was specific to MERS coronavirus.

Discussion

COVID19 is a newly discovered coronavirus that infects humans and whose spread has lead to a global pandemic (15) with zero available therapeutic strategies. To facilitate target

discovery and to further understanding of the basic transcriptional program of cells and tissues following coronavirus infection, we mined two independent microarray datasets containing transcriptome data of human cells infected with wild-type MERS-CoV and MERS-CoV London, in primary cells and in cell culture (3, 4). Across experimental models, we identified FBXL8 as one of the genes whose expression changed most significantly following infection with a MERS coronavirus. In both experimental models analyzed, FBXL8 expression significantly increased after infection with coronavirus.

In general, F-box proteins are named for their F-box motif that ranges in length from 42-48 amino acids known as a degron, and are most commonly known as part of a family of ubiquitin ligase complexes termed CRLs, or cullin RING ligases (16, 17). CRL1 is also known as SCF, which includes the S-phase kinase-associated protein Skp1, an E3 ligase core composed of the cullin CUL1 and RBX, as well as an F-box protein, of which there are approximately 70 in humans (17). Substrate recognition by SCF complexes occurs through the F-box subunit, by binding of a phosphorylated consensus sequence (“phosphodegron”) (16, 17). This provides modularity to SCF complexes in exerting their molecular function across a wide variety of proteins, the ubiquitination of substrates destined for the degradation by the proteasome (18, 19). F-box proteins are termed FBXLs if they possess an LRR (leucine-rich region), FBXWs if they possess a WD40 domain, and FBXOs if they possess some “other” domain (20). There is limited information regarding the specific functions of FBXL8 (21, 22), and we could not identify any literature regarding a role for FBXL8 in viral infection.

We found that FBXL8 was differentially expressed in a human cell line, CALU3 2B4 following infection with the human coronavirus MERS-CoV London as well as in primary human endothelial cells after infection with wild-type MERS-CoV. FBXL8 expression should be evaluated in the cells and tissues of patients infected with the novel coronavirus COVID19 to assess whether COVID19 infection similarly leads to induction of FBXL8 as seen after infection with MERS-CoVs. We suggest that transcriptional induction of FBXL8 is specific to MERS-CoV among the coronaviruses based on the relative lack of FBXL8 differential expression after

infection of mice with SARS-CoV and infection of a human liver cell line with human coronavirus 229E. Modulation of FBXL8 expression may represent a therapeutic strategy in MERS coronavirus infection, and the role of FBXL8 in COVID19 infection remains to be determined.

References

1. World Health Organization, 2020. Coronavirus disease 2019 (COVID-19): situation report, 49.
2. Wang, Y. and Zhu, L.Q., 2020. Pharmaceutical care recommendations for antiviral treatments in children with coronavirus disease 2019. *World Journal of Pediatrics*, pp.1-4.
3. GSE100509. Baric R, Sims A, Heller N, Waters KM, Einfeld AJ, Kawaoka Y. PNNL. 902 Battelle Blvd. Richland, USA.
4. Selinger, C., Tisoncik-Go, J., Menachery, V.D., Agnihothram, S., Law, G.L., Chang, J., Kelly, S.M., Sova, P., Baric, R.S. and Katze, M.G., 2014. Cytokine systems approach demonstrates differences in innate and pro-inflammatory host responses between genetically distinct MERS-CoV isolates. *BMC genomics*, 15(1), p.1161.
5. Baltimore, D., 1971. Expression of animal virus genomes. *Bacteriological reviews*, 35(3), p. 235.
6. Sawicki, S.G., Sawicki, D.L. and Siddell, S.G., 2007. A contemporary view of coronavirus transcription. *Journal of virology*, 81(1), pp.20-29.
7. Lai, M.M. and Cavanagh, D., 1997. The molecular biology of coronaviruses. In *Advances in virus research* (Vol. 48, pp. 1-100). Academic Press.
8. Peiris, J.S.M., Lai, S.T., Poon, L.L.M., Guan, Y., Yam, L.Y.C., Lim, W., Nicholls, J., Yee, W.K.S., Yan, W.W., Cheung, M.T. and Cheng, V.C.C., 2003. Coronavirus as a possible cause of severe acute respiratory syndrome. *The Lancet*, 361(9366), pp.1319-1325.
9. Lu, G., Hu, Y., Wang, Q., Qi, J., Gao, F., Li, Y., Zhang, Y., Zhang, W., Yuan, Y., Bao, J. and Zhang, B., 2013. Molecular basis of binding between novel human coronavirus MERS-CoV and its receptor CD26. *Nature*, 500(7461), pp.227-231.
10. van der Hoek, L., Pyrc, K., Jebbink, M.F., Vermeulen-Oost, W., Berkhout, R.J., Wolthers, K.C., Wertheim-van Dillen, P.M., Kaandorp, J., Spaargaren, J. and Berkhout, B., 2004. Identification of a new human coronavirus. *Nature medicine*, 10(4), pp.368-373.
11. van der Hoek, L., Pyrc, K., Jebbink, M.F., Vermeulen-Oost, W., Berkhout, R.J., Wolthers, K.C., Wertheim-van Dillen, P.M., Kaandorp, J., Spaargaren, J. and Berkhout, B., 2004. Identification of a new human coronavirus. *Nature medicine*, 10(4), pp.368-373.

12. Woo, P.C., Lau, S.K., Chu, C.M., Chan, K.H., Tsoi, H.W., Huang, Y., Wong, B.H., Poon, R.W., Cai, J.J., Luk, W.K. and Poon, L.L., 2005. Characterization and complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia. *Journal of virology*, 79(2), pp.884-895.
13. Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R. and Niu, P., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *New England Journal of Medicine*.
14. Bai, Y., Yao, L., Wei, T., Tian, F., Jin, D.Y., Chen, L. and Wang, M., 2020. Presumed asymptomatic carrier transmission of COVID-19. *Jama*.
15. World Health Organization, 2020. Coronavirus disease 2019 (COVID-19): situation report, 59.
16. Kipreos, E.T. and Pagano, M., 2000. The F-box protein family. *Genome biology*, 1(5), pp.reviews3002-1.
17. Skaar, J.R., Pagan, J.K. and Pagano, M., 2013. Mechanisms and function of substrate recruitment by F-box proteins. *Nature reviews Molecular cell biology*, 14(6), pp.369-381.
18. Hart, M., Concordet, J.P., Lassot, I., Albert, I., Del los Santos, R., Durand, H., Perret, C., Rubinfeld, B., Margottin, F., Benarous, R. and Polakis, P., 1999. The F-box protein β -TrCP associates with phosphorylated β -catenin and regulates its activity in the cell. *Current biology*, 9(4), pp.207-211.
19. Spencer, E., Jiang, J. and Chen, Z.J., 1999. Signal-induced ubiquitination of I κ B α by the F-box protein Slimb/ β -TrCP. *Genes & development*, 13(3), pp.284-294.
20. Jin, J., Cardozo, T., Lovering, R.C., Elledge, S.J., Pagano, M. and Harper, J.W., 2004. Systematic analysis and nomenclature of mammalian F-box proteins. *Genes & development*, 18(21), pp.2573-2580.
21. Li, Y., Yang, S., Cui, W., Mu, Y., Tang, Z., Chu, M., Peng, K. and Li, K., 2011. Cloning, sequencing and SNP site detection of the second intron of porcine FBXL8 gene. *Acta Agriculturae Universitatis Jiangxiensis*, 33(5), pp.938-941.
22. Randle, S.J. and Laman, H., 2016, February. F-box protein interactions with the hallmark pathways in cancer. In *Seminars in cancer biology* (Vol. 36, pp. 3-17). Academic Press.

Rank	ID	p-value	F	Gene	Gene name
1	A_33_P3284586/ NM_018378	3.42E-40	16915.7275	FBXL8	F-box and leucine-rich repeat protein 8

Table 1: FBXL8 is the most differentially expressed gene in primary human microvascular endothelial cells when comparing cells infected with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV EMC2012 and non-infected cells.

The rank of differential expression globally, the probe/transcript ID, the p-value of differential expression globally, F, a statistic used when more than two groups are compared in differential gene expression analysis (rather than the moderated t-statistic “t”), the gene and gene name are listed in this chart.

Rank	ID	p-value	F	Gene	Gene name
3	A_33_P3284586/ NM_018378	1.22E-16	574.2547	FBXL8	F-box and leucine-rich repeat protein 8

Table 2: FBXL8 is the third most differentially expressed in human lung adenocarcinoma cells (CALU3 2B4) when comparing cells infected with MERS-CoV London from 3 to 24 hours and cells infected at baseline.

The rank of differential expression globally, the probe/transcript ID, the p-value of differential expression globally, F, a statistic used when more than two groups are compared in differential gene expression analysis (rather than the moderated t-statistic “t”), the gene and gene name are listed in this chart.

Figure 1: FBXL8 is transcriptionally induced in primary human microvascular endothelial cells following infection with wild-type Middle East respiratory syndrome coronavirus (MERS-CoV), icMERS-CoV.

Expression of FBXL8 at the mRNA level is graphically represented with the mean value marked and the p-value listed for each comparison relative to baseline infection at 0 hours.

Figure 2: FBXL8 is transcriptionally induced in the human lung adenocarcinoma cell line CALU3 2B4 after infection with MERS-CoV London.

Expression of FBXL8 at the mRNA level is graphically represented with the mean value marked and the p-value listed for each comparison relative to baseline infection at 0 hours.