Over-expression of baculoviral IAP repeat containing 5 in human endometrial cancer.

Shahan Mamoor, MS¹
¹shahanmamoor@gmail.com
East Islip, NY USA

Gynecologic cancers including cancers of the endometrium are a clinical problem¹⁻⁴. We mined published microarray data^{5,6} to discover genes associated with endometrial cancers by comparing transcriptomes of the normal endometrium and endometrial tumors from humans. We identified baculoviral IAP repeat containing 5, encoded by BIRC5, as among the most differentially expressed genes, transcriptome-wide, in cancers of the endometrium. BIRC5 was expressed at significantly higher levels in endometrial tumor tissues as compared to the endometrium. Importantly, primary tumor expression of BIRC5 was correlated with overall survival in patients with endometrial cancer. BIRC5 may be a molecule of interest in understanding the etiology or progression of human endometrial cancer.

Keywords: endometrial cancer, gynecologic cancers, endometrium, BIRC5, baculoviral IAP repeat containing 5, systems biology of endometrial cancer, targeted therapeutics in endometrial cancer.

Endometrial cancer is the most common gynecologic cancer in the developed world¹. Over the last three decades, the incidence of endometrial cancer has increased 21%⁴ and the death rate has increased 100%³. We harnessed the power of independently published microarray datasets^{5,6} to determine in an unbiased fashion and at the systems-level genes most differentially expressed in endometrial tumors. We report here the differential and increased expression of the baculoviral IAP repeat containing 5 (BIRC5) in human endometrial cancer.

Methods

We utilized datasets GSE63678⁵ and GSE17025⁶ for this global differential gene expression analysis of human endometrial cancer in conjunction with GEO2R. GSE63678 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=5$ control endometrial tissues (including $n=4$ uterine myomas and $n=1$ benign cyst) and $n=7$ endometrial cancers (including $n=2$ endometrial adenocarcinomas, $n=3$ mixed endometrioid adenocarcinomas, and $n=2$ adenocarcinomas with squamous differentiation); analysis was performed using platform GPL571. GSE17025 was generated using Affymetrix Human Genome U133 Plus 2.0 Array technology with $n=5$ normal endometrial tissues (inactive endometrium) and $n=91$ endometrial cancers; analysis was performed using platform GPL570. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether BIRC5 gene expression was significantly different between control endometrial tissue and endometrial tumor tissue in humans using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter tool⁷ for correlation of BIRC5 mRNA expression levels with overall survival in $n=543$ endometrial cancer patients.

Results

We harnessed the power of blind comparative transcriptome analysis using published microarray data^{5,6} to discover in an unbiased fashion genes associated with endometrial cancer in humans.

BIRC5 is differentially expressed in endometrial cancer.

We identified baculoviral IAP repeat containing 5, encoded by BIRC5, as among the genes most differentially expressed in cancers of the endometrium when compared to benign endometrial tissues (Chart 1). When sorting each of the genes expressed in endometrial tumor tissue based on significance of change in expression as compared to benign endometrial tissue, BIRC5 ranked 51 out of 22273 transcripts, equating to 99.8% differential expression (Chart 1). Differential expression of BIRC5 in human endometrial cancers was statistically significant

(Chart 1; $p=3.77\text{E-}05$).

We queried a second microarray data to validate differential expression of BIRC5 in endometrial cancer. Again, we observed differential expression of BIRC5 when comparing endometrial tumor tissue to benign endometrial tissue (Chart 2). When sorting each of the genes expressed in endometrial tumor tissue based on significance of change in expression as compared to benign endometrial tissue, BIRC5 ranked 138 out of 54675 transcripts, equating to 99.7% differential expression (Chart 2). Differential expression of BIRC5 in human endometrial cancers was statistically significant (Chart 2; $p=2.03\text{E-}07$).

BIRC5 is expressed at significantly higher levels in endometrial cancers as compared to benign endometrial tissue.

We obtained exact mRNA expression levels for BIRC5 in endometrial tumor tissues and from benign endometrial tissue to evaluate direction and statistical significance of change in expression of BIRC5 in human endometrial cancer. BIRC5 was expressed at higher levels in endometrial tissue as compared to normal endometrial tissue, and this difference was statistically significant (Figure 1; $p<0.0001$). We calculated a mean fold change of 1.49 in BIRC5 mRNA levels in human endometrial cancer, as BIRC5 was expressed at 5.32 ± 0.58 arbitrary units (A.U.) in control endometrial tissue but at 7.92 ± 0.87 A.U. in endometrial tumor tissue.

BIRC5 expression is correlated with patient survival outcomes in endometrial cancer.

We performed Kaplan-Meier survival analysis to evaluate correlation between BIRC5 primary tumor expression and survival outcomes in 543 patients with endometrial cancer. We observed a correlation between primary tumor expression of BIRC5 and overall survival (OS) in patients with endometrial cancer (Figure 2). BIRC5 mRNA levels were a negative prognostic indicator in patients with endometrial cancer. Patients whose primary tumors expressed low levels of BIRC5 possessed median OS of 78.4 months as compared to a median OS of 47.43 months for patients whose tumors expressed high levels of BIRC5 (Figure 2). This difference in OS based on BIRC5 tumor expression in patients with endometrial cancer approached statistical significance (Figure 2, Chart 3; logrank p -value: 0.069; hazard ratio: 1.47 (0.97-2.24)).

Thus, by mining published microarray data^{5,6} in an unbiased and systematic fashion, we identified baculoviral IAP repeat containing 5, encoded by BIRC5, as among the genes whose expression was most different, transcriptome-wide, in the endometrial tumor tissue of patients with endometrial cancer when compared to benign endometrial tissue; we observed significantly increased expression of BIRC5 in endometrial tumor tissue as compared to benign endometrial tissue. Further, we found a correlation between BIRC5 expression and patient survival outcomes in human endometrial cancer, as overall survival was inferior in patients whose tumors expressed higher levels of BIRC5 as compared to patients whose tumors expressed lower levels of BIRC5.

Discussion

We provided evidence here that baculoviral IAP repeat containing 5 is among the genes most differentially expressed in human endometrial cancer, that mRNA for BIRC5 is present at significantly increased quantity in endometrial tumor tissue as compared to benign endometrium, and that BIRC5 primary tumor expression is correlated with overall survival in patients with endometrial cancer. These data suggest BIRC5 may be of importance to fundamental biological processes that underlie the initiation, progression or maintenance of human endometrial cancer.

References

1. Amant, F., Moerman, P., Neven, P., Timmerman, D., Van Limbergen, E. and Vergote, I., 2005. Endometrial cancer. *The Lancet*, 366(9484), pp.491-505.
2. Sorosky, J.I., 2008. Endometrial cancer. *Obstetrics & Gynecology*, 111(2), pp.436-447.
3. Morice, P., Leary, A., Creutzberg, C., Abu-Rustum, N. and Darai, E., 2016. Endometrial cancer. *The Lancet*, 387(10023), pp.1094-1108.
4. Sorosky, J.I., 2012. Endometrial cancer. *Obstetrics & Gynecology*, 120(2 Part 1), pp.383-397.
5. Pappa, K.I., Polyzos, A., Jacob-Hirsch, J., Amariglio, N., Vlachos, G.D., Loutradis, D. and Anagnou, N.P., 2015. Profiling of discrete gynecological cancers reveals novel transcriptional modules and common features shared by other cancer types and embryonic stem cells. *PLoS One*, 10(11), p.e0142229.
6. Day, R.S., McDade, K.K., Chandran, U.R., Lisovich, A., Conrads, T.P., Hood, B.L., Kolli, V.K., Kirchner, D., Litzi, T. and Maxwell, G.L., 2011. Identifier mapping performance for integrating transcriptomics and proteomics experimental results. *BMC bioinformatics*, 12(1), p.213.
7. Nagy, A., Munkacsy, G. and Gyorffy, B., 2020. Pancancer survival analysis of cancer hallmark genes. *bioRxiv*.

Rank: 51
Probe ID: 202095_s_at
p-value: 3.77E-05
t: -6.2894029
B: 2.54566
Gene: BIRC5
Gene name: baculoviral IAP repeat containing 5

Chart 1: BIRC5 is differentially expressed in endometrial cancer when comparing primary endometrial tumors to benign endometrial tissue.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 138
probe ID: 202095_s_at
p-value: 2.03E-07
t: -5.5897414
B: 6.87831
Gene: BIRC5
Gene name: baculoviral IAP repeat containing 5

Chart 2: BIRC5 is differentially expressed in endometrial cancer when comparing primary endometrial tumors to benign endometrial tissue.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: BIRC5 is expressed at significantly higher levels in the endometrial tumors of patients with endometrial cancer when compared to benign endometrium.

The mRNA expression level of BIRC5 in benign endometrial tissue (left) and in primary tumors of the endometrium (right) is graphically depicted; the result of a statistical test evaluating significance of difference in BIRC5 expression between benign endometrial tissue and primary tumors of the endometrial tissue is $p<0.0001$.

Figure 2: Correlation between BIRC5 primary tumor expression and overall survival in patients with endometrial cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival for $n=543$ total endometrial cancer patients stratified into two groups, based on low or high expression of BIRC5 in patient primary tumors. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on BIRC5 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low BIRC5 expression: 78.4 months
High BIRC5 expression: 47.43 months

Chart 3: Median overall survival is superior in endometrial cancer patients with low primary tumor expression of BIRC5.

The median OS (overall survival) of patients with low primary tumor expression of BIRC5 and high primary tumor expression of BIRC5 is listed in this chart, for $n=543$ endometrial cancer patients.