Biglycan (BGN) is differentially expressed in metastatic breast cancer, both in brain metastasis and in the primary tumor.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes to discover genes associated with brain metastasis in patients with metastatic breast cancer. We found that biglycan, encoded by BGN, was among the genes whose expression was most different in the brain metastases of with patients with metastatic breast cancer as compared to primary tumors of the breast. Interestingly, biglycan was also among the genes most differentially expressed transcriptome-wide when comparing primary tumors of the breast to normal breast tissue. We observed significant down-regulation of biglycan in metastasis to the brain. Molecular functions and down-regulation of BGN may be important for metastasis of primary tumor-derived cancer cells the brain in humans with metastatic breast cancer, and suggests a role for changes in biglycan expression during a spectrum of transformation from benign tissue of the breast, primary tumor and finally to metastasis of the brain.

Keywords: breast cancer, metastasis, brain metastasis, central nervous system metastasis, BGN, biglycan, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in trastuzumab-treated patients with breast cancer². This alarmingly high frequency of CNS metastasis events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with metastatic breast cancer^{4,5}. We discovered transcriptome-wide differential expression of BGN in brain metastases and primary tumors in human breast cancer.

Methods

We used datasets GSE125989⁴ and GSE29044⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE125989 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=16$ primary tumors from patients with breast cancer and $n=16$ brain metastases from patients with brain metastatic breast cancer. GSE29044, a microarray dataset, was generated using Affymetrix Human Genome U133 Plus 2.0 Array with $n=36$ normal breast tissues and $n=73$ primary tumors of the breast from patients with breast cancer; platform GPL570 was used for this analysis. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether BGN gene expression was significantly between primary breast tumors and brain metastases in humans with breast cancer using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

Results

We performed global comparative transcriptome analysis of the primary and metastatic tumor tissues of patients with brain metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

BGN is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

We identified biglycan, encoded by BGN, as among the genes whose expression was most significantly different in the brain metastases of patients with brain metastatic breast cancer⁴ (Table 1). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to the primary tumors of patients with brain metastatic breast cancer, BGN ranked 115 out of 22277 total transcripts (Table 1), equating to 99.48% differential expression. Differential expression of BGN in the brain metastases of patients with brain metastatic breast cancer was statistically significant (Table 1; $p=5.42E-04$).

We asked whether we could validate differential expression of BGN was specific to brain metastasis or if we could observe differential expression of BGN in primary tumors of the breast. Comparison of global gene expression profiles of 36 normal breast tissues to 73 primary tumors of the breast⁵ revealed that BGN was among the genes whose expression was most significantly different

transcriptome-wide between the primary tumors of the breast and benign breast tissue (Table 2). When sorting each of the genes expressed in primary tumors of the breast based on significance of difference in expression as compared to benign breast tissue, BGN ranked 516 out of 54675 total transcripts (Table 2), equating to 99.1% differential expression. Differential expression of BGN in primary tumors of the breast in human breast cancer was statistically significant (Table 2; $p=2.58E-13$).

BGN is expressed at significantly lower levels in the brain metastases of patients with brain metastatic breast cancer.

We obtained exact mRNA expression levels for BGN, in primary tumors of the breast and in brain metastasis of patients with brain metastatic breast cancer to determine magnitude and direction of change in BGN expression in brain metastatic tissues. BGN was expressed at lower levels in the brain metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 1; $p=0.0009$). We calculated a mean fold change of 0.84 ± 0.13 in BGN expression when comparing primary tumors of the breast to brain metastatic tissues in patients with breast cancer (Table 1).

Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified BGN as a differentially expressed gene in metastatic breast cancer, both in brain metastases when to primary tumors of the breast, and in primary tumors of the breast when compared to benign breast tissue; BGN was expressed at significantly lower levels in brain metastases as compared to primary tumors of the breast.

Discussion

We provided evidence here that biglycan, encoded by BGN is among the genes whose expression is most different in the tumor tissues of patients with metastatic breast cancer, in primary tumors of the breast and in brain metastases. BGN may be of relevance to any number of steps required for transformation and metastasis to the to the brain, and suggests significant changes in biglycan expression during oncogenesis of the breast and subsequent metastasis to the brain in humans with metastatic breast cancer.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Iwamoto, T., Niikura, N., Ogiya, R., Yasojima, H., Watanabe, K.I., Kanbayashi, C., Tsuneizumi, M., Matsui, A., Fujisawa, T., Iwasa, T. and Shien, T., 2019. Distinct gene expression profiles between primary breast cancers and brain metastases from pair-matched samples. *Scientific reports*, 9(1), pp. 1-8.
5. Colak, D., Nofal, A., AlBakheet, A., Nirmal, M., Jeprel, H., Eldali, A., Taher, A.T., Tulbah, A., Ajarim, D., Al Malik, O. and Inan, M.S., 2013. Age-specific gene expression signatures for breast tumors and cross-species conserved potential cancer progression markers in young women. *PloS one*, 8(5), p.e63204.

Rank	ID	p-value	t	B	FC	Gene	Gene name
115	201262_s_at	5.42E-04	-3.8130924	-0.1979	0.84 ± 0.13	BGN	biglycan

Table 1: BGN is differentially expressed in brain metastases in metastatic breast cancer when compared to primary tumors of the breast.

Rank of differential expression, probe ID, *p*-value with respect to differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change (FC) of BGN expression between the brain metastases of patients with breast cancer and primary tumors of the breast, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
516	201261_x_at	2.58E-13	-8.35	19.8948522	BGN	biglycan

Table 2: BGN is differentially expressed in primary tumors of the breast when compared to normal breast tissues.

Rank of differential expression, probe ID, *p*-value with respect to differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: BGN is expressed at significantly lower levels in the brain metastases of patients with brain metastatic breast cancer.

The mRNA expression level of BGN in primary tumors of the breast (left) and in the brain metastases of brain metastatic breast cancer patients (right) is graphically represented here with mean BGN levels marked and the result of a statistical test evaluating difference in mRNA expression level between brain metastases and primary tumors of the breast, a *p*-value, listed above.