The 65 kilodalton regulatory subunit of the protein phosphatase 2A is differentially expressed in brain metastatic human breast cancer.

Shahan Mamoor, MS¹
¹shahanmamoor@gmail.com
East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with brain metastasis in humans with metastatic breast cancer. We found that the 65 kilodalton regulatory subunit of the protein phosphatase 2A, encoded by PPP2R1A, was among the genes whose expression was most different in the brain metastases of patients with metastatic breast cancer as compared to primary tumors of the breast⁴. PPP2R1A was also differentially expressed in the tumor cells of patients with triple negative breast cancer⁵. PPP2R1A mRNA was present at increased quantities in brain metastatic tissues as compared to primary tumors of the breast. Importantly, expression of PPP2R1A in primary tumors was significantly correlated with patient recurrence-free survival. Modulation of PPP2R1A expression may be relevant to the biology by which tumor cells metastasize from the breast to the brain in humans with metastatic breast cancer.

Keywords: breast cancer, metastasis, brain metastases, central nervous system metastases, the 65 kilodalton regulatory subunit of the protein phosphatase 2A, PPP2R1A, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with brain metastatic breast cancer^{4,5}. We discovered differential and increased expression of the gene encoding the 65 kilodalton regulatory subunit of the protein phosphatase 2A, PPP2R1A, in brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE38959⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=11$ primary breast tumors and $n=3$ brain metastases from patients with breast cancer; analysis was performed using platform GPL885. GSE38959 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=13$ samples of normal mammary gland ductal cells and $n=30$ samples of primary tumor cells from patients with triple negative breast cancer; analysis was performed using platform GPL4133. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether PPP2R1A gene expression was significantly different between primary tumors of the breast and brain metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of PPP2R1A mRNA expression levels with recurrence-free survival in $n=3951$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of brain metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

PPP2R1A is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified the 65

1 kilodalton regulatory subunit of the protein phosphatase 2A, encoded by PPP2R1A, as a
2 differentially expressed gene in the breast metastatic tissues of humans with breast cancer (Table
3 1). When sorting each of the genes expressed in brain metastases based on significance of
4 difference as compared to primary tumors of the breast in patients with breast cancer, PPP2R1A
5 ranked 191 out of 17418 total transcripts (Chart 1), equating to 99.9% differential expression.
Differential expression of PPP2R1A in the brain metastases of patients with metastatic breast
cancer was statistically significant (Chart 1; $p=5.03E-04$).

6 To attempt to validate differential transcriptome-wide differential expression of PPP2R1A
7 in human breast cancer, we queried a second microarray dataset⁵, here comparing primary tumor
8 cells from patients with triple negative breast cancer to normal mammary gland ductal cells.
9 Again, we identified PPP2R1A as a differentially expressed gene in the tumor tissues of patients
10 with breast cancer (Chart 2). When sorting each of the genes expressed in triple negative breast
11 cancer tumor cells based on significance of difference as compared to normal mammary gland
12 ductal cells, PPP2R1A ranked 17837 out of 45015 total transcripts (Chart 2), equating to 60.4%
differential expression. Differential expression of PPP2R1A in the brain metastases of patients
with metastatic breast cancer trended towards statistical significance (Chart 2; $p=7.02E-02$). Thus,
differential expression of PPP2R1A, transcriptome-wide, in the tumor tissues of women with
breast cancer was conserved across two independent microarray datasets, both when in primary
and metastatic tumor tissues.

13 **PPP2R1A is expressed at higher levels in the brain metastases of patients with metastatic** 14 **breast cancer.**

15 We obtained exact mRNA expression levels for PPP2R1A, in primary tumors of the breast
16 and in brain metastasis of patients with brain metastatic breast cancer to determine direction and
17 statistical significance of change in PPP2R1A expression in brain metastatic tissues. PPP2R1A
18 was expressed at higher levels in the brain metastases of patients with breast cancer as compared
to primary tumors of the breast, and this difference was statistically significant (Figure 1;
 $p=0.000323$).

19 **PPP2R1A expression is significantly correlated with survival outcomes in human breast** 20 **cancer.**

21 We performed Kaplan-Meier survival analysis⁸ in 3951 breast cancer patients in total, to
22 evaluate whether PPP2R1A tumor expression was correlated with survival outcomes in human
23 breast cancer. We observed a statistically significant correlation between primary tumor
24 expression of PPP2R1A and recurrence-free survival (RFS) in patients with breast cancer in the
lower survival quartile (Figure 2). Patients whose primary tumors expressed low levels of
PPP2R1A possessed median RFS of 191.21 months, while patients whose tumors expressed high
levels of PPP2R1A possessed median RFS of 216.66 months. This difference in RFS based on
PPP2R1A tumor expression in patients with breast cancer was statistically significant (Figure 2,
Chart 3; logrank p -value: 0.021; hazard ratio: 0.87 (0.76-0.98)).

27 Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we
28 identified the 65 kilodalton regulatory subunit of the protein phosphatase 2A, encoded

1 by PPP2R1A, as among the genes whose expression was most different, transcriptome-wide, in
2 the brain metastases of patients with breast cancer, when compared to primary tumors of the
3 breast and in the primary tumor cells of patients with triple negative breast cancer when
4 compared to normal mammary gland ductal cells; we observed significantly increased expression
5 of PPP2R1A in brain metastases as compared to primary tumors of the breast. Further, we found
6 a significant correlation between PPP2R1A expression and patient survival outcomes, as
recurrence-free survival was significantly higher in patients whose primary
tumors expressed higher levels of PPP2R1A as compared to patients whose primary tumors
expressed lower levels of PPP2R1A.

7 **Discussion**

8 We provided evidence here that the 65 kilodalton regulatory subunit of the protein
9 phosphatase 2A, encoded by PPP2R1A, is among the genes whose expression is most different in
10 the brain metastases of patients with brain metastatic breast cancer, that PPP2R1A is also
11 differentially expressed in the primary tumor cells of patients with triple negative breast cancer,
12 that PPP2R1A mRNA is present at significantly increased quantities in brain metastatic tissues as
13 compared to primary tumors of the breast, and that primary tumor PPP2R1A expression is
14 significantly correlated with patient survival outcomes in human breast cancer. Evaluation of the
15 effects of genetic depletion of PPP2R1A in mouse models of metastatic breast cancer on
16 metastasis to the central nervous system is merited. Modulation of PPP2R1A expression may be
17 relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or
18 lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier
19 and colonize the brain.
20
21
22
23
24
25
26
27
28

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Komatsu, M., Yoshimaru, T., Matsuo, T., Kiyotani, K., Miyoshi, Y., Tanahashi, T., Rokutan, K., Yamaguchi, R., Saito, A., Imoto, S. and Miyano, S., 2013. Molecular features of triple negative breast cancer cells by genome-wide gene expression profiling analysis. *International journal of oncology*, 42(2), pp.478-506.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 191
Probe ID: 7021
P-value: 5.03E-04
t: -4.36
B: 2.641979
Gene: 0.04121
Gene name: protein phosphatase 2 (formerly 2A), regulatory subunit A , alpha isoform

Chart 1: PPP2R1A is differentially expressed in brain metastatic breast cancer when comparing brain metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 17837
probe ID: 10839
p-value: 7.02E-02
t: -1.8545269
B: -5.089554
Gene: PPP2R1A
Gene name: the 65 kilodalton regulatory subunit of the protein phosphatase 2A

Chart 2: PPP2R1A is differentially expressed in the tumor cells of patients with triple negative breast cancer when compared to normal mammary gland ductal cells.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: PPP2R1A is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of PPP2R1A in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in PPP2R1A expression between primary tumors and brain metastasis, a p -value, is $p=0.000323$.

Figure 2: Significant correlation between PPP2R1A primary tumor expression and recurrence-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=3951$ total patients stratified into two groups, based on low or high expression of PPP2R1A in patient primary tumors, in the lower survival quartile. The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on PPP2R1A expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low PPP2R1A expression: 191.21 months

High PPP2R1A expression: 216.66 months

Chart 3: Median recurrence-free survival is inferior in patients with low primary tumor expression of PPP2R1A.

The median RFS (recurrence-free survival) of patients with low primary tumor expression of PPP2R1A and high primary tumor expression of PPP2R1A is listed in this chart, for $n=3951$ breast cancer patients, in the lower survival quartile.