PTPN9 is differentially expressed in the tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab (Herceptin) is a monoclonal antibody targeting the extracellular domain of the human epidermal growth factor receptor 2 (HER2)¹ utilized for the treatment of adjuvant and metastatic breast cancer² in the United States and worldwide. We mined published microarray data^{3,4} to discover in an unbiased manner the most significant transcriptional changes associated with trastuzumab treatment. We identified PTPN9 among the genes most differentially expressed in the primary tumors of patients with breast cancer treated with trastuzumab. The primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of PTPN9 messenger RNA than did patients not treated with trastuzumab, and a single administration of trastuzumab was sufficient to result in differential expression of PTPN9 in primary tumors of the breast, demonstrating decreased expression of a negative regulator of the molecular target of trastuzumab⁵ as an apparent transcriptional consequence of trastuzumab administration in patients with breast cancer.

Keywords: trastuzumab, breast cancer, PTPN9, protein tyrosine phosphatase, non-receptor type 9, systems biology of breast cancer, targeted therapeutics in breast cancer.

The mechanism of action ascribed to trastuzumab function is thought to be binding of the F_{ab} domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2 (HER2)¹, but the transcriptional consequences resulting from signaling downstream of trastuzumab binding to HER2 is less well understood. Through unbiased, systems-level comparative transcriptome analysis of published microarray data^{3,4}, we report here that PTPN9 is a differentially expressed gene in the tumors of trastuzumab-treated patients: patients treated with trastuzumab expressed significantly higher levels of PTPN9 than did those of patients not treated with trastuzumab.

Methods

We used datasets GSE130788³, and GSE114082⁴ for this differential gene expression analysis of the transcriptional response to trastuzumab treatment in patients with breast cancer. GSE130788 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with $n=11$ primary breast tumors from patients treated with trastuzumab and $n=36$ primary tumors from patients treated with trastuzumab, analysis performed using platform GPL6480. GSE114082 was generated using Illumina HumanHT-12 WG-DASL V4.0 R2 expression beadchip technology with $n=17$ primary breast tumors from patients before trastuzumab treatment and $n=17$ primary tumors from patients after trastuzumab treatment, analysis performed using platform GPL14951. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global or multiplexed differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether PTPN9 gene expression was significantly between primary breast tumors of patients receiving trastuzumab using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

Results

We performed unbiased comparative transcriptome analysis using published microarray data^{3,4} to identify the most significant transcriptional features of the primary breast tumors of patients treated with trastuzumab (Herceptin).

PTPN9 is differentially expressed in the tumors of patients treated with trastuzumab.

We identified protein tyrosine phosphatase, non-receptor type 9, encoded by PTPN9 as among the genes whose expression changed most significantly in the tumors of breast cancer patients treated with trastuzumab³ (Table 1). When sorting each of the genes expressed in the primary tumors of breast cancer patients treated with trastuzumab based on significance of change as compared to that of patients not treated with trastuzumab, PTPN9 ranked 237 out of 41000 total transcripts (Table 1), equating to 99.4% differential expression. PTPN9 differential expression in the tumors of patients treated with trastuzumab trended towards statistical significance (Table 1; $p=1.88E-44$).

We turned to a second microarray dataset to validate our findings of PTPN9 differential expression in the patients treated with trastuzumab. The tumors compared in this dataset were sampled from the same patient before and after a single, brief exposure to trastuzumab, not comparing tumors from different patients treated with a course of trastuzumab or not. Thus, while differentially expressed genes when comparing tumors based on trastuzumab treatment can result from a variety of sources, including gene expression changes resulting from tumor evolution over the course of treatment, changes identified

1 in this dataset, occurring after only a single treatment of trastuzumab, are more likely to be a direct
2 transcriptional result of trastuzumab binding events. We found again that PTPN9 was among the genes
3 most differentially expressed in the tumors of patients administered a single treatment of trastuzumab
4 (Table 2). When sorting each of the genes expressed in the tumors of patients administered a single
5 treatment of trastuzumab, PTPN9 ranked 2709 out of 29377 total transcripts (Table 2), equating to 90.8%
6 differential expression. Differential expression of PTPN9 in the primary tumors of breast cancer patients
7 following treatment with trastuzumab was statistically significant (Table 2; $p=1.12E-03$).

8 **PTPN9 is expressed at significantly lower levels in the tumors of breast cancer patients treated with**
9 **trastuzumab than in the tumors of those not treated with trastuzumab.**

10 We obtained exact mRNA expression levels for PTPN9 from tumors of patients treated with
11 trastuzumab and from the tumors of patients not treated with trastuzumab to directly evaluate direction
12 and statistical significance of change in expression of PTPN9 based on history of trastuzumab treatment.
13 PTPN9 was expressed at lower levels in the tumors of patients treated with trastuzumab as compared to
14 those of patients not treated with trastuzumab, and this difference was statistically significant (Figure 1;
15 $p<0.0001$).

16 Thus, we found using published microarray data^{3,4} that PTPN9 was among the genes most
17 differentially expressed in the tumors of patients treated with trastuzumab, that PTPN9 was expressed at
18 lower levels in the tumors of patients treated with trastuzumab as compared to those not treated with
19 trastuzumab, and that this transcriptional change was likely a direct result of trastuzumab treatment as
20 opposed to a gene expression change associated with clonal selection and evolution of the tumor across
21 the course of treatment.

22 **Discussion**

23 We provided evidence here that PTPN9 is a differentially expressed gene in the tumors of patients
24 treated with trastuzumab, and expressed at significantly lower levels in the primary tumors of breast
25 cancer patients treated with trastuzumab as compared to patients not treated with trastuzumab. Thus, a
26 direct transcriptional consequence of trastuzumab administration, a treatment that has been associated
27 with a 34% rate of central nervous system metastases⁶, is down-regulation of a negative regulator of the
28 molecular target of trastuzumab, ErbB2 (HER2)⁵.

References

1. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
2. Herceptin (Trastuzumab). Indications. <https://www.herceptin.com/hcp/indications.html>.
3. GSE130788. A Phase II Randomized Trial of Neoadjuvant Trastuzumab or Lapatinib or the Combination of Trastuzumab and Lapatinib, Followed by Docetaxel and Carboplatin with Trastuzumab and/or Lapatinib in Patients with HER2+ Breast Cancer. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788>. Hsiaowang Chen. UCLA, Department of Medicine. Santa Monica, CA.
4. Triulzi, T., Regondi, V., De Cecco, L., Cappelletti, M.R., Di Modica, M., Paolini, B., Lollini, P.L., Di Cosimo, S., Sfondrini, L., Generali, D. and Tagliabue, E., 2018. Early immune modulation by single-agent trastuzumab as a marker of trastuzumab benefit. *British journal of cancer*, 119(12), pp. 1487-1494.
5. Yuan, T., Wang, Y., Zhao, Z.J. and Gu, H., 2010. Protein-tyrosine phosphatase PTPN9 negatively regulates ErbB2 and epidermal growth factor receptor signaling in breast cancer cells. *Journal of Biological Chemistry*, 285(20), pp.14861-14870.
6. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.

Rank	ID	p-value	t	B	Gene	Gene name
237	A_24_P687372	1.88E-44	-18.29257	90.520044	PTPN9	protein tyrosine phosphatase, non-receptor type 9

Table 1: PTPN9 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene
2709	ILMN_1789504	1.12E-03	-3.5418971	-0.9833	PTPN9

Table 2: A single exposure to trastuzumab is sufficient to drive differential expression of PTPN9 in the primary tumors of patients with breast cancer.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: PTPN9 is expressed at significantly higher levels in the primary tumors of patients treated with trastuzumab.

The mRNA expression level of PTPN9 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between PTPN9 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.