One Year Surveillance of Body Mass Index and Cardiorespiratory Fitness in Primary School Children in North West England and the Impact of School Deprivation Level

Post-print article published on: 18/12/2018
(doi: [10.1136/archdischild-2018-315567](https://doi.org/10.1136/archdischild-2018-315567))

Research conducted by the ukactive Research Institute

Final article published in Archives of Disease in Childhood

Citation: Mann, S., Wade, M., Jones, M., Sandercock, G., Beedie, C., & Steele, J. (2019). One Year Surveillance of Body Mass Index and Cardiorespiratory Fitness in Primary School Children in North West England and the Impact of School Deprivation Level. *Archives of Disease in Childhood*. [10.1136/archdischild-2018-315567](https://doi.org/10.1136/archdischild-2018-315567)

SUMMARY OF KEY FINDINGS

- Both body mass index (BMI) and cardiorespiratory fitness (CRF) vary over the period of one year, including the school term and summer recess, in primary school children aged 9-10 years.
- CRF significantly increased throughout the school year, yet decreased back to baseline during the summer recess.
- This effect was influenced by school deprivation levels and pupils from the most deprived areas saw significantly greater reductions in CRF compared with pupils from affluent areas
- These results highlight a need for public health interventions aimed at increasing children's PA levels, particularly vigorous PA, to be targeted towards the summer holiday period.

One Year Surveillance of Body Mass Index and Cardiorespiratory Fitness in Primary School Children in North West England and the Impact of School Deprivation Level

Steven Mann Ph.D.^{1,2}, Matthew Wade MSc^{1,3}, Michelle Jones Ph.D.⁴, Gavin Sandercock Ph.D.⁵, Chris Beedie Ph.D.⁶, James Steele Ph.D.^{1,4}

¹Ukactive Research Institute, London, UK; ²Centre for Applied Biological and Exercise Sciences, Faculty of Health and Life Sciences, Coventry University, Coventry, UK; ³School of Sport Health and Applied Science, St Mary's University, Twickenham, UK; ⁴School of Sport, Health, and Social Sciences, Southampton Solent University, UK; ⁵Department of Biological Sciences, University of Essex, Wivenhoe Park, Colchester CO4 3SQ; ⁶School of Human and Life Sciences, Canterbury Christ Church University, Canterbury, Kent, CT1 1QU
Contact for Correspondence jamessteele@ukactive.org.uk
Twitter: [@jamessteeleii](https://twitter.com/jamessteeleii)

SCIENTIFIC ABSTRACT

Objectives: Cardiorespiratory fitness (CRF) is independently associated with health and academic attainment in childhood and adolescence. Yet overweight/obesity remain the focus in public health policy. Surveillance of BMI and CRF considering school deprivation levels is limited. Therefore, we examined this in English Primary Schools. **Methods:** Participants (n=409) were students (9-10 years), from 13 schools. BMI and CRF (20 m shuttle-run) were measured at three time-points across the academic year and a fourth after summer recess. **Results:** BMI z-scores significantly decreased ($p = 0.015$) from autumn ($z = 0.336$ [CI 0.212 to 0.460]) to spring ($z = 0.252$ [CI 0.132 to 0.371]), and then significantly increased ($p = 0.010$) to summer ($z = 0.327$ [CI 0.207 to 0.447]). CRF significantly increased ($p < 0.001$) from autumn ($z = 0.091$ [CI -0.014 to 0.196]) to spring ($z = 0.492$ [CI 0.367 to 0.616]), no change ($p = 0.308$) into summer ($z = 0.411$ [CI 0.294 to 0.528]), and a significant decrease ($p < 0.001$) into the following autumn term ($z = 0.125$ [CI 0.021 to 0.230]). BMI was unaffected by deprivation; however, pupils from the most deprived areas saw significantly greater reductions in CRF compared with pupils from affluent areas. No time, or deprivation level, by sex interactions were found. **Conclusion:** Significant reductions in children's CRF occurred over the summer recess and was greater among children from schools in the most deprived areas. This may help inform future research into interventions targeting physical activity of school children, particularly over the summer recess.

Key words: Physical Fitness; Sport; Health Policy; Obesity

INTRODUCTION

Cardiorespiratory fitness (CRF) is modifiable by physical activity (PA) and independently associated with health [1], and higher academic attainment [2,3]; in children and adolescents. Evidence supports tracking CRF from childhood to adulthood [4], and that childhood CRF predicts adult cardiovascular disease and metabolic syndrome risk [5], highlighting the importance of maintaining and improving children's CRF [6]. Lang et al. [7] proposed these factors considered together indicate that population status of CRF in children may help predict future non-communicable disease burden.

Cross-sectional data and meta-analyses have demonstrated rapid secular decline in CRF globally of around 0.36% per annum increasing in magnitude since

1970 [8]. Whilst public health agencies encourage novel interventions to increase childhood PA, widespread lack of routine health data hampers evaluation of their impact [9]. Indeed, the UK Chief Medical Officer has stated, "*The introduction of a standardised school based fitness assessment in England may have multiple benefits that extend beyond the benefits for the individual*" [10].

Currently, the only mandated measure of children's health status in England is the National Child Measurement Programme (NCMP [11]). The NCMP is an annual programme that measures height and weight of children in Reception (aged 4–5 years) and Year 6 (aged 10–11 years). Ease of implementation is a key consideration for school based measurement with height and weight easily measured to determine body mass index

(BMI). Tracking of CRF levels should also follow a feasible and scalable approach.

The 20 m shuttle run test (20mSRT) is the most commonly employed field based measurement of CRF in children, is low cost, simple, and can test large groups simultaneously [7,12,13]. There is strong evidence the 20mSRT has criterion validity in children [12] and as a surveillance instrument could help identify populations with either low or high CRF (based upon international normative values for children [13]) and provide indicators that inform policy or intervention practice [7].

Physical fitness, including muscular power and CRF, increases from school year to school year [14]. Both BMI and CRF vary seasonally in children [15,16] with increases seen throughout the school year [16,17]. However, improvements in CRF do not continue and are often lost over the summer recess [16,17]. Despite sedentary time and PA both potentially affecting BMI and CRF [18-21], the loss of CRF over the summer is particularly surprising as PA levels often increase during the summer periods [21].

Socioeconomic status also influences BMI with children from more deprived areas having greater obesogenic growth trajectories than more affluent peers [22-24]. Yet, PA levels and CRF may explain the impact of deprivation levels upon BMI [23,24]. Children from more deprived areas often have greater barriers to PA in general potentially influencing their BMI and CRF [25]. However, it is unknown whether seasonal variation seen in CRF, including the typical loss over summer holiday periods, is also influenced by deprivation levels. It may be that children from more deprived areas have greater barriers to summer PA opportunities [25] and thus, though in general children's PA levels increase through the summer [21], this may not be the case for those in more deprived areas. Though there are currently programmes examining anthropometric measures in children in the UK, there is a lack of CRF

testing in primary schools. Considering the importance of CRF, recent scoping work highlighted that it can be measured in schools in a simple scalable way using the 20mSRT [9]. In light of this and the seasonal variation seen in CRF, in addition to the current lack of research examining the impact of deprivation upon CRF, the current study had the following aims:

1. To investigate the BMI and CRF in primary school pupils in North West England over a one-year period.
2. Examine the impact of school deprivation levels upon CRF and BMI.

METHODS

Participants

Participants were (n=409) children aged 9-10 (Mean±SD; Height 132±6cm, Weight 30.75±7.15kg; 222 males and 187 females) from 13 schools in North West England. Uptake to the testing at the individual level was 100% from children in the age groups tested across the schools involved. Participants were recruited through schools' involvement with the investigation, and schools were themselves recruited via National Teaching Schools in the local area and existing links with Premier Sport. All schools approached agreed to participate in the investigation.

Ethics and informed consent

As the result of guidance from our institutional ethics panel, the protocol of the present investigation deviated from existing procedures for governance of child measurement. The norm is for schools to opt in to an investigation and provide consent on behalf of pupils with parents provided the opportunity to opt their children out of the investigation. In the present investigation schools and parents/guardians were required to provide consent and opt in to the investigation. Institution ethics was granted for this study

by the ethics committee at Aberystwyth University.

Consent was obtained from schools and subsequently parents/guardians of all potential participants. Children were provided the opportunity to not participate in testing sessions if they did not wish to. This model was been developed through consultation with Public Health England and the Office for Standards in Education, Children's Services and Skills (Ofsted).

Confidentiality

All data were anonymised immediately following collection and stored on secure systems approved by Ofsted. Opt-out forms and paper files were stored in locked cabinets within schools, and electronic files stored on password protected computers, in both cases in accordance with the Ofsted information governance team. All files were destroyed upon completion of the investigation. Data were anonymised when transferred to the research team, and stored on password protected computers.

Measures

All testing was completed by *in-situ* service staff delivering curricular PE lessons and other extra-curricular physical activities in schools. All staff were employees of Premier Sport. This organisation was identified as being well placed to deliver the fitness testing described above on the scale necessary as they were already situated in the schools included in the study and delivering the 20m-SRT as part of their core business. Measurements were taken at four time points throughout a single calendar year: autumn term (October 2014), spring term (March 2015), summer term (June 2015), and the following autumn term (October 2015). All protocols and procedures were overseen by a member of the research team at the first time point, and thereafter random visits made to ensure standard protocols were followed. Further, delivery staff were

provided with standard electronic forms to record all data.

BMI: Height was measured to the nearest centimetre and weight measured to the nearest 0.1kg. Body Mass Index (BMI) was calculated from height and weight scores ($\text{kg}\cdot\text{m}^{-2}$). BMI z-scores specific to age and sex were calculated using UK 1990 growth reference data [26]. International Obesity Task Force (IOTF) classifications were then established [27].

Fitness: The 20mSRT was conducted involving 20m shuttles starting at an initial speed of $8.5 \text{ km}\cdot\text{h}^{-1}$ increasing by $0.5 \text{ km}\cdot\text{h}^{-1}$ each minute. An audible signal indicated when a participant should initiate each shuttle. The final shuttle was recorded once a participant failed to maintain the audible controlled pace or stopped due to volitional exhaustion. End speed ($\text{km}\cdot\text{h}^{-1}$) was determined by the final stage of the 20-mSRT based on the number of completed shuttles. The speed for the final completed stage was used as the end speed [28]. Age and sex specific z-scores were calculated using age global reference values for children [13].

Area-Level Deprivation: The area-level deprivation of each school was classified using the English Index Deprivation 2015 (EID) [29] obtained from the school post-code (zip-code). The EID is a multidimensional measure of relative deprivation for small areas in England. The UK Government ranks areas based on decile of EID score (from decile 1 representing areas with the highest level of deprivation to decile 10 - the lowest). Areas falling in EID deciles 1 and 2 are considered deprived with area-level deciles 9 and 10 considered non-deprived (affluent). Participants were grouped from "Most Deprived" (deciles 1-2; $n = 153$), "Mid Deprived" (deciles 3-8; $n = 124$), to "Least Deprived" (deciles 9-10; $n = 132$).

Data treatment.

Missing data resulted at some time points due to child sickness, injury, or absence from school on the day of testing.

Table 1. Number of participants in each IOTF grade at each of the four measurement points.

Grade	Autumn 2014		Spring 2015		Summer 2015		Autumn 2015	
	%	n	n	%	n	%	n	%
Underweight (BMI <18.5 kg.m ²)	8.1%	33	41	9.0%	35	8.6%	35	8.6%
Normal weight (BMI 18.5 to <25 kg.m ²)	68.9%	282	285	69.7%	286	69.9%	276	67.5%
Overweight (BMI 25 to <30 kg.m ²)	14.9%	61	60	14.7%	64	15.6%	70	17.1%
Obesity (BMI > 30 kg.m ²)	8.1%	33	23	5.6%	24	5.9%	28	6.8%

The last observation carried forward (LOCF) method was used to provide a full data set. LOCF was used if a participant had two or more data entries producing a total sample of 409 for analysis. LOCF can provide biased estimates where it is expected that dependent variables might change over time (producing conservative or liberal estimates dependent upon whether it is expected that improvement or reduction respectively will occur). However, as standardised z-scores were used based upon age appropriate normative data as the dependent variables for analysis, thus providing a stable variable over time, it was considered that LOCF was appropriate. The total number of missing points, interpolated points, and comparison between the groups before and after data treatment are presented in the supplementary material. Independent samples t-test indicated no statistical difference between the original data and LOCF data, therefore the 409 participants with full data were used in subsequent analysis.

Statistical Analysis.

Data were analysed using IBM SPSS Statistics 24 for Windows (IBM., Chicago, IL.). Significance was accepted at an α level of $p \leq 0.05$. Assumptions of sphericity were examined using Mauchly's Test of Sphericity. A two-way 'time' x 'group' ANOVA was used to compare main effects for 'time' (Autumn 2014, Spring 2015, Summer 2015, Autumn 2015), 'group' (Most Deprived, Mid

Deprived, Least Deprived), and 'time' x 'group' interaction effects. Effects by 'sex' were also examined in addition to 'time' x 'sex' and 'group' x 'sex' interactions. Where sphericity was violated a Greenhouse-Geisser correction used. Post hoc pairwise comparisons using bonferroni adjusted confidence intervals were performed using estimated marginal means.

RESULTS

Body Mass Index

IOTF classifications are reported in Table 1. The majority of participants were classified as normal weight. For BMI, ANOVA revealed a significant effect by 'time' ($F_{(2.393,971.538)} = 6.469$, $p = 0.001$), though no significant effect by 'group' ($F_{(2,406)} = 1.171$, $p = 0.311$). There was a significant 'time' x 'group' interaction effect ($F_{(4.786,971.538)} = 3.916$, $p = 0.002$), though no significant post hoc pairwise comparisons at any time point. BMI significantly decreased ($p = 0.015$) from autumn ($z = 0.336$ [CI 0.212 to 0.460]) to spring ($z = 0.252$ [CI 0.132 to 0.371]), and then significantly increased ($p = 0.010$) to summer ($z = 0.327$ [CI 0.207 to 0.447]). Thus there were no significant differences between the first and following autumn terms ($p > 0.999$). Figure 1 shows mean z-scores over time for BMI and figure 2 shows BMI z-scores by deprivation group. There was no significant between group effect by 'sex' for BMI ($F_{(1,403)} = 0.008$, $p = 0.930$) and there were no 'group' x 'sex' ($F_{(2,403)} = 0.139$, $p = 0.870$) or 'time' x 'sex'

($F_{(2.395,965.074)} = 0.536, p = 0.617$) interactions.

Cardiorespiratory Fitness

For CRF, ANOVA revealed a significant effect by 'time' ($F_{(2.868,1164.256)} = 41.043, p < 0.001$), a significant effect by 'group' ($F_{(2,406)} = 4.351, p = 0.013$), and a significant 'time' x 'group' interaction effect ($F_{(5.735,1164.256)} = 5.987, p < 0.001$). CRF significantly increased ($p < 0.001$) from autumn ($z = 0.091$ [CI -0.014 to 0.196]) to spring ($z = 0.492$ [CI 0.367 to 0.616]), no change ($p = 0.308$) into summer ($z = 0.411$ [CI 0.294 to 0.528]), and a significant decrease ($p < 0.001$) into the following autumn term ($z = 0.125$ [CI 0.021 to 0.230]). Thus there were no significant differences between the first and following autumn terms ($p > 0.999$). Figure 3 shows z-scores over time for CRF.

Post hoc pairwise comparisons revealed pupils in the Most Deprived group had significantly lower CRF compared with the Least Deprived group in the autumn ($p < 0.001$). In the spring and summer there were no differences between groups. However, in the following autumn pupils in the Most Deprived group had significantly lower CRF compared with the Least Deprived group ($p < 0.001$) and the Mid Deprived group ($p = 0.003$). Figure 4 shows CRF scores by deprivation group.

There was a significant between group effect by 'sex' for CRF ($F_{(1,403)} = 23.394, p < 0.001$). However, there were no 'group' x 'sex' ($F_{(2,403)} = 0.318, p = 0.728$) or 'time' x 'sex' ($F_{(2.871,965.074)} = 1.913, p = 0.128$).

Figure 1. Mean BMI z-scores and 95% CIs by time point for entire sample

Figure 2. Mean BMI z-scores and 95% CIs by time point grouped by deprivation level

Figure 3. Mean CRF z-scores and 95% CIs by time point for entire sample

Figure 4. Mean CRF z-scores and 95% CIs by time point grouped by deprivation level

DISCUSSION

Main Findings – BMI, CRF, and Deprivation Levels

The primary purpose of this investigation was to examine BMI and CRF of children over a one-year period including academic year and summer recess, and the influence of school deprivation levels upon this. CRF increased (+3.8% end speed for 20-mSRT) from autumn to spring and was maintained throughout the school year. However, CRF levels decreased (-2.3% end speed for 20-mSRT) over the summer holidays to a level similar to the previous autumn. Maturation may impact CRF however normative data indicate an annual increase in boys and girls 9-10 years of age [13] so was unlikely to have caused the variation seen. Further, climate correlates with 20mSRT performance with higher CRF in colder countries [7] though was also unlikely to have caused the variation as meteorological data of the nearest UK station suggests mean temperature remained <20°C over the summer [30]. CRF changes were compounded by school area deprivation level increasing towards spring and decreasing over summer in all deprivation groups, yet with schools in the poorest areas diverging from others post summer recess. The Most Deprived group had initially lower CRF suggesting there may be the greatest potential for improvement in these children, yet similarly the greatest potential for CRF loss over the summer. BMI followed a similar pattern with initial decreases followed by increases over the year such that BMI was similar at both the beginning and end of the academic year and summer recess. BMI however was relatively unaffected by deprivation level. Further, though CRF was affected by sex, there were no interactions for BMI or CRF by sex with either time of year, or deprivation level.

These findings may have implications for policy makers and those responsible for the health and wellbeing of children. Firstly, from a positive perspective, school based PA delivery

programmes may be positively impacting upon CRF of children during the academic year. Indeed, children from schools in all deprivation levels increased CRF. Such increases are not dissimilar from specific training to increase CRF in children (~ 4-5% [31]). Other work has reported CRF increases over the school year in Greek school children [16] and Carrel et al. [32] found a school based PA intervention in US children improved CRF over the school academic year. However, some studies show CRF plateaus over summer recess [31], and Carrel et al. [17] reported follow-up analysis that improvements from their intervention were eradicated during this period. Recent work on seasonal variation in PA levels suggest children are least active during the autumn and winter seasons [33], and PA levels increase consistently during the summer season [21]. It seems counterintuitive that children in the present study lost CRF over this period.

It appears moderate-vigorous PA rather than light PA is more strongly associated with sustained health, fitness, and wellbeing in children [1,3,34,35]. Indeed, a recent study reported reallocation of sedentary time to light or moderate PA was not associated with improved CRF whereas vigorous PA was [20]. Though PA tends to increase over the summer this is mainly due to increased moderate PA, whereas vigorous PA tends to vary far less [21]. Thus, building CRF during youth through vigorous PA should be a focus of school based and summer PA programs. Summer day camps may provide opportunities to increase vigorous PA by 15-18% [36] and as such could represent a potentially valuable intervention.

Although PA levels were not measured, a notable limitation, anecdotally parents of children in this study suggested accessible opportunities to keep children active are limited during school holidays, and often unaffordable or difficult to manage around working hours. This may be a barrier that is amplified for those from

deprived areas [25] and involved in the greater loss of CRF in the most deprived children. As such, future work should examine whether interventions addressing barriers such as accessibility and cost can facilitate uptake in the most deprived.

Limitations

Although all students consented to participate, we did not have complete data for all time points (see supplementary material). This was primarily due to sickness, injury, and school absence, though it is possible some children may have been absent because of awareness of the forthcoming testing and thus there may be some bias to estimates produced from LOCF data. PA levels were not measured in this study and so it is also not clear whether changes in CRF were a result of PA levels. Further, deprivation was considered on the basis of school area. Individual participant level data was not available for deprivation levels. Some participants may have been based in areas outside of the schools that represented differing EID levels. Future work should consider this and also other factors indicative of deprivation level such as eligibility for free school meals. Lastly, this research was conducted in the North West of England. Thus the generalizability to other schools across England is not clear. Future work should similarly examine seasonal change in CRF in other areas of the UK and worldwide.

CONCLUSION

Our data highlight that, although CRF increases throughout the academic year, reductions in children's CRF levels occur over the summer holiday period – an effect which is significantly more apparent among children from the most deprived schools and unaffected by children's sex. As such, these results highlight a need for interventions aimed at increasing children's PA levels, particularly vigorous PA, targeted towards the summer holidays to maintain the improvements produced during the school year.

ACKNOWLEDGEMENTS

The authors wish to thank Premier Sport for their funding and support in the delivery of this investigation. Specific thanks goes to Luke Miles, Danny Melling, John Goner, and Andy Heald without whom the data collection would not have been possible, and for their contribution to the conceptualisation of this manuscript. Further, we wish to acknowledge each of the schools that participated, the parents of the children for providing required consent, and of course the children themselves.

REFERENCES

1. Barker AR, Gracia-Marco L, Ruiz JR, et al. Physical activity, sedentary time, TV viewing, physical fitness and cardiovascular disease risk in adolescents: The HELENA study. *Int J Cardiol.* 2018; 254: 303-309 doi: 10.1016/j.ijcard.2017.11.080.
2. Sardinha LB, Marques A, Minderico C, et al. Longitudinal relationship between cardiorespiratory fitness and academic achievement. *Med Sci Sports Exerc.* 2016; 48(5): 839-844 doi: 10.1249/MSS.0000000000000830.
3. Oliveira T, Pizarro A, Costa M, et al. Cardiorespiratory fitness, but not physical activity, is associated with academic achievement in children and adolescents. *Ann Hum Biol.* 2017; 44(4): 309-315 doi: 10.1080/03014460.2017.1308010.
4. Ruiz JR, Castro-Piñero J, Artero EG, et al. Predictive Validity of Health-Related Fitness in Youth: A Systematic Review. *Br J Sports Med.* 2009; 43(12): 909-923. doi: 10.1136/bjsm.2008.056499
5. Malina RM, Physical activity and fitness: Pathways from childhood to adulthood. *Am J Hum Biol.* 2001;13: 162–172 doi:10.1002/1520-6300(200102/03)13:2<162::AID-AJHB1025>3.0.CO;2-T

6. Case A, Fertig A, Paxson C. The lasting impact of childhood health and circumstance. *J Health Econ.* 2005; 24(2): 365-89
7. Lang JJ, Tremblay MS, Léger L, Olds T, Tomkinson GR. International variability in 20 m shuttle run performance in children and youth: who are the fittest from a 50-country comparison? A systematic literature review with pooling of aggregate results. *Br J Sports Med.* 2018; 52: 276. doi: 10.1136/bjsports-2016-096224
8. Tomkinson GR, Olds TS. Secular changed in pediatric aerobic fitness test performance: The global picture. In: Tomkinson GR, Olds TS, editors. *Pediatric Fitness. Secular Trends and Geographic Variability.* Med Sport Sci: Basel, Karger; 2007, p. 46-66
<https://doi.org/10.1159/000101075>
9. Domone S, Mann S, Sandercock G, Wade M, Beedie C. A Method by Which to Assess the Scalability of Field-Based Fitness Tests of Cardiorespiratory Fitness Among Schoolchildren. *Sports Med.* 2016; 46(12): 1819-1831 doi: 10.1007/s40279-016-0553-6.
10. Department of Health. *On the State of the Public Health. The 2009 Annual Report of the Chief Medical Officer.* London, UK; Department of Health; 2010.
11. NHS Digital. National Child Measurement Programme [Internet]. NHS Digital; n.d. Available from: <http://content.digital.nhs.uk/NCMP>
12. Castro-Piñero J, Artero EG, España-Romero V, et al. Criterion-related validity of field-based fitness tests in youth: a systematic review. *Br J Sports Med.* 2010; 44: 934-943. Doi: 10.1136/bjsm.2009.058321
13. Tomkinson GR, Lang JJ, Tremblay MS, et al. International normative 20 m shuttle run values from 1 142 026 children and youth representing 50 countries. *Br J Sports Med* 2017; 51: 1545-1554. Doi: 10.1136/bjsports-2016-095987.
14. Caldwell HA, Proudfoot NA, King-Dowling S, Di Cristofaro NA, Cairney J, Timmons BW. Tracking of physical activity and fitness during the early years. *Apply Physiol Nutr Metab.* 2016; 41(5): 504-510 doi: 10.1139/apnm-2015-0338
15. Moreno JP, Johnston CA, Chen T, et al. Seasonal variability in weight change during elementary school. *Obesity.* 2015; 23(2): 422-428
16. Christodoulos AD, Flouris AD, Tokmakidis SP. Obesity and physical fitness of pre-adolescent children during the academic year and the summer period: effects of organized physical activity. *J Child Health Care* 2006; 10(3): 199-212
17. Carrel AL, Clark RR, Peterson S, Eickhoff J, Allen DB. School-based fitness changes are lost during the summer vacation. *Arch Pediatr Adolesc Med.* 2007; 161(6): 561-4. doi:10.1001/archpedi.161.6.561
18. Sandercock GR, Ogunleye AA. Independence of physical activity and screen time as predictors of cardiorespiratory fitness in youth. *Pediatr Res.* 2013; 73(5): 692-697 doi: 10.1038/pr.2013.37.
19. Júdice PB, Silva AM, Berria J, Petroski EL, Ekelund U, Sardinha LB. Sedentary patterns, physical activity and health-related physical fitness in youth: a cross-sectional study. *Int J Behav Nutr Phys Act.* 2017; 14(1): 25 doi: 10.1186/s12966-017-0481-3.
20. Santos DA, Marques A, Minderico CS, Ekelund U, Sardinha LB. A cross-sectional and prospective analyses of reallocating sedentary time to physical activity on childrens cardiorespiratory fitness. *J Sports Sci.* 2017; Epub ahead of print. doi: 10.1080/02640414.2017.1411176.

21. Kornides ML, Rimm EB, Chavarro JE, Gillman MW, Rosner B, Field AE. Seasonal variations in meeting physical activity recommendations and development of overweight during adolescence. *Child Obes* 2018; 14(1): 33-40
22. Nau C, Schwartz BS, Bandeen-Roche K, et al. Community socioeconomic deprivation and obesity trajectories in children using electronic health records. *Obesity* 2015; 23(1): 207-212
23. Nevill AM, Duncan MJ, Lahart I, Sandercock G. Modelling the association between weight status and social deprivation in English school children: Can physical activity and fitness affect the relationship? *Ann Hum Biol.* 2016; 43(6): 497-504.
24. Nevill AM, Duncan MJ, Lahart I, Sandercock G. Cardiorespiratory fitness and activity explains the obesity-deprivation relationship in children. *Health Promot Int* 2017; Epub ahead of print. Doi: 10.1093/heapro/daw106.
25. Eyre EL, Duncan MJ, Birch SL, Cox VM. Low socio-economic environmental determinants of children's physical activity in Coventry, UK: A qualitative study in parents. *Prev Med Rep* 2014; 1: 32-42
26. Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Dis Child.* 1995; 73(1): 25-9.
27. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ.* 2000; 320(7244): 1240-1243.
28. Léger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for aerobic fitness. *J Sports Sci.* 1988; 6(2): 93-101.
29. Department for Communities and Local Government. *The English Index of Multiple Deprivation (IMD) 2015 – Guidance*. London, UK: Department for Communities and Local Government; 2015. 7. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/464430/English_Index_of_Multiple_Deprivation_2015_-_Guidance.pdf
30. Met Office Web site [internet]. Exeter, UK: Met Office; (cited March 2018). Available from: <https://www.metoffice.gov.uk/pub/data/weather/uk/climate/stationdata/bradforddata.txt>
31. Baquet G, Van Praagh E, Berthoin S. Endurance training and aerobic fitness in young people. *Sports Med.* 2003; 33(15): 1127-1143.
32. Carrel AL, Clark RR, Peterson SE, Nemeth BA, Sullivan J, Allen DB. Improvement of fitness, body composition, and insulin sensitivity in overweight children in a school-based exercise program: a randomized, controlled study. *Arch Pediatr Adolesc Med.* 2005; 159(10): 963-968
33. Atkin AJ, Sharp SJ, Harrison F, Brage S, van Sluijs EMF. Seasonal variation in children's physical activity and sedentary time. *Med Sci Sports Exerc.* 2014; 48(3): 449-456 doi: 10.1249/MSS.0000000000000786
34. Gralla MH, McDonald SM, Breneman C, Beets MW, Moore JB. Associations of objectively measured vigorous physical activity with body composition, cardiorespiratory fitness, and cardiometabolic health in youth: A review. *Am J Lifestyle Med.* 2016; Epub ahead of print doi.org/10.1177/1559827615624417

35. Aires L, Silva P, Silva G, Santos MP, Ribeiro JC, Mota J. Intensity of physical activity, cardiorespiratory fitness, and body mass index in youth. *J Phys Act Health*. 2010; 7(1): 54-59
36. Beets MW, Weaver RG, Beighle A, Webster C, Pate RR. How physically active are children attending summer day camps? *J Phys Act Health* 2013; 10(6): 850-855