ASCL1 is differentially expressed in the tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab (Herceptin) is a monoclonal antibody targeting the extracellular domain of the human epidermal growth factor receptor 2 (HER2)¹ utilized for the treatment of adjuvant and metastatic breast cancer² in the United States and worldwide. We mined published microarray and gene expression data^{3,4} to discover in an unbiased manner the most significant transcriptional changes associated with trastuzumab treatment. We identified achaete-scute family bHLH transcription factor 1, encoded by ASCL1 among the genes most differentially expressed in the primary tumors of patients with breast cancer treated with trastuzumab. The primary tumors of breast cancer patients treated with trastuzumab expressed higher levels of ASCL1 messenger RNA than did patients not treated with trastuzumab demonstrating increased expression of a transcription whose expression is sufficient to re-program mouse embryonic fibroblasts into cells of the neuronal lineage⁵⁻⁹ as a transcriptional consequence of treatment with trastuzumab.

Keywords: trastuzumab, breast cancer, ASCL1, systems biology of breast cancer, targeted therapeutics in breast cancer.

1 The mechanism of action ascribed to trastuzumab function is thought to be binding of the F_{ab}
2 domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2
3 (HER2)¹, but the transcriptional consequences resulting from signaling downstream of trastuzumab
4 binding to HER2 is less well understood. Through unbiased, systems-level comparative transcriptome
analysis of published microarray and gene expression data^{3,4}, we report here that ASCL1 is a differentially
expressed gene in the tumors of trastuzumab-treated patients: patients treated with trastuzumab expressed
significantly higher levels of ASCL1 than did those of patients not treated with trastuzumab.

5 **Methods**

6 We used datasets GSE130788³, and GSE50948⁴ for this differential gene expression analysis of
7 the transcriptional response to trastuzumab treatment in patients with breast cancer. GSE130788 was
8 generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with
9 *n*=11 primary breast tumors from patients treated with trastuzumab and *n*=36 primary tumors from
10 patients treated with trastuzumab, analysis performed using platform GPL6480. GSE50948 was
11 generated using Affymetrix Human Genome U133 Plus 2.0 Array technology with *n*=93 primary tumors
12 of the breast from patients not treated with trastuzumab and *n*=63 primary tumors of the breast from
13 patients treated with trastuzumab; platform GPL570 was utilized for analysis. The Benjamini and
14 Hochberg method of *p*-value adjustment was used for ranking of differential expression but raw *p*-values
were used to assess statistical significance of global or multiplexed differential expression. Log-
transformation of data was auto-detected, and the NCBI generated category of platform annotation was
used. A statistical test was performed to evaluate whether ASCL1 gene expression was significantly
between primary breast tumors of patients receiving trastuzumab using a two-tailed, unpaired t-test with
Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human
breast cancer (Version 8.4.0)(455).

15 **Results**

16 We performed unbiased comparative transcriptome analysis using published microarray and gene
17 expression data^{3,4} to identify the most significant transcriptional features of the primary breast tumors of
18 patients treated with trastuzumab (Herceptin).

19 **ASCL1 is differentially expressed in the tumors of patients treated with trastuzumab.**

20 We identified achaete-scute family bHLH transcription factor 1, encoded by ASCL1 as among the
21 genes whose expression changed most significantly in the tumors of breast cancer patients treated with
22 trastuzumab in a multiplexed gene expression dataset³ (Table 1). When sorting each of the genes
23 expressed in the primary tumors of breast cancer patients treated with trastuzumab based on significance
of change as compared to that of patients not treated with trastuzumab, ASCL1 ranked 108 out of 41000
total transcripts (Table 1), equating to 99.7% differential expression. ASCL1 differential expression in the
tumors of patients treated with trastuzumab trended towards statistical significance (Table 1; *p*=1.92E-54).

24 We turned to a second microarray dataset to validate our findings of ASCL1 differential
25 expression in the patients treated with trastuzumab. We found again that ASCL1 was among the genes
26 most differentially expressed in the tumors of patients administered a single treatment of trastuzumab
27 (Table 2). When sorting each of the genes expressed in the tumors of patients administered a single
28 treatment of trastuzumab, ASCL1 ranked 4621 out of 54675 total transcripts (Table 2). Differential
expression of ASCL1 in the primary tumors of breast cancer patients treated with trastuzumab, though
equating to 91.5% differential expression, was not deemed statistically significant (Table 2; *p*=7.84E-02).

ASCL1 is expressed at significantly higher levels in the tumors of breast cancer patients treated with trastuzumab than in the tumors of those not treated with trastuzumab.

We obtained exact mRNA expression levels for ASCL1 from tumors of patients treated with trastuzumab and from the tumors of patients not treated with trastuzumab to directly evaluate direction and statistical significance of change in expression of ASCL1 based on history of trastuzumab treatment. ASCL1 was expressed at higher levels in the tumors of patients treated with trastuzumab as compared to those of patients not treated with trastuzumab, and this difference was deemed statistically significant (Figure 1; $p < 0.0001$).

Thus, we found using published multiplexed gene expression and microarray data^{3,4} that ASCL1 was among the genes most differentially expressed in the tumors of patients treated with trastuzumab, that ASCL1 was expressed at higher levels in the tumors of patients treated with trastuzumab as compared to those not treated with trastuzumab, though we were not able to discern given the data analyzed whether this transcriptional change was a direct result of trastuzumab treatment or a gene expression change associated with clonal selection and evolution of the tumor across the course of treatment.

Discussion

We provided evidence here that ASCL1 is a differentially expressed gene in the tumors of patients treated with trastuzumab, and expressed at significantly higher levels in the primary tumors of breast cancer patients treated with trastuzumab as compared to patients not treated with trastuzumab. Thus, in patients with breast with breast cancer, administration of trastuzumab, a treatment associated with a 34% rate of central nervous system metastasis¹⁰, results in increased primary tumor expression of a transcription factor whose expression is sufficient to induce re-programming of mouse embryonic fibroblasts into induced neuronal cells⁵⁻⁹.

References

1. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
2. Herceptin (Trastuzumab). Indications. <https://www.herceptin.com/hcp/indications.html>.
3. GSE130788. A Phase II Randomized Trial of Neoadjuvant Trastuzumab or Lapatinib or the Combination of Trastuzumab and Lapatinib, Followed by Docetaxel and Carboplatin with Trastuzumab and/or Lapatinib in Patients with HER2+ Breast Cancer. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788>. Hsiaowang Chen. UCLA, Department of Medicine. Santa Monica, CA.
4. Prat, A., Bianchini, G., Thomas, M., Belousov, A., Cheang, M.C., Koehler, A., Gómez, P., Semiglazov, V., Eiermann, W., Tjulandin, S. and Byakhov, M., 2014. based PAM50 subtype predictor identifies higher responses and improved survival outcomes in HER2-positive breast cancer in the NOAH study. *Clinical Cancer Research*, 20(2), pp.511-521.
5. Chanda, S., Ang, C.E., Davila, J., Pak, C., Mall, M., Lee, Q.Y., Ahlenius, H., Jung, S.W., Südhof, T.C. and Wernig, M., 2014. Generation of induced neuronal cells by the single reprogramming factor ASCL1. *Stem cell reports*, 3(2), pp.282-296.
6. Pattyn, A., Simplicio, N., van Doorninck, J.H., Goridis, C., Guillemot, F. and Brunet, J.F., 2004. Ascl1/Mash1 is required for the development of central serotonergic neurons. *Nature neuroscience*, 7(6), pp. 589-595.
7. Battiste, J., Helms, A.W., Kim, E.J., Savage, T.K., Lagace, D.C., Mandyam, C.D., Eisch, A.J., Miyoshi, G. and Johnson, J.E., 2007. Ascl1 defines sequentially generated lineage-restricted neuronal and oligodendrocyte precursor cells in the spinal cord. *Development*, 134(2), pp.285-293.
8. Raposo, A.A., Vasconcelos, F.F., Drechsel, D., Marie, C., Johnston, C., Dolle, D., Bithell, A., Gillotin, S., van den Berg, D.L., Ettwiller, L. and Flicek, P., 2015. Ascl1 coordinately regulates gene expression and the chromatin landscape during neurogenesis. *Cell reports*, 10(9), pp.1544-1556.
9. Aydin, B., Kakumanu, A., Rossillo, M., Moreno-Estellés, M., Garipler, G., Ringstad, N., Flames, N., Mahony, S. and Mazzoni, E.O., 2019. Proneural factors Ascl1 and Neurog2 contribute to neuronal subtype identities by establishing distinct chromatin landscapes. *Nature neuroscience*, 22(6), pp. 897-908.
10. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.

Rank	ID	p-value	t	B	Gene	Gene name
108	A_23_P25305	1.92E-54	21.756168	113.415568	ASCL1	achaete-scute family bHLH transcription factor 1

Table 1: ASCL1 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
4621	209985_s_at	7.84E-02	1.7716838	-4.1084	ASCL1	achaete-scute family bHLH transcription factor 1

Table 2: ASCL1 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, gene and gene name are listed in this chart.

Figure 1: ASCL1 is expressed at significantly higher levels in the primary tumors of patients treated with trastuzumab.

The mRNA expression level of ASCL1 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between ASCL1 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.