A new family of markers to characterize the human $\gamma\delta$ T cell lineage

Shahan Mamoor, MS

Abstract

Prospective isolation of $\gamma\delta$ T lymphocytes demands a comprehensive description of the molecules that distinguish T cells with $\gamma\delta$ T-cell receptors (TCRs) ($\gamma\delta$ T cells, or $T\gamma\delta$) from those with $\alpha\beta$ TCRs ($T\alpha\beta$). Here I describe some of the most differentially expressed genes in the $\gamma\delta$ T cell when compared to the developmentally proximal but lineage-distinct $T\alpha\beta$ CD4⁺ and CD8⁺ lymphocytes. These genes encode cluster of differentiation markers, transcription factors, cell surface receptors and non-coding RNAs. As hematopoietic stem cells (HSCs) have been prospectively isolated based on the analysis of differentially expressed genes (1), any combination of these molecules may potentially be used to isolate $T\gamma\delta$, perhaps even independent of the $\gamma\delta$ TCR. This description of the most striking identifying features of the $T\gamma\delta$ will be a resource for the isolation of a multi-potent common $\gamma\delta$ T-cell progenitor.

Keywords: $\gamma\delta$ T-cells, $T\gamma\delta$, $\alpha\beta$ T-cells, $T\alpha\beta$, differential gene expression analysis, transcriptome

Introduction

$\gamma\delta$ T-cells can be classified according to their anatomical site of residence (e.g., the skin), the cytokines they secrete (e.g., IL-17), by the $V\gamma$ and $V\delta$ chains they utilize at the $\gamma\delta$ TCR (e.g., $V\gamma 9V\delta 2$), or by the types of markers they express on their cell surface (2, 3, 4).

Prospective isolation of $T\gamma\delta$ to a high degree of homogeneity has not been achieved, nor has a $\gamma\delta$ -specific common ancestor been defined that is capable of multi-lineage $T\gamma\delta$ reconstitution: one that has reached a stage where $T\alpha\beta$ lineage choice is no longer an option but still capable of differentiating into the myriad varieties of $T\gamma\delta$ that humans possess. A long-standing unresolved question in the $\gamma\delta$ T-cell field is whether the presence of a $\gamma\delta$ TCR on the surface of a lymphocyte is the developmentally defining event of the $\gamma\delta$ T-cell lineage (and similarly, whether the presence of an $\alpha\beta$ TCR defines the $\alpha\beta$ lineage) (5, 6). The main arguments from multiple studies utilizing over-expression or deletion of $\gamma\delta/\alpha\beta$ -Tcr genes can be summarized by the instructive and stochastic models of $T\gamma\delta$ development (5, 6, 8, 9). The instructive model argues TCR expression by the developing T-lymphocyte developmentally defines the $T\gamma\delta$ cell, while the stochastic model argues that though the expression of the $\gamma\delta$ TCR occurs during a developmentally defined window, TCR signaling (at most) reinforces $T\gamma\delta$ lineage fate decisions but does not specify the $T\gamma\delta$ lineage. Still, it remains unclear whether the presence of a $\gamma\delta$ TCR absolutely restricts a T cell to the $\gamma\delta$ lineage (5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16).

If the presence of a $\gamma\delta$ TCR does not necessarily define the fate of a $\gamma\delta$ T-cell, or necessarily restrict a T cell to the $\gamma\delta$ lineage, then it follows that a $T\gamma\delta/T\gamma\delta$ progenitor might exist in some cellular state (in development or in homeostasis) without a $\gamma\delta$ TCR. Assessing the validity of the instructive model of $\gamma\delta$ T-cell development requires an assessment of whether cells that are otherwise phenotypically $\gamma\delta$ T-cells exist without a $\gamma\delta$ TCR, and thus whether a $\gamma\delta$ TCR by definition both identifies and developmentally defines the $\gamma\delta$ lineage. Unfortunately, this is not currently possible as all methods hitherto have utilized antibodies against the $\gamma\delta$ TCR for $\gamma\delta$ T-cell isolation. Unique, non-TCR markers are needed for isolation and prospective characterization of $\gamma\delta$ T-cells.

In addition to its use for the identification of common progenitor, a standard panel of markers that label $T\gamma\delta$ could facilitate resolution of $\gamma\delta$ T-cells into distinct developmental stage groups independent of $V\gamma/V\delta$ usage, IL-17/IFN γ production, or site of anatomical residence. Similarly, answering the question of whether a common $\gamma\delta$ T-progenitor with the capacity for multi-lineage $\gamma\delta$ T-cell reconstitution exists would be facilitated by descriptions of the transcriptional landscape of $\gamma\delta$ T-cells in comparative fashion.

I present here a series of more than thirty molecules expressed by the $\gamma\delta$ T-cell, identified by comparing the global gene expression profiles of $T\gamma\delta$ cells against two T-cell types that developmentally follow $\gamma\delta$ T-cell lineage instruction: CD4 $^{+}$ and CD8 $^{+}$ $\alpha\beta$ T-cells ($T\alpha\beta$ cells). In total, the differential expression of five transcription factors, six differentiation markers, nineteen cell surface receptors, two non-coding RNA and six cytokine/chemokine receptors is outlined. Some of these molecules have been described in relationship to the $\gamma\delta$ T-cell lineage but many are completely unexplored with respect to $T\gamma\delta$ cells.

Methods

Comparative Gene Expression Profiling

GEO (Gene Expression Omnibus) Dataset GSE47920 was utilized for this analysis. The GEO2R web application identifies the most differentially expressed transcripts in a given comparison by ranking the log-fold change of RNA expression value between samples when provided with a dataset. Log transformation of data was set to “Yes”, and the p-value, was set to “Benjamini-Hochberg”. The “Submitter supplied” category of platform annotation was used to identify gene/RNA names.

Statistics

Statistical analyses comparing the mean RNA expression value of each differentially expressed transcript from flow-cytometry sorted peripheral blood CD3⁺ T $\gamma\delta$ versus T $\alpha\beta$ CD4⁺ or CD8⁺, were performed using GraphPad Prism Version 8.0. A standard one-way ANOVA with multiple comparisons was performed for each differentially expressed transcript, comparing the mean expression RNA expression value of CD4⁺ T $\alpha\beta$ to T $\gamma\delta$ as well as CD8⁺ T $\alpha\beta$ to T $\gamma\delta$. P-values for each comparison are denoted on each graph.

Results

I. Cluster of Differentiation Markers

I identified the most differentially expressed genes between cells that express $\gamma\delta$ or $\alpha\beta$ TCRs by comparing the transcriptomes of $\gamma\delta$ T-cells (T $\gamma\delta$) to CD4⁺ and CD8⁺ single positive T-lymphocytes of the $\alpha\beta$ lineage, using a dataset from a study investigating chimeric antigen receptors (17). This analysis revealed the two hundred genes in the lymphocyte transcriptome whose expression was most significantly different between $\gamma\delta$ T-cells and CD4⁺ and CD8⁺ $\alpha\beta$ T-cells. Of these, five transcripts corresponded to “traditional” cluster of differentiation (CD) markers.

CD36, also known as fatty acid translocase (FAT) or glycoprotein IIIb/IV was initially identified as a receptor for oxidized low-density lipoprotein and later as a scavenger receptor that recognizes a variety of molecules including native lipoproteins and long-chain fatty acids (21, 22, 23). Its expression by bovine $\gamma\delta$ T cells has been described (25). CD36 functions in a TLR4/6 heterodimer to promote innate immune signaling in response to amyloid beta or oxidized lipoprotein (24). I found that CD36 was expressed at significantly higher levels in $\gamma\delta$ T-cell transcriptome than in CD4⁺ or CD8⁺ $\alpha\beta$ T-lymphocytes (Figure 1).

CD160, or BY55 is a negative regulator of CD4⁺ T-cells expressed by natural killer (NK) cells and on intestinal intraepithelial lymphocytes (18, 19). It has been described as a receptor for classical and non-classical MHC class I molecules (20). These analyses revealed CD160 to be among the most differentially expressed molecules in the $\gamma\delta$ T-cell transcriptome (Figure 1).

CD86 is a co-stimulatory molecule belonging to the B7 family (26). Dysregulated expression of CD86/B7.2 has been reported in lesions in patients with multiple sclerosis, macrophage of patients with inflammatory bowel disease and in the plasma of patients with systematic lupus erythematosus (27, 28, 29). A study that examined the expression of a panel of co-stimulatory molecules in peripheral blood mononuclear cells reported that while unstimulated $\gamma\delta$ T cells did not express CD86, stimulation of PBMC-derived $\gamma\delta$ T cells with isopentyl pyrophosphate and IL-2 induced CD86 expression (30). My findings are different and indicate that not only is CD86 expressed in purified $\gamma\delta$ T cells but that, in the unstimulated state it is expressed in significant amounts relative to single-positive (CD4⁺/CD8⁺) $\alpha\beta$ T-lymphocytes (Figure 1).

CD163 is a class B scavenger receptor that is highly expressed on activated macrophage and is the receptor for the hemoglobin-haptoglobin complex (31). Bovine $\gamma\delta$ T-cells express WC1 which is evolutionarily related to CD163 (32). A meta-analysis of peripheral T-cell lymphomas (PTCLs) reported that expression of CD163 is induced in PTCLs (33) but a role for CD163 in $\gamma\delta$ T-cell biology has not been thoroughly examined. CD163 is among the genes whose expression is most significantly higher in $\gamma\delta$ than in $CD4^+/CD8^+$ T $\alpha\beta$ (Figure 1).

CD1d is a lipid-presenting molecule, and has been studied to some extent by multiple groups as a ligand/ligand-presenting complex that interacts with the $\gamma\delta$ TCR on gut $V\delta 1^+$ $\gamma\delta$ T-cells (34, 35, 36, 37). However, these studies all describe CD1d as being expressed on the cell that interacts with the $\gamma\delta$ T-cell, not the $\gamma\delta$ T-cell itself. These data indicate that not only is CD1d expressed by $\gamma\delta$ but in significantly higher quantities than in $CD4^+$ or $CD8^+$ T $\alpha\beta$ (Figure 1).

II. Cell Surface Receptors

Leukocyte Ig-like receptors

Leukocyte Ig-like receptors (LILRs) were originally identified as HLA/MHC class I interactors but are now known to engage a variety of ligands and have both activating and inhibitory effects on cells that engage them (37, 38, 39). LILRB1 has been reported to be expressed by $V\delta 2^-$ $\gamma\delta$ T-cells, but a role for LILRA2, LILRA3, or LILRB3 in $\gamma\delta$ T-cell biology has not been described. This analysis revealed that LILRA2, LILRA3, LILRB1 and LILRB3 are among the most differentially transcribed genes when comparing the transcriptomes of $\gamma\delta$ T-cells to $CD4^+$ and $CD8^+$ T-cells (Figure 2).

The LILRA genes represent a new class of immunoglobulin-like receptors that are expressed at high levels in $\gamma\delta$ T-cells and that distinguish $\gamma\delta$ T-cells from their $\alpha\beta$ lineage counterparts.

Membrane spanning 4A domains subfamily

Membrane-spanning 4 domains, subfamily A is a group of tetraspannin-like proteins that each have unique expression patterns and cellular functions (41).

MS4A7 is not well-characterized but was reported to be expressed in B-lymphocyte cell lines (41).

MS4A4A is expressed in cells of the macrophage lineage, particularly M2 polarized macrophages. MS4A4A was recently reported to be important in cancer via dectin-1 activation and NK cell-mediated resistance to metastasis (40).

MS4A6A is expressed in monocytes, macrophage and microglia. A single nucleotide variant of MS4A6A is associated with risk of developing Alzheimer's Disease (43).

The expression of MS4A7, MS4A4A or MS4A6A in $\gamma\delta$ T-cells has not been described nor have the MS4A family been implicated in $\gamma\delta$ -T cell biology. I found that MS4A7, MS4A4A and MS4A6A were among the most differentially transcribed genes when comparing $\gamma\delta$ T-cells to single positive $\alpha\beta$ T-lymphocytes, at significantly higher levels than their $CD4^+$ and $CD8^+$ counterparts (Figure 2). MS4A receptors, along with LILRA/B, belong to a new family of receptors that positively distinguish the identity of the $\gamma\delta$ T-cell.

C-type lectin and lectin receptors

Clec7a (Dectin-1) is expressed in IL17+ $\gamma\delta$ T-cells at significantly lower levels relative to macrophage but comparable to that of dendritic cells (46).

Clec4e (also known as Mincle) is expressed by δ -T cells in the skin that produce IL-17 (dermal $\gamma\delta$ T17), but it is not required for the induction of these cells by *Corynebacterium* microbiota in the skin (47).

No direct examination of the role of Clec10A in $\gamma\delta$ -T cell development or biology has been performed, but Tcr δ ^{-/-} mice fail to up-regulate Clec10A and M2-polarize macrophage as wild-type mice do in response to ozone exposure (45).

Neither Clec7a/Dectin-1, Clec4e/Mincle nor Clec10A have been thoroughly examined with respect to their expression throughout $\gamma\delta$ T-cell development, nor with respect to their role in the development of $\gamma\delta$ T-cells. I found that Clec7a, Clec4e and Clec10A were all expressed in Ty δ and at levels significantly higher than in Ta β (Figure 2). These members of the C-type lectin receptor family are among the genes most differentially expressed by Ty δ when compared to CD4⁺ and CD8⁺ Ta β .

Siglec10 is an innate immune lectin-type inhibitory receptor that contains an ITIM, or immunoreceptor tyrosine-based inhibitory motif. Siglec10 functions as a T-cell inhibitory molecule by interacting with CD52 (92). It serves critical functions as a negative regulator of tissue-damage induced immune activation (93), and in tumor immunotherapy by its expression on tumor-associated macrophages and interaction with CD24 over-expressing breast and ovarian cancers (94). I found that Siglec10 is expressed at significantly higher levels in Ty δ cells than in Ta β cells, and is among the genes most differentially expressed in Ty δ cells when compared to Ta β cells (Figure 2).

Wnt receptors

Frizzled 2 (FZD2) and Frizzled 5 (FZD) are receptors of the Wnt family, a morphogenic signaling pathway that canonically signals through beta-catenin and the Tcf1/Lef transcription factors (48). The putative ligand for Fzd2 and Fzd5 is Wnt5a (49). Human peripheral blood mononuclear cells express Frizzled 5 (50), but a role for Fzd2 or Fzd5 in the development or functioning of Ty δ cells has not been established. I found that Fzd2 and Fzd5 are expressed at significantly higher levels in Ty δ cells than in single-positive Ta β cells (Figure 2). Wnt receptors Fzd2 and Fzd5 are among the genes most differentially expressed between Ta β cells and Ty δ cells.

Assorted Transmembrane Receptors

TMEM144 is an orphan 10 transmembrane family receptor and has not been described in relation to $\gamma\delta$ T cells (51). I found that TMEM144 is among the genes whose expression is most differentially increased when comparing Ty δ cells to single-positive Ta β cells (Figure 2).

FCAR encodes for the transmembrane receptor for IgA, I (also known as CD89). IgA binding to Fc α RI can transduce either inhibitory or activating signals depending on the intracellular molecules that are recruited to its immunoreceptor tyrosine-based activation motif, or ITAM (52, 53). By recruiting the Shp1 phosphatase to the ITAM on its intracellular signaling tail, Fc α RI is able to transmit an inhibitory signal downstream of what otherwise would be an IgA-mediated activation cascade (53). Fc α RI is expressed on a variety of myeloid cells, but its expression in the $\gamma\delta$ T-cell lineage has not been extensively reported (52, 53). I found that Fc α RI is expressed on Ty δ cells, at significantly higher levels than in CD4⁺ and CD8⁺ Ta β cells

(Figure 2). It is among the most differentially expressed genes in the $\gamma\delta$ cell transcriptome relative to the $T\alpha\beta$ transcriptome (Figure 2).

LRRC25 is a negative regulator of the NF-kappaB signaling pathway, accomplishing this by targeting RelA for autophagic degradation (54). LRRC25 can also negatively regulate RIG-I-like receptor (RLR)-induced type I interferon signaling by targeting RIG-I for autophagic degradation (55). I found that LRRC25 is expressed in $\gamma\delta$ cells and at significantly higher levels than in $T\alpha\beta$ cells (Figure 2). A role for LRRC25 in $\gamma\delta$ cell development or function has not yet been reported.

Sphingosine-1 phosphate receptor

Sphingosine is a unique 18-carbon lipid molecule that possesses innate signaling properties when phosphorylated as sphingosine-1 phosphate (S1P) (56). S1P is recognized by a family of sphingosine-1 phosphate receptors, or S1PRs.

S1PR5 is expressed in human fetal blood V γ 9V δ 2 $\gamma\delta$ T-cells and is among the genes whose log-fold expression was highest in V γ 9V δ 2 $\gamma\delta$ T cells when comparing the transcriptomes of fetal blood $\gamma\delta$ and $\alpha\beta$ T cells (56, 57). Similarly, I found that S1PR5 is among the genes most differentially expressed in $\gamma\delta$ T-cells when compared to single-positive $\alpha\beta$ CD4⁺ or CD8⁺ T-cells (Figure 2). A direct role for S1PR5 in $\gamma\delta$ -T cell development or function has not been described.

TREM1

Triggering receptor expressed on myeloid cells, or TREM1 is expressed on neutrophils and monocytes and functions in a pro-inflammatory manner by inducing secretion of IL-8, TNF- α and MCP-1 (monocyte chemotactic protein 1) when engaged (59). The peritoneal neutrophils of patients with sepsis increase the expression of TREM1, and blockade of TREM1 prevents LPS-induced shock in mice (59). One study reported that TREM1 was one of the genes whose expression was the highest in the ears of Tcr δ ^{-/-} mice when compared to wild-type counterparts (58). TREM1 has not been described as a marker for $\gamma\delta$ identity. I found that TREM1 is expressed at significantly higher levels in $\gamma\delta$ cells than in $T\alpha\beta$ cells (Figure 2).

Complement Receptor C5aR

The complement pathway is an evolutionarily conserved innate immune defense system that generates a signaling cascade similar to thrombin in that activation of the complement pathway leads to proteolytic cleavage events and subsequent downstream effector functions. It has been previously reported that C5aR is expressed by $\gamma\delta$ cells (60). In support of this data, I found C5aR to be a differentially expressed gene in $\gamma\delta$ cells, with significantly higher expression in $\gamma\delta$ than in CD4⁺ and CD8⁺ $T\alpha\beta$ (Figure 2).

III. Transcription Factors

Transcription factor KLF4 (kruppel-like factor 4) is one of four “Yamanaka factors” whose combined expression can convert adult somatic cells to embryonic stem cells (61). Cd4-Cre Klf4 knockout mice possess normal amounts and percentages of $\gamma\delta$ T-cells, but transgenic mice that over-express Klf4 are blocked at the DN2-DN3 transition and have a reduction in the number of $\gamma\delta$ T-cells (62, 63). Klf4 was among the genes most differentially expressed, and at higher levels, by $\gamma\delta$ T-cells when compared to CD4⁺ or CD8⁺ T-cells of the $\alpha\beta$ lineage (Figure 3).

TFEC (transcription factor EC), a member of the microphthalmia family, has been not described with respect to $\gamma\delta$ T-cell development or functions (64). Specific, IL-4-inducible

expression of TFEC in the macrophage lineage was previously reported but *Tfec*^{-/-} mice lack obvious myeloid cell phenotypes (65). These analyses indicate that TFEC is expressed in $\gamma\delta$ T-cells and at significantly higher levels in $\gamma\delta$ T-cells as compared to single-positive $\alpha\beta$ T-cells (Figure 3).

ZEB2 (zinc finger E box-binding homeobox 2) is expressed in CD8⁺ T cells and promotes maturation of CD8⁺ effector and memory T cell populations during infection (66). ZEB2 was also reported to be important for NK cell maturation and survival and its expression is regulated by T-bet (67). There is no literature investigating the role of ZEB2 in $\gamma\delta$ T-cell development or function. These analyses revealed ZEB2 as one of five transcription factors whose expression was most significantly higher in $\gamma\delta$ T-cells when compared to CD4⁺ and CD8⁺ T-cells (Figure 3).

The homeobox A10 transcription factor HOXA10 is expressed in primitive hematopoietic stem cells, and it is over-expressed in most myeloid leukemias but not in acute or chronic lymphoid leukemias (68, 69, 71). Over-expression of HOXA10 impairs erythroid and myeloid development and leads to a reduction in B-cell numbers in NOD/SCID mice that are repopulated with HOXA10 retrovirally-transduced cord blood/fetal liver-derived CD34⁺ hematopoietic stem progenitor cells (70). HOXA10 is also expressed in the human endometrium (72). There is no literature describing a role for HOXA10 in $\gamma\delta$ T-cell development or function. I found that HOXA10 is expressed at higher levels in $\gamma\delta$ T-cells than in CD4⁺ or CD8⁺ T-cells. Differential expression of HoxA10 is deemed statistically significant by GEO2R but a one-way ANOVA of expression values indicates that this differential expression is of less statistical significance than the other four transcription factors revealed by comparative transcriptome analyses (Figure 3).

MYCL is a basic helix-loop-helix leucine zipper transcription factor that can bind to E-box elements after heterodimerization with MAX (73). There is no literature describing a role for MYCL (L-myc) in $\gamma\delta$ -T cell development or function. Though c-Myc and n-Myc deficient mice die at embryonic days 10 and 12, respectively, l-myc deficient mice are viable and have no obvious phenotype (73). I found that MYCL was among the genes whose expression was most significantly different between $\gamma\delta$ T-cells and $\alpha\beta$ CD4⁺ and CD8⁺ T-cells, with significantly higher expression in $\gamma\delta$ T cells (Figure 3).

IV. Non-coding RNAs

LINC00484 is a linear, 454 bp long intergenic non-coding RNA (lincRNA) located on chromosome 9, also known as C9orf73. Its function is not well-characterized and there are no studies describing a relationship between LINC00484 and $\gamma\delta$. I found that LINC00484 was one of the most differentially expressed genes when comparing $\gamma\delta$ cells to $\alpha\beta$ cells, and expressed at significantly higher levels in $\gamma\delta$ cells (Figure 4).

LINC01234 (also known as onco-lincRNA-32) is a long intergenic non-coding RNA located on chromosome 12 that is expressed at higher levels in multiple cancers (74, 75, 76, 77, 78). Higher expression of LINC01234 has been reported to correlate with poorer survival in breast and colon cancer. LINC01234 has not been examined with respect to $\gamma\delta$ cell function or development. LINC01234 is expressed at significantly higher levels in $\gamma\delta$ cells than in CD4⁺/CD8⁺ single-positive $\alpha\beta$ cells, and is among the genes most differentially expressed when comparing $\gamma\delta$ cells to $\alpha\beta$ cells (Figure 4).

V. Chemokine and Cytokine Receptors and Ligands

Chemokine CC motif receptor like-2 (CCRL2) is a less-studied chemokine receptor that displays most homology to CCR1 (80). It plays a key role in IgE-mediated cutaneous

anaphylaxis through mast cells and also in trafficking of dendritic cells to the lung (80, 81). Ccr12 has not been investigated with respect to $\gamma\delta$ T-cell development or function. I found that Ccr12 was expressed at significantly higher levels in $\gamma\delta$ versus $\alpha\beta$ CD4⁺ or $\alpha\beta$ CD8⁺ and was among the genes whose expression was most different between $\gamma\delta$ and $\alpha\beta$ CD4⁺/CD8⁺ (Figure 5).

Chemokine (C-C) motif-like 4 (CCL4), also known macrophage inflammatory protein 1 β (MIP-1 β) displays specificity for the CCR5 chemokine receptor (82). CCL4 is expressed by a wide variety of immune cells including monocytes, neutrophils, as well as B and T cells (82, 83). CCL4 functions in the recruitment of regulatory T cells by antigen-presenting cells and B-cells, and can act in a chemoattractant fashion for monocytes and NK cells as well. It was reported that engagement of NKp30 on V δ 1 could trigger the production of CCL4 (82, 83). The potential role of CCL4/CCL4 signaling in the function and/or development of $\gamma\delta$ has not been thoroughly examined (Figure 5).

Chemokine (C-C) CCR1 is a receptor for the "regulation on activation T-cell expressed and secreted" chemokine RANTES. A study investigating the expression of chemokine receptors on $\gamma\delta$ found that bulk $\gamma\delta$ TCR⁺ peripheral blood cells express CCR1 at higher levels than $\alpha\beta$ TCR⁺ peripheral blood cells (84). I found that both CCL4 and CCR1 are expressed at significantly higher levels in $\gamma\delta$ cells than in single-positive CD4 or CD8 $\alpha\beta$ cells (Figure 5).

CSF2RA is the ligand-binding subunit for the granulocyte-macrophage colony-stimulating factor receptor (85, 86). It is expressed by almost all myeloid cells (85, 86). A role for CSF2RA in $\gamma\delta$ cell development or function has not been described. I found here that $\gamma\delta$ cells express significantly higher levels of CSF2RA than $\alpha\beta$ cells do, and that CSF2RA is one of the most differentially expressed genes when comparing the transcriptomes of $\gamma\delta$ cell and CD4⁺/CD8⁺ $\alpha\beta$ cells (Figure 5).

IL-1B is a pro-inflammatory cytokine that is the canonical downstream effector of the inflammasomes. $\gamma\delta$ cells have been documented to participate in inflammasome signaling in trans through their ability to produce IL-17 (87), and at least one group has reported the production of proIL-1 β by $\gamma\delta$ cells (92). I found here that IL-1 β is among the most differentially expressed genes when comparing the transcriptomes of $\gamma\delta$ cells to CD4⁺/CD8⁺ $\alpha\beta$ cells, and that $\gamma\delta$ cells express significantly higher levels of IL-1 β than $\alpha\beta$ cells do (Figure 5).

IL-18RAP, IL-18 receptor accessory protein (CDw218b) is a subunit of the interleukin-18 receptor (90). Single nucleotide polymorphisms the IL18rap gene are susceptibility loci for Crohn's Disease, celiac disease and leprosy (88, 89, 90). An IL18R α ⁺ $\gamma\delta$ cell population has previously been reported, and that study documented a higher expression of IL18RAP in CD8⁺ $\alpha\beta$ cells than in $\gamma\delta$ cells (91). On the contrary, these analyses revealed that IL18RAP is expressed at significantly higher levels in $\gamma\delta$ cells than in CD8⁺ and CD4⁺ $\alpha\beta$ cells (Figure 5).

Discussion

In this study I compared the transcriptional profiles of primary human CD3⁺ $\gamma\delta$ T-cells ($\gamma\delta$ cells) to CD4⁺ and CD8⁺ $\alpha\beta$ T-cells ($\alpha\beta$ cells). This global gene expression comparison study (mining control data from a study that investigated the transcriptional profile of chimeric antigen receptor-transduced cells) revealed a multitude of cell surface receptors, cluster of differentiation markers, transcription factors and two non-coding RNA molecules that are expressed by $\gamma\delta$ cells. Significantly, these $\gamma\delta$ cell-expressed transcripts represent the most

differentially expressed genes when comparing the transcriptional profiles of $\gamma\delta$ cells to $\alpha\beta$ cells. Previous studies such as the serial analysis of gene expression, or SAGE (96) are landmarks in global gene expression analyses of $\gamma\delta$, and others like the Immunological Consortium (11) have followed. These large-scale studies are powerful in that their systems-level assessment (e.g., principal component analysis) present a broad view of the transcriptional landscape within which $\gamma\delta$ T-cells operate. This study differs in that it utilizes one comparison between fewer samples, providing a more focused view of the most differentially expressed genes between $\gamma\delta$ and $\alpha\beta$ as opposed to a host of different cell types from multiple developmental stages.

Surprisingly, this global differential gene expression analysis revealed that CD1d is expressed by $\gamma\delta$ as opposed to (or, in addition to) being expressed on the antigen-presenting cells that interact with $\gamma\delta$. This challenges the prevailing view of CD1d function and merits experimental validation.

The $\gamma\delta$ cell differential expression of Wnt receptors Fzd2 and Fzd5, MS4A receptors, LILRA/B receptors, LRRC25 and non-coding RNAs LINC01234 and LINC00484 has not been reported previously.

The enriched expression of the new family of $\gamma\delta$ cell markers described here suggests that valuable insights can be gained by isolation of cells with antibodies that recognize any of these molecules. A “traditional” $\gamma\delta$ cell analysis could then be performed, ascertaining through flow cytometry what percentage of cells that have been isolated are $\gamma\delta$ TCR⁺. In this manner, an entire series of $\gamma\delta$ cell markers could be quantitatively indexed for their capacity to prospectively isolate some amount of cells that express the $\gamma\delta$ TCR.

Moreover, iterated analysis of a panel of cells that have been isolated without the use of $\gamma\delta$ TCR antibodies could be harnessed for transcriptional and functional evaluation of $\gamma\delta$ cell identity: the cytokines they secreted, the cell surface receptors they express, and the anatomical regions to which they home. More advanced methods utilizing GFP knock-in variants of selected markers would enable tracking of $\gamma\delta$ cells through development to more precisely and accurately address the question of whether the $\gamma\delta$ TCR instructs or simply reinforces $\gamma\delta$ cell lineage choice.

References

1. Kiel, M.J., Yilmaz, Ö.H., Iwashita, T., Yilmaz, O.H., Terhorst, C. and Morrison, S.J., 2005. SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell*, 121(7), pp.1109-1121.
2. Rothenberg, E.V., Moore, J.E. and Yui, M.A., 2008. Launching the T-cell-lineage developmental programme. *Nature Reviews Immunology*, 8(1), p.9.
3. Bonneville, M., O'Brien, R.L. and Born, W.K., 2010. $\gamma\delta$ T cell effector functions: a blend of innate programming and acquired plasticity. *Nature Reviews Immunology*, 10(7), p.467.
4. Ciofani, M. and Zúñiga-Pflücker, J.C., 2010. Determining $\gamma\delta$ versus $\alpha\beta$ T cell development. *Nature Reviews Immunology*, 10(9), p.657.
5. Mombaerts, P., Clarke, A.R., Rudnicki, M.A., Iacomini, J., Itohara, S., Lafaille, J.J., Wang, L., Ichikawa, Y., Jaenisch, R., Hooper, M.L. and Tonegawa, S., 1992. Mutations in T-cell antigen

- receptor genes α and β block thymocyte development at different stages. *Nature*, 360(6401), p.225.
6. Itohara, S., Mombaerts, P., Lafaille, J., Iacomini, J., Nelson, A., Clarke, A.R., Hooper, M.L., Farr, A. and Tonegawa, S., 1993. T cell receptor δ gene mutant mice: independent generation of $\alpha\beta$ T cells and programmed rearrangements of $\gamma\delta$ TCR genes. *Cell*, 72(3), pp. 337-348.
 7. Dave, V.P., Cao, Z., Browne, C., Alarcon, B., Fernandez-Miguel, G., Lafaille, J., de la Hera, A., Tonegawa, S. and Kappes, D.J., 1997. CD3 δ deficiency arrests development of the $\alpha\beta$ but not the $\gamma\delta$ T cell lineage. *The EMBO Journal*, 16(6), pp.1360-1370.
 8. Washburn, T., Schweighoffer, E., Gridley, T., Chang, D., Fowlkes, B.J., Cado, D. and Robey, E., 1997. Notch activity influences the $\alpha\beta$ versus $\gamma\delta$ T cell lineage decision. *Cell*, 88(6), pp. 833-843.
 9. Hayes, S.M., Li, L. and Love, P.E., 2005. TCR signal strength influences $\alpha\beta/\gamma\delta$ lineage fate. *Immunity*, 22(5), pp.583-593.
 10. Ciofani, M., Knowles, G.C., Wiest, D.L., von Boehmer, H. and Zúñiga-Pflücker, J.C., 2006. Stage-specific and differential notch dependency at the $\alpha\beta$ and $\gamma\delta$ T lineage bifurcation. *Immunity*, 25(1), pp.105-116.
 11. Narayan, K., Sylvia, K.E., Malhotra, N., Yin, C.C., Martens, G., Vallerskog, T., Kornfeld, H., Xiong, N., Cohen, N.R., Brenner, M.B. and Berg, L.J., 2012. Intrathymic programming of effector fates in three molecularly distinct $\gamma\delta$ T cell subtypes. *Nature immunology*, 13(5), p. 511.
 12. Coffey, F., Lee, S.Y., Buus, T.B., Lauritsen, J.P.H., Wong, G.W., Joachims, M.L., Thompson, L.F., Zúñiga-Pflücker, J.C., Kappes, D.J. and Wiest, D.L., 2014. The TCR ligand-inducible expression of CD73 marks $\gamma\delta$ lineage commitment and a metastable intermediate in effector specification. *Journal of Experimental Medicine*, 211(2), pp.329-343.
 13. Zarin, P., Wong, G.W., Mohtashami, M., Wiest, D.L. and Zúñiga-Pflücker, J.C., 2014. Enforcement of $\gamma\delta$ -lineage commitment by the pre-T-cell receptor in precursors with weak $\gamma\delta$ -TCR signals. *Proceedings of the National Academy of Sciences*, 111(15), pp.5658-5663.
 14. Sumaria, N., Grandjean, C.L., Silva-Santos, B. and Pennington, D.J., 2017. Strong TCR $\gamma\delta$ signaling prohibits thymic development of IL-17A-secreting $\gamma\delta$ T cells. *Cell reports*, 19(12), pp.2469-2476.
 15. In, T.S., Trotman-Grant, A., Fahl, S., Chen, E.L., Zarin, P., Moore, A.J., Wiest, D.L., Zúñiga-Pflücker, J.C. and Anderson, M.K., 2017. HEB is required for the specification of fetal IL-17-producing $\gamma\delta$ T cells. *Nature communications*, 8(1), p.2004.
 16. Lee, S.Y., Coffey, F., Fahl, S.P., Peri, S., Rhodes, M., Cai, K.Q., Carleton, M., Hedrick, S.M., Fehling, H.J., Zúñiga-Pflücker, J.C. and Kappes, D.J., 2014. Noncanonical mode of ERK action controls alternative $\alpha\beta$ and $\gamma\delta$ T cell lineage fates. *Immunity*, 41(6), pp.934-946.
 17. Themeli, M., Kloss, C.C., Ciriello, G., Fedorov, V.D., Perna, F., Gonen, M. and Sadelain, M., 2013. Generation of tumor-targeted human T lymphocytes from induced pluripotent stem cells for cancer therapy. *Nature biotechnology*, 31(10), p.928.

18. Cai, G., Anumanthan, A., Brown, J.A., Greenfield, E.A., Zhu, B. and Freeman, G.J., 2008. CD160 inhibits activation of human CD4+ T cells through interaction with herpesvirus entry mediator. *Nature immunology*, 9(2), p.176.
19. Anumanthan, A., Bensussan, A., Boumsell, L., Christ, A.D., Blumberg, R.S., Voss, S.D., Patel, A.T., Robertson, M.J., Nadler, L.M. and Freeman, G.J., 1998. Cloning of BY55, a novel Ig superfamily member expressed on NK cells, CTL, and intestinal intraepithelial lymphocytes. *The Journal of Immunology*, 161(6), pp.2780-2790.
20. Maeda, M., Carpenito, C., Russell, R.C., Dasanji, J., Veinotte, L.L., Ohta, H., Yamamura, T., Tan, R. and Takei, F., 2005. Murine CD160, Ig-like receptor on NK cells and NKT cells, recognizes classical and nonclassical MHC class I and regulates NK cell activation. *The Journal of Immunology*, 175(7), pp.4426-4432.
21. Endemann, G., Stanton, L.W., Madden, K.S., Bryant, C.M., White, R.T. and Protter, A.A., 1993. CD36 is a receptor for oxidized low density lipoprotein. *Journal of Biological Chemistry*, 268(16), pp.11811-11816.
22. Hoebe, K., Georgel, P., Rutschmann, S., Du, X., Mudd, S., Crozat, K., Sovath, S., Shamel, L., Hartung, T., Zähringer, U. and Beutler, B., 2005. CD36 is a sensor of diacylglycerides. *Nature*, 433(7025), p.523.
23. Aitman, T.J., Glazier, A.M., Wallace, C.A., Cooper, L.D., Norsworthy, P.J., Wahid, F.N., Al-Majali, K.M., Trembling, P.M., Mann, C.J., Shoulders, C.C. and Graf, D., 1999. Identification of Cd36 (Fat) as an insulin-resistance gene causing defective fatty acid and glucose metabolism in hypertensive rats. *Nature genetics*, 21(1), p.76.
24. Stewart, C.R., Stuart, L.M., Wilkinson, K., Van Gils, J.M., Deng, J., Halle, A., Rayner, K.J., Boyer, L., Zhong, R., Frazier, W.A. and Lacy-Hulbert, A., 2010. CD36 ligands promote sterile inflammation through assembly of a Toll-like receptor 4 and 6 heterodimer. *Nature immunology*, 11(2), p.155.
25. Lubick, K. and Jutila, M.A., 2006. LTA recognition by bovine $\gamma\delta$ T cells involves CD36. *Journal of leukocyte biology*, 79(6), pp.1268-1270.
26. Qureshi, O.S., Zheng, Y., Nakamura, K., Attridge, K., Manzotti, C., Schmidt, E.M., Baker, J., Jeffery, L.E., Kaur, S., Briggs, Z. and Hou, T.Z., 2011. Trans-endocytosis of CD80 and CD86: a molecular basis for the cell-extrinsic function of CTLA-4. *Science*, 332(6029), pp.600-603.
27. Windhagen, A., Newcombe, J., Dangond, F., Strand, C., Woodroffe, M.N., Cuzner, M.L. and Hafler, D.A., 1995. Expression of costimulatory molecules B7-1 (CD80), B7-2 (CD86), and interleukin 12 cytokine in multiple sclerosis lesions. *Journal of Experimental Medicine*, 182(6), pp.1985-1996.
28. Wong, C.K., Lit, L.C.W., Tam, L.S., Li, E.K. and Lam, C.W.K., 2005. Aberrant production of soluble costimulatory molecules CTLA-4, CD28, CD80 and CD86 in patients with systemic lupus erythematosus. *Rheumatology*, 44(8), pp.989-994.
29. Rugtveit, J., Bakka, A. and Brandtzaeg, P., 1997. Differential distribution of B7. 1 (CD80) and B7. 2 (CD86) costimulatory molecules on mucosal macrophage subsets in human inflammatory bowel disease (IBD). *Clinical & Experimental Immunology*, 110(1), pp.104-113.

30. Maniar, A., Zhang, X., Lin, W., Gastman, B.R., Pauza, C.D., Strome, S.E. and Chapoval, A.I., 2010. Human $\gamma\delta$ T lymphocytes induce robust NK cell-mediated antitumor cytotoxicity through CD137 engagement. *Blood*, 116(10), pp.1726-1733.
31. Fabriek, B.O., Dijkstra, C.D. and van den Berg, T.K., 2005. The macrophage scavenger receptor CD163. *Immunobiology*, 210(2-4), pp.153-160.
32. Herzig, C.T., Waters, R.W., Baldwin, C.L. and Telfer, J.C., 2010. Evolution of the CD163 family and its relationship to the bovine gamma delta T cell co-receptor WC1. *BMC evolutionary biology*, 10(1), p.181.
33. Pont, F., Familiades, J., Déjean, S., Fruchon, S., Cendron, D., Poupot, M., Poupot, R., L'Faqihi-Olive, F., Prade, N., Ycart, B. and Fournié, J.J., 2012. The gene expression profile of phosphoantigen-specific human $\gamma\delta$ T lymphocytes is a blend of $\alpha\beta$ T-cell and NK-cell signatures. *European journal of immunology*, 42(1), pp.228-240.
34. Luoma, A.M., Castro, C.D., Mayassi, T., Bembinster, L.A., Bai, L., Picard, D., Anderson, B., Scharf, L., Kung, J.E., Sibener, L.V. and Savage, P.B., 2013. Crystal structure of V δ 1 T cell receptor in complex with CD1d-sulfatide shows MHC-like recognition of a self-lipid by human $\gamma\delta$ T cells. *Immunity*, 39(6), pp.1032-1042.
35. Uldrich, A.P., Le Nours, J., Pellicci, D.G., Gherardin, N.A., McPherson, K.G., Lim, R.T., Patel, O., Beddoe, T., Gras, S., Rossjohn, J. and Godfrey, D.I., 2013. CD1d-lipid antigen recognition by the $\gamma\delta$ TCR. *Nature immunology*, 14(11), p.1137.
36. Benlagha, K. and Bendelac, A., 2000, December. CD1d-restricted mouse V α 14 and human V α 24 T cells: lymphocytes of innate immunity. In *Seminars in immunology* (Vol. 12, No. 6, pp. 537-542). Academic Press.
37. Lafarge, X., Pitard, V., Ravet, S., Roumanes, D., Halary, F., Dromer, C., Vivier, E., Paul, P., Moreau, J.F. and Déchanet-Merville, J., 2005. Expression of MHC class I receptors confers functional intraclonal heterogeneity to a reactive expansion of $\gamma\delta$ T cells. *European journal of immunology*, 35(6), pp.1896-1905.
38. Takeda, K. and Nakamura, A., 2017. Regulation of immune and neural function via leukocyte Ig-like receptors. *The Journal of Biochemistry*, 162(2), pp.73-80.
39. Burshtyn, D.N. and Morcos, C., 2016. The expanding spectrum of ligands for leukocyte Ig-like receptors. *The Journal of Immunology*, 196(3), pp.947-955.
40. Mattioli, I., Tomay, F., De Pizzol, M., Silva-Gomes, R., Savino, B., Gulic, T., Doni, A., Lonardi, S., Boutet, M.A., Nerviani, A. and Carrierio, R., 2019. The macrophage tetraspan MS4A4A enhances dectin-1-dependent NK cell-mediated resistance to metastasis. *Nature immunology*, 20(8), p.1012.
41. Eon Kuek, L., Leffler, M., Mackay, G.A. and Hulett, M.D., 2016. The MS4A family: counting past 1, 2 and 3. *Immunology and cell biology*, 94(1), pp.11-23.

42. Zuccolo, J., Deng, L., Unruh, T., Sanyal, R., Bau, J.A., Storek, J., Demetrick, D.J., Luider, J.M., Auer-Grzesiak, I., Mansoor, A. and Deans, J.P., 2013. Expression of MS4A and TMEM176 genes in human B lymphocytes. *Frontiers in immunology*, 4, p.195.
43. Hollingworth, P., Harold, D., Sims, R., Gerrish, A., Lambert, J.C., Carrasquillo, M.M., Abraham, R., Hamshere, M.L., Pahwa, J.S., Moskvin, V. and Dowzell, K., 2011. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nature genetics*, 43(5), p.429.
44. Howie, D., Nolan, K.F., Daley, S., Butterfield, E., Adams, E., Garcia-Rueda, H., Thompson, C., Saunders, N.J., Cobbold, S.P., Tone, Y. and Tone, M., 2009. MS4A4B is a GPCR-associated membrane adapter, expressed by regulatory T cells, which modulates T cell activation. *The Journal of Immunology*, 183(7), pp.4197-4204.
45. Mathews, J.A., Kasahara, D.I., Ribeiro, L., Wurmbrand, A.P., Ninin, F.M. and Shore, S.A., 2015. $\gamma\delta$ T cells are required for M2 macrophage polarization and resolution of ozone-induced pulmonary inflammation in mice. *PLoS One*, 10(7), p.e0131236.
46. Martin, B., Hirota, K., Cua, D.J., Stockinger, B. and Veldhoen, M., 2009. Interleukin-17-producing $\gamma\delta$ T cells selectively expand in response to pathogen products and environmental signals. *Immunity*, 31(2), pp.321-330.
47. Ridaura, V.K., Bouladoux, N., Claesen, J., Chen, Y.E., Byrd, A.L., Constantinides, M.G., Merrill, E.D., Tamoutounour, S., Fischbach, M.A. and Belkaid, Y., 2018. Contextual control of skin immunity and inflammation by *Corynebacterium*. *Journal of Experimental Medicine*, 215(3), pp.785-799.
48. Slusarski, D.C., Corces, V.G. and Moon, R.T., 1997. Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature*, 390(6658), p.410.
49. He, X., Saint-Jeannet, J.P., Wang, Y., Nathans, J., Dawid, I. and Varmus, H., 1997. A member of the Frizzled protein family mediating axis induction by Wnt-5A. *Science*, 275(5306), pp.1652-1654.
50. Blumenthal, A., Ehlers, S., Lauber, J., Buer, J., Lange, C., Goldmann, T., Heine, H., Brandt, E. and Reiling, N., 2006. The Wingless homolog WNT5A and its receptor Frizzled-5 regulate inflammatory responses of human mononuclear cells induced by microbial stimulation. *Blood*, 108(3), pp.965-973.
51. El-Heliebi, A., Kroneis, T., Wagner, K., Meditz, K., Kolb, D., Feichtinger, J., Thallinger, G.G., Quehenberger, F., Liegl-Atzwanger, B. and Rinner, B., 2014. Resolving tumor heterogeneity: genes involved in chordoma cell development identified by low-template analysis of morphologically distinct cells. *PloS one*, 9(2), p.e87663.
52. Bakema, J.E. and Van Egmond, M., 2011. The human immunoglobulin A Fc receptor Fc α RI: a multifaceted regulator of mucosal immunity. *Mucosal immunology*, 4(6), p.612.
53. Pasquier, B., Launay, P., Kanamaru, Y., Moura, I.C., Pfirsch, S., Ruffié, C., Hénin, D., Benhamou, M., Pretolani, M., Blank, U. and Monteiro, R.C., 2005. Identification of Fc α RI as

- an inhibitory receptor that controls inflammation: dual role of FcRγ ITAM. *Immunity*, 22(1), pp.31-42.
54. Feng, Y., Duan, T., Du, Y., Jin, S., Wang, M., Cui, J. and Wang, R.F., 2017. LRRC25 functions as an inhibitor of NF-κB signaling pathway by promoting p65/RelA for autophagic degradation. *Scientific reports*, 7(1), p.13448.
 55. Du, Y., Duan, T., Feng, Y., Liu, Q., Lin, M., Cui, J. and Wang, R.F., 2018. LRRC25 inhibits type I IFN signaling by targeting ISG15-associated RIG-I for autophagic degradation. *The EMBO journal*, 37(3), pp.351-366.
 56. Liu, Y., Wada, R., Yamashita, T., Mi, Y., Deng, C.X., Hobson, J.P., Rosenfeldt, H.M., Nava, V.E., Chae, S.S., Lee, M.J. and Liu, C.H., 2000. Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation. *The Journal of clinical investigation*, 106(8), pp.951-961.
 57. Mandala, S., Hajdu, R., Bergstrom, J., Quackenbush, E., Xie, J., Milligan, J., Thornton, R., Shei, G.J., Card, D., Keohane, C. and Rosenbach, M., 2002. Alteration of lymphocyte trafficking by sphingosine-1-phosphate receptor agonists. *Science*, 296(5566), pp.346-349.
 58. Girardi, Michael, Julia M. Lewis, Renata B. Filler, Adrian C. Hayday, and Robert E. Tigelaar. "Environmentally responsive and reversible regulation of epidermal barrier function by γδ T cells." *Journal of Investigative Dermatology* 126, no. 4 (2006): 808-814.
 59. Bouchon, A., Facchetti, F., Weigand, M.A. and Colonna, M., 2001. TREM-1 amplifies inflammation and is a crucial mediator of septic shock. *Nature*, 410(6832), p.1103.
 60. Han, G., Geng, S., Li, Y., Chen, G., Wang, R., Li, X., Ma, Y., Shen, B. and Li, Y., 2011. γδT-cell function in sepsis is modulated by C5a receptor signalling. *Immunology*, 133(3), pp. 340-349.
 61. Takahashi, K. and Yamanaka, S., 2006. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*, 126(4), pp.663-676.
 62. An, J., Golech, S., Klaewsongkram, J., Zhang, Y., Subedi, K., Huston, G.E., Wood III, W.H., Wersto, R.P., Becker, K.G., Swain, S.L. and Weng, N., 2011. Krüppel-like factor 4 (KLF4) directly regulates proliferation in thymocyte development and IL-17 expression during Th17 differentiation. *The FASEB Journal*, 25(10), pp.3634-3645.
 63. Wen, X., Liu, H., Xiao, G. and Liu, X., 2011. Downregulation of the transcription factor KLF4 is required for the lineage commitment of T cells. *Cell research*, 21(12), p.1701.
 64. Rehli, M., Lichanska, A., Cassady, A.I., Ostrowski, M.C. and Hume, D.A., 1999. TFEC is a macrophage-restricted member of the microphthalmia-TFE subfamily of basic helix-loop-helix leucine zipper transcription factors. *The Journal of Immunology*, 162(3), pp.1559-1565.
 65. Rehli, M., Sulzbacher, S., Pape, S., Ravasi, T., Wells, C.A., Heinz, S., Söllner, L., El Chartouni, C., Krause, S.W., Steingrimsson, E. and Hume, D.A., 2005. Transcription factor Tfec contributes to the IL-4-inducible expression of a small group of genes in mouse macrophages including the granulocyte colony-stimulating factor receptor. *The Journal of Immunology*, 174(11), pp.7111-7122.

66. van Helden, M.J., Goossens, S., Daussy, C., Mathieu, A.L., Faure, F., Marçais, A., Vandamme, N., Farla, N., Mayol, K., Viel, S. and Degouve, S., 2015. Terminal NK cell maturation is controlled by concerted actions of T-bet and Zeb2 and is essential for melanoma rejection. *Journal of Experimental Medicine*, 212(12).
67. Omilusik, K.D., Best, J.A., Yu, B., Goossens, S., Weidemann, A., Nguyen, J.V., Seuntjens, E., Stryjewska, A., Zweier, C., Roychoudhuri, R. and Gattinoni, L., 2015. Transcriptional repressor ZEB2 promotes terminal differentiation of CD8⁺ effector and memory T cell populations during infection. *Journal of Experimental Medicine*, 212(12), pp.2027-2039.
68. Buske, C., Feuring-Buske, M., Antonchuk, J., Rosten, P., Hogge, D.E., Eaves, C.J. and Humphries, R.K., 2001. Overexpression of HOXA10 perturbs human lymphomyelopoiesis in vitro and in vivo. *Blood*, 97(8), pp.2286-2292.
69. Lawrence, H.J., Sauvageau, G., Ahmadi, N., Lopez, A.R., LeBeau, M.M., Link, M., Humphries, K. and Largman, C., 1995. Stage-and lineage-specific expression of the HOXA10 homeobox gene in normal and leukemic hematopoietic cells. *Experimental hematology*, 23(11), pp.1160-1166.
70. Thorsteinsdottir, U., Sauvageau, G., Hough, M.R., Dragowska, W., Lansdorp, P.M., Lawrence, H.J., Largman, C. and Humphries, R.K., 1997. Overexpression of HOXA10 in murine hematopoietic cells perturbs both myeloid and lymphoid differentiation and leads to acute myeloid leukemia. *Molecular and cellular biology*, 17(1), pp.495-505.
71. Magnusson, M., Brun, A.C., Miyake, N., Larsson, J., Ehinger, M., Bjornsson, J.M., Wutz, A., Sigvardsson, M. and Karlsson, S., 2007. HOXA10 is a critical regulator for hematopoietic stem cells and erythroid/megakaryocyte development. *Blood*, 109(9), pp.3687-3696.
72. Taylor, H.S., Arici, A., Olive, D. and Igarashi, P., 1998. HOXA10 is expressed in response to sex steroids at the time of implantation in the human endometrium. *The Journal of clinical investigation*, 101(7), pp.1379-1384.
73. Hatton, K.S., Mahon, K., Chin, L., Chiu, F.C., Lee, H.W., Peng, D., Morgenbesser, S.D., Horner, J. and DePinho, R.A., 1996. Expression and activity of L-Myc in normal mouse development. *Molecular and cellular biology*, 16(4), pp.1794-1804.
74. Lin, C., Zhang, Y., Chen, Y. and Bai, Y., 2019. Long noncoding RNA LINC01234 promotes serine hydroxymethyltransferase 2 expression and proliferation by competitively binding miR-642a-5p in colon cancer. *Cell death & disease*, 10(2), p.137.
75. Chen, X., Chen, Z., Yu, S., Nie, F., Yan, S., Ma, P., Chen, Q., Wei, C., Fu, H., Xu, T. and Ren, S., 2018. Long noncoding RNA LINC01234 functions as a competing endogenous RNA to regulate CBFB expression by sponging miR-204-5p in gastric cancer. *Clinical Cancer Research*, 24(8), pp.2002-2014.
76. Lin, C., Zhang, Y., Chen, Y. and Bai, Y., 2019. Long noncoding RNA LINC01234 promotes serine hydroxymethyltransferase 2 expression and proliferation by competitively binding miR-642a-5p in colon cancer. *Cell death & disease*, 10(2), p.137.
77. Ghaffar, M., Khodahemmati, S., Li, J., Shahzad, M., Wang, M., Wang, Y., Li, C., Chen, S. and Zeng, Y., 2018. Long Non-coding RNA LINC01234 Regulates Proliferation, Invasion and Apoptosis in Esophageal Cancer Cells. *Journal of Cancer*, 9(22), p.4242.

78. Xie, J.J., Guo, Q.Y., Jin, J.Y. and Jin, D., 2019. SP1-mediated overexpression of lncRNA LINC01234 as a ceRNA facilitates non-small-cell lung cancer progression via regulating OTUB1. *Journal of cellular physiology*.
79. Dimova, T., Brouwer, M., Gosselin, F., Tassignon, J., Leo, O., Donner, C., Marchant, A. and Vermijlen, D., 2015. Effector V γ 9V δ 2 T cells dominate the human fetal $\gamma\delta$ T-cell repertoire. *Proceedings of the National Academy of Sciences*, 112(6), pp.E556-E565.
80. Zabel, B.A., Nakae, S., Zúñiga, L., Kim, J.Y., Ohyama, T., Alt, C., Pan, J., Suto, H., Soler, D., Allen, S.J. and Handel, T.M., 2008. Mast cell-expressed orphan receptor CCRL2 binds chemerin and is required for optimal induction of IgE-mediated passive cutaneous anaphylaxis. *Journal of Experimental Medicine*, 205(10), pp.2207-2220.
81. Otero, K., Vecchi, A., Hirsch, E., Kearley, J., Vermi, W., Del Prete, A., Gonzalvo-Feo, S., Garlanda, C., Azzolino, O., Salogni, L. and Lloyd, C.M., 2010. Nonredundant role of CCRL2 in lung dendritic cell trafficking. *Blood*, 116(16), pp.2942-2949.
82. Bystry, R.S., Aluvihare, V., Welch, K.A., Kallikourdis, M. and Betz, A.G., 2001. B cells and professional APCs recruit regulatory T cells via CCL4. *Nature immunology*, 2(12), p.1126.
83. Hudspeth, K., Fogli, M., Correia, D.V., Mikulak, J., Roberto, A., Della Bella, S., Silva-Santos, B. and Mavilio, D., 2012. Engagement of NKp30 on V δ 1 T cells induces the production of CCL3, CCL4, and CCL5 and suppresses HIV-1 replication. *Blood*, 119(17), pp.4013-4016.
84. Tagawa, T., Nishimura, H., Yajima, T., Hara, H., Kishihara, K., Matsuzaki, G., Yoshino, I., Maehara, Y. and Yoshikai, Y., 2004. V δ 1+ $\gamma\delta$ T cells producing CC chemokines may bridge a gap between neutrophils and macrophages in innate immunity during Escherichia coli infection in mice. *The Journal of Immunology*, 173(8), pp.5156-5164.
85. Glatzel, A., Wesch, D., Schiemann, F., Brandt, E., Janssen, O. and Kabelitz, D., 2002. Patterns of chemokine receptor expression on peripheral blood $\gamma\delta$ T lymphocytes: strong expression of CCR5 is a selective feature of V δ 2/V γ 9 $\gamma\delta$ T cells. *The Journal of Immunology*, 168(10), pp.4920-4929.
86. Croxford, A.L., Lanzinger, M., Hartmann, F.J., Schreiner, B., Mair, F., Pelczar, P., Clausen, B.E., Jung, S., Greter, M. and Becher, B., 2015. The cytokine GM-CSF drives the inflammatory signature of CCR2+ monocytes and licenses autoimmunity. *Immunity*, 43(3), pp.502-514.
87. Lalor, S.J., Dungan, L.S., Sutton, C.E., Basdeo, S.A., Fletcher, J.M. and Mills, K.H., 2011. Caspase-1-processed cytokines IL-1 β and IL-18 promote IL-17 production by $\gamma\delta$ and CD4 T cells that mediate autoimmunity. *The Journal of Immunology*, 186(10), pp.5738-5748.
88. Zhernakova, A., Festen, E.M., Franke, L., Trynka, G., van Diemen, C.C., Monsuur, A.J., Bevova, M., Nijmeijer, R.M., van't Slot, R., Heijmans, R. and Boezen, H.M., 2008. Genetic analysis of innate immunity in Crohn's disease and ulcerative colitis identifies two susceptibility loci harboring CARD9 and IL18RAP. *The American Journal of Human Genetics*, 82(5), pp.1202-1210.
89. Festen, E.A., Goyette, P., Green, T., Boucher, G., Beauchamp, C., Trynka, G., Dubois, P.C., Lagace, C., Stokkers, P.C., Hommes, D.W. and Barisani, D., 2011. A meta-analysis of genome-wide association scans identifies IL18RAP, PTPN2, TAGAP, and PUS10 as shared risk loci for Crohn's disease and celiac disease. *PLoS genetics*, 7(1), p.e1001283.

90. Liu, H., Irwanto, A., Tian, H., Fu, X.A., Yu, Y., Yu, G., Low, H., Chu, T., Li, Y., Shi, B. and Chen, M., 2012. Identification of IL18RAP/IL18R1 and IL12B as leprosy risk genes demonstrates shared pathogenesis between inflammation and infectious diseases. *The American Journal of Human Genetics*, 91(5), pp.935-941.
91. Fergusson, J.R., Smith, K.E., Fleming, V.M., Rajoriya, N., Newell, E.W., Simmons, R., Marchi, E., Björkander, S., Kang, Y.H., Swadling, L. and Kurioka, A., 2014. CD161 defines a transcriptional and functional phenotype across distinct human T cell lineages. *Cell reports*, 9(3), pp.1075-1088.
92. McCandless, E.E., Budde, M., Lees, J.R., Dorsey, D., Lyng, E. and Klein, R.S., 2009. IL-1R signaling within the central nervous system regulates CXCL12 expression at the blood-brain barrier and disease severity during experimental autoimmune encephalomyelitis. *The Journal of Immunology*, 183(1), pp.613-620.
93. Bandala-Sanchez, E., Zhang, Y., Reinwald, S., Dromey, J.A., Lee, B.H., Qian, J., Böhmer, R.M. and Harrison, L.C., 2013. T cell regulation mediated by interaction of soluble CD52 with the inhibitory receptor Siglec-10. *Nature immunology*, 14(7), p.741.
94. Chen, G.Y., Tang, J., Zheng, P. and Liu, Y., 2009. CD24 and Siglec-10 selectively repress tissue damage–induced immune responses. *Science*, 323(5922), pp.1722-1725.
95. Barkal, A.A., Brewer, R.E., Markovic, M., Kowarsky, M., Barkal, S.A., Zaro, B.W., Krishnan, V., Hatakeyama, J., Dorigo, O., Barkal, L.J. and Weissman, I.L., 2019. CD24 signalling through macrophage Siglec-10 is a target for cancer immunotherapy. *Nature*, p.1.
96. Shires, J., Theodoridis, E. and Hayday, A.C., 2001. Biological insights into TCR $\gamma\delta$ ⁺ and TCR $\alpha\beta$ ⁺ intraepithelial lymphocytes provided by serial analysis of gene expression (SAGE). *Immunity*, 15(3), pp.419-434.

Fig.1 : Expression of CD36, CD160, CD86, CD163 and CD1d distinguishes the $\gamma\delta$ T-cell lineage from the CD4/CD8 $\alpha\beta$ T cell lineage. Floating bars are plotted, grouped by cell subset with each differentially expressed gene displayed (log values). Bar graphs depict the mean RNA expression values for each individual gene in T $\gamma\delta$, T $\gamma\delta$ CD4+ and T $\gamma\delta$ CD8+ (linear values). The errors bars denote the standard deviation, with $n=2$ for each cell subset. P-values, as determined by one-way ANOVA, are shown with respect to T $\gamma\delta$.

Cell Surface Receptors

Figure 2: A new family of cell surface receptors expressed by T $\gamma\delta$ cells. Floating bars are plotted, grouped by cell subset with each differentially expressed gene displayed (log values). Bar graphs depict the mean RNA expression values for each individual gene in T $\gamma\delta$, T $\gamma\delta$ CD4⁺ and T $\gamma\delta$ CD8⁺ (linear values). The error bars denote the standard deviation, with $n=2$ for each cell subset. P-values, as determined by one-way ANOVA, are shown with respect to T $\gamma\delta$.

Figure 3: Differential expression of transcription factors Zeb2, Mycl, Tfec, Klf4 and Hoxa10 in the $\gamma\delta$ T-cell lineage. Floating bars are plotted, grouped by cell subset with each differentially expressed gene displayed (log values). Bar graphs depict the mean RNA expression values for each individual gene in $\gamma\delta$, $\gamma\delta$ CD4+ and $\gamma\delta$ CD8+ (linear values). The error bars denote the standard deviation, with $n=2$ for each cell subset. P-values, as determined by one-way ANOVA, are shown with respect to $\gamma\delta$.

Figure 4: Two non-coding RNAs, LINC01234 and LINC00484 are expressed in $\gamma\delta$ cells at high levels relative to $\alpha\beta$ cells. Floating bars are plotted, grouped by cell subset with each differentially expressed gene displayed (log values). Bar graphs depict the mean RNA expression values for each individual gene in $\gamma\delta$, $\gamma\delta$ CD4+ and $\gamma\delta$ CD8+ (linear values). The error bars denote the standard deviation, with $n=2$ for each cell subset. P-values, as determined by one-way ANOVA, are shown with respect to $\gamma\delta$.

Figure 5: Chemokine receptors CCRL2 and CCR1, chemokine CCL4, the alpha subunit of the GM-CSF receptor, IL-1 β and the IL-18 accessory protein are among the genes most differentially expressed in T $\gamma\delta$ cells and are expressed at significantly higher levels in T $\gamma\delta$ cells than in T $\alpha\beta$ cells. Floating bars are plotted, grouped by cell subset with each differentially expressed gene displayed (log values). Bar graphs depict the mean RNA expression values for each individual gene in T $\gamma\delta$, T $\gamma\delta$ CD4 $^{+}$ and T $\gamma\delta$ CD8 $^{+}$ (linear values). The error bars denote the standard deviation, with $n=2$ for each cell subset. P-values, as determined by one-way ANOVA, are shown with respect to T $\gamma\delta$.