TNFSF8 (CD153 / CD30L) is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab, a monoclonal antibody targeted against the human epidermal growth factor receptor 2 (HER2) is utilized for the treatment of human breast cancer^{1,2}, but a complete understanding of how tumor signal transduction is modulated by trastuzumab treatment is lacking. By mining published and public microarray and gene expression data^{3,4} from the primary tumors of patients treated with trastuzumab, we found that TNFSF8, also known as CD153 and CD30 ligand (CD30L) was among the genes most differentially expressed in the primary tumors of patients treated with trastuzumab. TNFSF8 messenger RNA expression was significantly enhanced in the primary tumors of patients treated with trastuzumab. Thus, trastuzumab treatment in patients with breast cancer is associated with increased expression, in primary tumors of the breast, of a marker with broad expression in multiple human cancers of the hematopoietic system^{5,6}, and with the capacity to regulate immunoglobulin class switching⁷.

Keywords: breast cancer, trastuzumab, TNFSF8, systems biology of breast cancer, targeted therapeutics in breast cancer.

The mechanism of action by which trastuzumab exerts its therapeutic function in human breast cancer is said to be binding of the F_{ab} domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2 (HER2)², possibly resulting in decreased cellular proliferation⁸. Transcriptional modulation resulting from trastuzumab binding events in the primary tumors of treated patients is less well understood. We mined published and public microarray and gene expression data^{3,4} generated using primary tissues from patients treated with trastuzumab to compare tumor global gene expression with that of patients not treated with trastuzumab. Using this unbiased, systems level methodology, we identified differential expression and up-regulation of TNFSF8 in primary tumors of the breast in patients treated with trastuzumab.

Methods

We used datasets GSE38057³ and GSE130788⁴ for this differential gene expression analysis of the transcriptional response to trastuzumab treatment in patients with breast cancer. GSE38057 was generated using DASL Human Cancer Panel by Probe technology with $n=6$ primary breast tumors from patients treated with induction chemotherapy and/or hormone therapy without trastuzumab and $n=35$ primary breast tumors from patients treated with induction chemotherapy and/or hormone therapy in addition to trastuzumab, both of which developed brain metastases; analysis was performed using platform GPL5826. GSE130788 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with $n=11$ primary breast tumors from patients treated with trastuzumab and $n=36$ primary tumors from patients treated with trastuzumab, analysis performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global or multiplexed differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether TNFSF8 gene expression was significantly between primary breast tumors of patients receiving trastuzumab using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

Results

We performed comparative transcriptome analysis of primary tumors from patients treated or not with trastuzumab (Herceptin), using published multiplexed gene expression data³ and a microarray dataset⁴, in order to identify the most significant transcriptional changes in the primary tumors of breast cancer patients following treatment with trastuzumab.

TNFSF8 is differentially expressed in the primary tumors of patients treated with trastuzumab.

We found that the tumor necrosis factor superfamily member 8, TNFSF8, also known as CD153 and CD30 ligand (CD30L), was among the genes whose expression was most significantly different in the primary tumors of patients with breast cancer treated with trastuzumab³. When sorting each of the genes expressed in trastuzumab-treated breast tumors based on significance of difference in expression as compared to the primary breast tumors of patients not treated with trastuzumab, TNFSF8 ranked 2 out of 1506 total transcripts, equating to 99.9% statistical significance (Table 1). Differential expression of TNFSF8 in the primary tumors of breast cancer patients treated with trastuzumab was trended towards statistical significance (Table 1; $p=0.00439$).

We queried a second dataset⁶ to validate our findings of TNFSF8 differential expression upon trastuzumab treatment. In a second microarray dataset analyzing gene expression of patients treated with trastuzumab, we again found that TNFSF8 was among the genes whose expression was most significantly

different transcriptome-wide when comparing the global gene expression profiles of tumors from patients treated with trastuzumab as compared to those not treated with trastuzumab. When sorting each of the genes expressed in the tumors of patients treated with trastuzumab based on significance of change as compared to the tumors of patients not treated with trastuzumab, TNFSF8 ranked 16926 out of 41000 total transcripts, equating to 58.7% differential expression. Differential expression of TNFSF8 in the primary tumors of patients treated with trastuzumab was statistically significant (Table 2; $p=3.41E-04$).

TNFSF8 is expressed at higher levels in the tumors of patients treated with trastuzumab.

We obtained exact mRNA expression levels for TNFSF8 from the tumors of patients treated with trastuzumab and from the tumors of patients not receiving trastuzumab to determine magnitude and direction of TNFSF8 change in the tumors of patients treated with trastuzumab. TNFSF8 expression was significantly higher in the tumors of patients treated with trastuzumab as compared to the tumors of patients not treated with trastuzumab (Figure 1). Increased expression of TNFSF8 trended was statistically significant (Figure 1: $p<0.0001$). We calculated a mean fold change of 2.23 ± 1.22 when comparing the expression of TNFSF8 in the tumors of patients treated with trastuzumab as compared to the tumors of patients treated with trastuzumab.

Thus, we found that TNFSF8 / CD153 / CD30L was among the genes most differentially expressed in the primary tumors of patients with breast cancer and that TNFSF8 mRNA was present at higher quantities in the primary tumors of patients treated with trastuzumab as compared to those not treated.

Discussion

We provided evidence here that TNFSF8, also known as CD153 and CD30L, is expressed at higher levels in the tumors of patients treated with trastuzumab and among the genes most differentially expressed in the primary tumors of patients treated with trastuzumab. We recently demonstrated that both TAL1 and LMO2 are transcriptionally induced in the primary tumors of breast cancer patients treated with trastuzumab^{8,9}. These data indicate that a marker frequently expressed in human myeloid and lymphoid malignancies^{5,6} is also transcriptionally induced in the primary tumors of breast cancer patients treated with trastuzumab, demonstrating that in the breast, transcriptional activation of two leukemogenic oncogenes by trastuzumab is accompanied by cell surface expression of a marker observed in cancers of the blood.

References

1. Slamon, D., Eiermann, W., Robert, N., Pienkowski, T., Martin, M., Press, M., Mackey, J., Glaspy, J., Chan, A., Pawlicki, M. and Pinter, T., 2011. Adjuvant trastuzumab in HER2-positive breast cancer. *New England Journal of Medicine*, 365(14), pp.1273-1283.
2. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
3. Duchnowska, R., Jassem, J., Goswami, C.P., Dundar, M., Gökmen-Polar, Y., Li, L., Woditschka, S., Biernat, W., Sosińska-Mielcarek, K., Czartoryska-Arłukowicz, B. and Radecka, B., 2015. Predicting early brain metastases based on clinicopathological factors and gene expression analysis in advanced HER2-positive breast cancer patients. *Journal of Neuro-oncology*, 122(1), pp.205-216.
4. GSE130788. A Phase II Randomized Trial of Neoadjuvant Trastuzumab or Lapatinib or the Combination of Trastuzumab and Lapatinib, Followed by Docetaxel and Carboplatin with Trastuzumab and/or Lapatinib in Patients with HER2+ Breast Cancer. [https:// www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788). Hsiaowang Chen. UCLA, Department of Medicine. Santa Monica, CA.
5. Gattei, V., Degan, M., Gloghini, A., De Iuliis, A., Improta, S., Rossi, F.M., Aldinucci, D., Perin, V., Serraino, D., Babare, R. and Zagonel, V., 1997. CD30 ligand is frequently expressed in human hematopoietic malignancies of myeloid and lymphoid origin. *Blood, The Journal of the American Society of Hematology*, 89(6), pp.2048-2059.
6. Gruss, H.J., DaSilva, N., Hu, Z.B., Uphoff, C.C., Goodwin, R.G. and Drexler, H.G., 1994. Expression and regulation of CD30 ligand and CD30 in human leukemia-lymphoma cell lines. *Leukemia*, 8(12), p. 2083.
7. Cerutti, A., Schaffer, A., Goodwin, R.G., Shah, S., Zan, H., Ely, S. and Casali, P., 2000. Engagement of CD153 (CD30 ligand) by CD30+ T cells inhibits class switch DNA recombination and antibody production in human IgD+ IgM+ B cells. *The Journal of Immunology*, 165(2), pp.786-794.
8. Mamoor, S., The LMO2 oncogene is up-regulated in the primary tumors of breast cancer patients treated with trastuzumab.
9. Mamoor, S., TAL1 is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

Rank	ID	p-value	t	B	FC	Gene
2	GI_24119162-S-3	0.00439	3.007632	-4.02	2.23 ± 1.22	TNFSF8 (CD153)

Table 1: TNFSF8 (CD30L / CD153) is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in TNFSF8 between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
16926	A_23_P169257	3.41E-04	-3.644954	-1.165619	TNFSF8	tumor necrosis factor superfamily member 8

Table 2: TNFSF8 (CD153 / CD30L) is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in TNFSF8 between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Figure 1: TNFSF8 is expressed at higher levels in the primary tumors of patients treated with trastuzumab as compared to those not treated with trastuzumab.

The mRNA expression level of TNFSF8 is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between TNFSF8 expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.