Recurrent differential expression of open reading frames on chromosome 14 in human brain metastatic breast cancer.

Shahan Mamoor, MS¹
¹shahanmamoor@gmail.com
East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We leveraged the power of systems-level comparative transcriptome analysis in human tissues to discover genes associated with brain metastasis in human breast cancer by integration of multiple independent published microarray datasets⁴⁻⁷. We identified five independent open reading frames on chromosome 14 that were among the most differentially expressed, transcriptome-wide, in the brain metastatic tissues of patients with metastatic breast cancer when compared to primary tumors of the breast: C14orf179, C14orf112, C14orf169, C14orf109 and C14orf130⁴. Importantly, primary tumor expression of each of these open reading frames was significantly correlated with either overall, distant metastasis-free, or recurrence-free survival in patients with breast cancer. Analysis of three separate microarray datasets⁵⁻⁷ revealed at least five other open reading frames on chromosome 14 that were recurrently transcriptionally modulated in brain metastatic tissues of humans with metastatic breast cancer as evidenced by transcriptome-wide differential expression relative to primary tumors of the breast. It is possible that structural or epigenetic changes on chromosome 14 are a major feature associated with human brain metastatic breast cancer.

Keywords: breast cancer, metastatic breast cancer, metastasis, brain metastasis, central nervous system metastasis, chromosome 14, C14orf179, C14orf112, C14orf169, C14orf109, C14orf130, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁸ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁸. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁸ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment-related or not. We performed unbiased systems-level discovery of genes associated with brain metastatic breast cancer using global comparative analysis of primary and metastatic patient tumor tissues and report here the recurrent, differential and increased expression of multiple open reading frames on chromosome 14 in human brain metastatic breast cancer.

Methods

We used datasets GSE10893⁴, GSE124648⁵, GSE125989⁶ and GSE52604⁷ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=11$ primary breast tumors and $n=3$ or $n=4$ brain metastases from patients with breast cancer; analysis was performed using platform GPL885. GSE124648 was generated using Affymetrix Human Genome U133A array technology with $n=10$ normal breast tissues and $n=44$ lymph node metastases from patients with breast cancer; analysis was performed using platform GPL96. GSE125989 was generated using Affymetrix Human Genome U133A 2.0 array technology with $n=16$ primary breast tumors and $n=16$ brain metastases from patients with breast cancer; the samples utilized by investigators for this microarray dataset were paired, matched tumor and metastasis samples; analysis was performed using platform GPL571. GSE52604 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer; analysis was performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether C14orf gene expression was significantly different between primary tumors of the breast and brain metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁹ for correlation of C14orf mRNA expression levels in primary tumors of the breast with overall survival in $n=1402$ breast cancer patients, with distant metastasis-free survival in $n=1803$ breast cancer patients, and recurrence-free survival in $n=3951$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic and primary tumor tissues of patients with metastatic breast, as well as normal tissues of the breast using

published microarray data⁴⁻⁷ to describe the transcriptional landscape of brain metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

Multiple open reading frames on chromosome 14 are differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified open reading frame 179 on chromosome 14, encoded by C14orf179, as a differentially expressed gene in the brain metastatic tissues of humans with breast cancer (Chart 1). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, C14orf179 ranked 39 out of 17418 total transcripts (Chart 1), equating to 99.8% differential expression. Differential expression of C14orf179 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=7.43E-05$). Four further open reading frames on chromosome 14 were among those most differentially expressed transcriptome-wide in the brain metastases of patients with brain metastatic breast cancer when compared to primary tumors of the breast (Chart 1). These included C14orf112, C14orf169, C14orf109, and C14orf130.

When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, C14orf112 ranked 46 out of 17418 total transcripts (Chart 1), equating to 99.7% differential expression. Differential expression of C14orf112 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=1.05E-04$).

When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, C14orf169 ranked 66 out of 17418 total transcripts (Chart 1), equating to 99.6% differential expression. Differential expression of C14orf169 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=1.63E-04$).

When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, C14orf109 ranked 190 out of 17418 total transcripts (Chart 1), equating to 98.9% differential expression. Differential expression of C14orf109 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=8.23E-04$).

When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, C14orf130 ranked 227 out of 17418 total transcripts (Chart 1), equating to 98.7% differential expression. Differential expression of C14orf130 in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=6.71E-04$).

Thus, we identified five separate open-reading frames on chromosome 14 as among the genes whose expression was most quantitatively different, transcriptome-wide, when comparing brain metastases to primary tumors of the breast.

Open reading frames on chromosome 14 are differentially expressed in the lymph node metastases of patients with breast cancer.

To determine whether differential expression of open reading frames on chromosome 14 in human breast cancer was specific to brain metastases or could be observed in other anatomical locations, we queried a second microarray dataset⁵, here comparing normal breast tissues to lymph node metastases. Again, we identified chromosome 14 open reading frames

as differentially expressed gene in the lymph node metastatic tissues of patients with breast cancer (Chart 2): these included C14orf132, C14orf1, C14orf2, C14orf169, and C14orf93.

When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, C14orf132 ranked 1408 out of 22283 total transcripts (Chart 2), equating to 93.7% differential expression. Differential expression of C14orf132 in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=8.59E-06$).

When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, C14orf1 ranked 3775 out of 22283 total transcripts (Chart 2), equating to 83.1% differential expression. Differential expression of C14orf1 in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=1.81E-03$).

When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, C14orf2 ranked 4944 out of 22283 total transcripts (Chart 2), equating to 77.8% differential expression. Differential expression of C14orf2 in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=6.40E-03$).

When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, C14orf169 ranked 5625 out of 22283 total transcripts (Chart 2), equating to 74.8% differential expression. Differential expression of C14orf169 in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=1.07E-02$).

When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, C14orf93 ranked 8673 out of 22283 total transcripts (Chart 2), equating to 61.1% differential expression. Differential expression of C14orf93 in the lymph node metastases of patients with metastatic breast cancer was trended towards statistical significance (Chart 2; $p=6.13E-02$).

Thus, transcriptome-wide differential expression of open reading frames on chromosome 14, transcriptome-wide, in metastatic tissues of women with metastatic breast cancer was conserved across two independent microarray datasets, in the brain and in the lymph nodes.

Open reading frames on chromosome 14 differentially expressed in the lymph node metastases of patients with breast cancer are also differentially expressed in the brain metastases of patients with breast cancer.

To determine whether we could validate differential expression of open reading frames on chromosome 14 in human breast cancer, we queried two further microarray datasets^{6,7}, here comparing primary tumors of the breast to brain metastases. We blindly identified three of five open reading frames differentially expressed in the lymph node metastases of patients with breast cancer as also differentially expressed in the brain metastases of patients with breast cancer. These included C14orf132, C14orf1, and C14orf2 (Chart 3). In addition, we observed differential expression of C14orf159 (Chart 3). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast, C14orf132 ranked 1964 out of 22277 total transcripts (Chart 3), equating to 91.2% differential expression. Differential expression of C14orf132 in the brain metastases of patients with metastatic breast cancer trended towards statistical significance (Chart 3; $p=5.74E-02$).

1 When sorting each of the genes expressed in brain metastases based on significance of
2 difference as compared to primary tumors of the breast, C14orf1 ranked 2449 out of 22277 total
3 transcripts (Chart 3), equating to 89.01% differential expression. Differential expression of
4 C14orf1 in the brain metastases of patients with metastatic breast cancer trended towards
statistical significance (Chart 3; $p=7.54E-02$).

5 When sorting each of the genes expressed in brain metastases based on significance of
6 difference as compared to primary tumors of the breast, C14orf2 ranked 2467 out of 22277 total
7 transcripts (Chart 3), equating to 88.9% differential expression. Differential expression of
8 C14orf2 in the brain metastases of patients with metastatic breast cancer trended towards
9 statistical significance (Chart 3; $p=7.60E-02$).

10 When sorting each of the genes expressed in brain metastases based on significance of
11 difference as compared to primary tumors of the breast, C14orf159 ranked 5867 out of 22277
12 total transcripts (Chart 3), equating to 73.7% differential expression. Differential expression of
13 C14orf159 in the brain metastases of patients with metastatic breast cancer was not deemed
14 statistically significant (Chart 3; $p=2.19E-01$).

15 Differential gene expression analyses that utilize primary and metastatic tumor tissues to
16 discover genes associated with metastasis can be challenging due to the relatively heterogenous
17 cellular composition and constitution of the tumor as compared to normal, untransformed
18 (benign) tissues. Analysis of a fourth microarray dataset revealed that C14orf159, C14orf1,
19 C14orf93, C14orf169, and C14orf2 were again differentially expressed in the brain metastatic
20 tissues in independent patient tissues (Chart 4), here when comparing brain metastases to normal
21 (benign) tissues of the breast. When sorting each of the genes expressed in brain metastases
22 based on significance of difference as compared to normal breast tissues, C14orf159 ranked 5936
23 out of 41093 total transcripts (Chart 4), equating to 85.6% differential expression. Differential
24 expression of C14orf159 in the brain metastases of patients with metastatic breast cancer was
25 statistically significant (Chart 4; $p=4.79E-04$).

26 When sorting each of the genes expressed in brain metastases based on significance of
27 difference as compared to normal breast tissues, C14orf1 ranked 7673 out of 41093 total
28 transcripts (Chart 4), equating to 81.3% differential expression. Differential expression of
C14orf1 in the brain metastases of patients with metastatic breast cancer was statistically
significant (Chart 4; $p=1.50E-03$).

When sorting each of the genes expressed in brain metastases based on significance of
difference as compared to normal breast tissues, C14orf93 ranked 12263 out of 41093 total
transcripts (Chart 4), equating to 70.2% differential expression. Differential expression of
C14orf93 in the brain metastases of patients with metastatic breast cancer was statistically
significant (Chart 4; $p=1.22E-02$).

When sorting each of the genes expressed in brain metastases based on significance of
difference as compared to normal breast tissues, C14orf93 ranked 14115 out of 41093 total
transcripts (Chart 4), equating to 65.7% differential expression. Differential expression of
C14orf93 in the brain metastases of patients with metastatic breast cancer was statistically
significant (Chart 4; $p=2.20E-02$).

When sorting each of the genes expressed in brain metastases based on significance of
difference as compared to normal breast tissues, C14orf93 ranked 14198 out of 41093 total
transcripts (Chart 4), equating to 65.4% differential expression. Differential expression of
C14orf93 in the brain metastases of patients with metastatic breast cancer was statistically
significant (Chart 4; $p=2.25E-02$).

Thus, we were able to validate transcriptome-wide differential expression of multiple open reading frames on chromosome 14 in the brain metastatic tissues of patients with metastatic breast cancer when compared to primary tumors of the breast, in independently published microarray datasets^{6,7}.

Expression of differentially expressed ORFs on chromosome 14 is up-regulated in the brain metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for C14orfs in primary tumors of the breast and in brain metastasis of patients with brain metastatic breast cancer to determine direction and statistical significance of change in C14orf expression in brain metastatic tissues. C14orf112 was expressed at higher levels in the brain metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 1; $p=0.0001$). C14orf169 was expressed at higher levels in the brain metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 2; $p=0.00142$). C14orf109 was expressed at higher levels in the brain metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 3; $p=0.000183$). C14orf179 was expressed at higher levels in the brain metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 4; $p=0.001325$). C14orf130 was expressed at higher levels in the brain metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 5; $p=0.001585$).

Expression of open reading frames on chromosome 14 is significantly correlated with survival outcomes in human breast cancer.

We performed Kaplan-Meier survival analysis⁹ in to evaluate whether C14orf primary tumor expression was correlated with survival outcomes in human breast cancer, in 3951 breast cancer patients for recurrence-free survival, in 1803 breast cancer patients for distant metastasis-free survival, and 1402 breast cancer patients for overall survival. We observed a statistically significant correlation between primary tumor expression of C14orf169 primary tumor expression and distant metastasis-free survival (DMFS) in patients with breast cancer (Figure 6). Patients whose primary tumors expressed low levels of C14orf169 possessed median DMFS of 78.64 months, while patients whose tumors expressed high levels of C14orf169 possessed median DMFS of 125.65 months. This difference in DMFS based on C14orf169 tumor expression in patients with breast cancer was statistically significant (Figure 6, Chart 5; logrank p -value: 0.019; hazard ratio: 0.79 (0.66-0.96)) (Chart 5).

We also observed a statistically significant correlation between primary tumor expression of C14orf169 and distant metastasis-free survival in patients with breast cancer (Figure 7, Chart 6; logrank p -value: 0.0043; hazard ratio: 0.76 (0.62-0.92)). Patients whose primary tumors expressed low levels of C14orf169 possessed median DMFS of 78 months, while patients whose tumors expressed high levels of C14orf169 possessed median DMFS of 118.82 months (Chart 6).

We also observed a statistically significant correlation between primary tumor expression of C14orf109 and recurrence-free survival (RFS) in patients with breast cancer in the upper survival quartile (Figure 8, Chart 7; logrank p -value: $3.3e-5$; hazard ratio: 1.29 (1.14-1.46)). Patients whose primary tumors expressed low levels of C14orf109 possessed median RFS of 54 months, while patients whose tumors expressed high levels of C14orf109 possessed median

RFS of 38 months (Chart 7).

We also observed a statistically significant correlation between primary tumor expression of C14orf130 and overall survival (OS) in patients with breast cancer (Figure 9, Chart 8; logrank p -value: 0.027; hazard ratio: 0.79 (0.63-0.97)). Patients whose primary tumors expressed low levels of C14orf130 possessed median OS of 81 months, while patients whose tumors expressed high levels of C14orf130 possessed median OS of 130 months (Chart 8).

We also observed a statistically significant correlation between primary tumor expression of C14orf132 and overall survival (OS) in patients with breast cancer (Figure 10, Chart 9; logrank p -value: 0.0028; hazard ratio: 0.72 (0.58-0.9)). Patients whose primary tumors expressed low levels of C14orf132 possessed median OS of 75.6 months, while patients whose tumors expressed high levels of C14orf132 possessed median OS of 136.8 months (Chart 9).

We also observed a statistically significant correlation between primary tumor expression of C14orf1 and distant metastasis-free survival in patients with breast cancer in the upper tertile (Figure 11, Chart 10; logrank p -value: 0.039; hazard ratio: 0.8 (0.65-0.99)). Patients whose primary tumors expressed low levels of C14orf1 possessed median DMFS of 87.6 months, while patients whose tumors expressed high levels of C14orf1 possessed median DMFS of 133.68 months (Chart 10).

We also observed a statistically significant correlation between primary tumor expression of C14orf2 and distant metastasis-free survival in patients with breast cancer (Figure 12, Chart 11; logrank p -value: 0.027; hazard ratio: 0.81 (0.67-0.98)). Patients whose primary tumors expressed low levels of C14orf2 possessed median DMFS of 85.81 months, while patients whose tumors expressed high levels of C14orf2 possessed median DMFS of 110 months (Chart 11).

Finally, we also observed a statistically significant correlation between primary tumor expression of C14orf93 and recurrence-free survival in patients with breast cancer (Figure 13, Chart 12; logrank p -value: $4e-5$; hazard ratio: 0.8 (0.71-0.89)). Patients whose primary tumors expressed low levels of C14orf93 possessed median RFS of 228.85 months, while patients whose tumors expressed high levels of C14orf93 possessed median RFS of 216.66 months (Chart 12).

Discussion

We provided evidence here, through analysis of published microarray datasets⁴⁻⁷, that a series of open reading frames on chromosome 14 is among the genes whose expression is most different in the brain metastases of patients with brain metastatic breast cancer, that C14orf mRNA is present at significantly increased quantities in brain metastatic tissues as compared to primary tumors of the breast, and that primary tumor C14orf expression is significantly correlated with patient survival outcomes in human breast cancer. Evaluation of the effects of genetic depletion of ORFs on chromosome 14 in mouse models of metastatic breast cancer on metastasis to the central nervous system is merited. It is possible that major structural aberrations on chromosome 14 in the brain metastases of women with breast cancer have thusfar escaped the notice of scientists and physicians, but it is more likely that epigenetic changes on chromosome 14 which are more difficult to detect and measure contribute to perturbations in C14orf expression we have demonstrated here; one group has described a translocation between chromosome 12 and 14 in the primary and brain metastatic tumor tissues of a woman with basal-like breast cancer¹⁰. The differentially expressed open reading frames on chromosome 14 we have identified here, including Cox16 and Ubr7, are united in their genomic localization but diverse in their functions. C14orf functions may be relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier and colonize the brain.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Sinn, B.V., Fu, C., Lau, R., Litton, J., Tsai, T.H., Murthy, R., Tam, A., Andreopoulou, E., Gong, Y., Murthy, R. and Gould, R., 2019. SET ER/PR: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ breast cancer*, 5(1), pp.1-8.
6. Iwamoto, T., Niikura, N., Ogiya, R., Yasojima, H., Watanabe, K.I., Kanbayashi, C., Tsuneizumi, M., Matsui, A., Fujisawa, T., Iwasa, T. and Shien, T., 2019. Distinct gene expression profiles between primary breast cancers and brain metastases from pair-matched samples. *Scientific reports*, 9(1), pp.1-8.
7. Salhia, B., Kiefer, J., Roas, J.T., Metapally, R., Martinez, R.A., Johnson, K.N., DiPerna, D.M., Paquette, K.M., Jung, S., Nasser, S. and Wallstrom, G., 2014. Integrated genomic and epigenomic analysis of breast cancer brain metastasis. *Plos one*, 9(1), p.e85448.
8. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
9. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.
10. Ding, L.I., Ellis, M.J., Li, S., Larson, D.E., Chen, K., Wallis, J.W., Harris, C.C., McLellan, M.D., Fulton, R.S., Fulton, L.L. and Abbott, R.M., 2010. Genome remodelling in a basal-like breast cancer metastasis and xenograft. *Nature*, 464(7291), pp.999-1005.

Chart 1: Differential expression of open reading frames on chromosome 14 in the brain metastatic tissues of patients with metastatic breast cancer.

Rank: 39
Probe ID: 22051
P-value: 7.43E-05
t: -5.43
B: 1.8129
Gene: C14orf112
Gene name: chromosome 14 open reading frame 112

Rank: 46
Probe ID: 19912
P-value: 1.05E-04
t: -5.24
B: 1.4986
Gene: C14orf169
Gene name: chromosome 14 open reading frame 169

Rank: 66
Probe ID: 20243
P-value: 1.63E-04
t: -5.01
B: 1.0995
Gene: C14orf109
Gene name: chromosome 14 open reading frame 109

Rank: 190
Probe ID: 2933
P-value: 8.23E-04
t: -4.18
B: -0.3863
Gene: C14orf179
Gene name: chromosome 14 open reading frame 179

Rank: 227
Probe ID: 3031
P-value: 6.71E-04
t: -4.22
B: -0.224575
Gene: C14orf130
Gene name: chromosome 14 open reading frame 130

The rank of differential expression, probe ID, p-value, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name is listed in this chart.

Chart 2: Differential expression of open reading frames on chromosome 14 in the lymph node metastatic tissues of patients with metastatic breast cancer.

Rank: 1408
Probe ID: 218820_at
P-value: 8.59E-06
t: 4.903434
B: 3.092892
Gene: C14orf132
Gene name: chromosome 14 open reading frame 132

Rank: 3775
Probe ID: 217188_s_at
P-value: 1.81E-03
t: 3.277519
B: -2.017784
Gene: C14orf1
Gene name: chromosome 14 open reading frame 1

Rank: 4944
Probe ID: 210532_s_at
P-value: 6.40E-03
t: -2.83382
B: -3.18654
Gene: C14orf2
Gene name: chromosome 14 open reading frame 2

Rank: 5625
Probe ID: 219526_at
P-value: 1.07E-02
t: -2.641223
B: -3.654708
Gene: C14orf169
Gene name: chromosome 14 open reading frame 169

Rank: 8673
Probe ID: 219009_at
P-value: 6.13E-02
t: -1.909882
B: -5.186593
Gene: C14orf93
Gene name: chromosome 14 open reading frame 93

The rank of differential expression, probe ID, p-value, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name is listed in this chart.

Chart 3: Differential expression of open reading frames on chromosome 14 in the lymph node metastatic tissues of patients with metastatic breast cancer.

Rank: 1964
Probe ID: 218820_at
P-value: 5.74E-02
t: -1.9598669
B: -4.2408
Gene: C14orf132
Gene name: chromosome 14 open reading frame 132

Rank: 2449
Probe ID: 217188_s_at
P-value: 7.54E-02
t: -1.8284797
B: -4.4587
Gene: C14orf1
Gene name: chromosome 14 open reading frame 1

Rank: 2467
Probe ID: 202279_at
P-value: 7.60E-02
t: -1.8244513
B: -4.4652
Gene: C14orf2
Gene name: chromosome 14 open reading frame 2

Rank: 5867
Probe ID: 218298_s_at
P-value: 2.19E-01
t: 1.2491672
B: -5.2632
Gene: C14orf159
Gene name: chromosome 14 open reading frame 159

The rank of differential expression, probe ID, p-value, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name is listed in this chart.

Chart 4: Differential expression of open reading frames on chromosome 14 in the brain metastatic tissues of patients with metastatic breast cancer.

Rank: 5936
Probe ID: A_23_P48771
P-value: 4.79E-04
t: 3.7537885
B: -0.534606
Gene: C14orf159
Gene name: chromosome 14 open reading frame 159

Rank: 7673
Probe ID: A_23_P25935
P-value: 1.50E-03
t: -3.3724595
B: -1.606865
Gene: C14orf1
Gene name: chromosome 14 open reading frame 1

Rank: 12263
Probe ID: A_23_P48581
P-value: 1.22E-02
t: 2.6065983
B: -3.534238
Gene: C14orf93
Gene name: chromosome 14 open reading frame 93

Rank: 14115
Probe ID: A_23_P163117
P-value: 2.20E-02
t: -2.3684783
B: -4.060089
Gene: C14orf169
Gene name: chromosome 14 open reading frame 169

Rank: 14198
Probe ID: A_23_P205281
P-value: 2.25E-02
t: -2.3599392
B: -4.078223
Gene: C14orf2
Gene name: chromosome 14 open reading frame 2

The rank of differential expression, probe ID, p-value, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name is listed in this chart.

C14orf112

Figure 1: C14orf112 is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of C14orf112 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in C14orf112 expression between primary tumors and brain metastasis, a p -value, is $p=0.0001$.

C14orf169

Figure 2: C14orf169 is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of C14orf169 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in C14orf169 expression between primary tumors and brain metastasis, a p -value, is $p=0.000142$.

C14orf109

Figure 3: C14orf109 is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of C14orf109 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in C14orf109 expression between primary tumors and brain metastasis, a p -value, is $p=0.000183$.

C14orf179

Figure 4: C14orf179 is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of C14orf179 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in C14orf179 expression between primary tumors and brain metastasis, a p -value, is $p=0.001325$.

C14orf130

Figure 5: C14orf130 is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of C14orf130 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in C14orf130 expression between primary tumors and brain metastasis, a p -value, is $p=0.001585$.

Figure 6: Significant correlation between C14orf169 primary tumor expression and distant metastasis-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of distant metastasis-free survival for $n=1803$ total patients stratified into two groups, based on low or high expression of C14orf169 in patient primary tumors. The log rank p -value denoting statistical significance of difference in distant metastasis-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf169 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf169 expression: 78.64 months

High C14orf169 expression: 125.65 months

Chart 5: Median distant metastasis-free survival is inferior in patients with low primary tumor expression of C14orf169.

The median DMFS of patients with low primary tumor expression of C14orf169 and high primary tumor expression of C14orf169 is listed in this chart, for $n=1803$ breast cancer patients.

Figure 7: Significant correlation between C14orf112 primary tumor expression and distant metastasis-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of distant metastasis-free survival for $n=1803$ total patients stratified into two groups, based on low or high expression of C14orf112 in patient primary tumors. The log rank p -value denoting statistical significance of difference in distant metastasis-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf112 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf112 expression: 78 months

High C14orf112 expression: 118.82 months

Chart 6: Median distant metastasis-free survival is inferior in patients with low primary tumor expression of C14orf112.

The median DMFS of patients with low primary tumor expression of C14orf112 and high primary tumor expression of C14orf112 is listed in this chart, for $n=1803$ breast cancer patients.

Figure 7: Significant correlation between C14orf109 primary tumor expression and recurrence-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=3951$ total patients stratified into two groups, based on low or high expression of C14orf109 in patient primary tumors. The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf109 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf109 expression: 54 months

High C14orf109 expression: 38 months

Chart 6: Median recurrence-free survival is inferior in patients with high primary tumor expression of C14orf109.

The median RFS of patients with low primary tumor expression of C14orf109 and high primary tumor expression of C14orf109 is listed in this chart, for $n=3951$ breast cancer patients.

Figure 8: Significant correlation between C14orf130 primary tumor expression and overall survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival for $n=1402$ total patients stratified into two groups, based on low or high expression of C14orf109 in patient primary tumors. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf109 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf130 expression: 81 months

High C14orf130 expression: 130 months

Chart 7: Median overall survival is inferior in patients with low primary tumor expression of C14orf130.

The median OS of patients with low primary tumor expression of C14orf130 and high primary tumor expression of C14orf130 is listed in this chart, for $n=1402$ breast cancer patients.

Figure 9: Significant correlation between C14orf132 primary tumor expression and overall survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival for $n=1402$ total patients stratified into two groups, based on low or high expression of C14orf132 in patient primary tumors. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf132 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf132 expression: 75.6 months

High C14orf132 expression: 136.8 months

Chart 8: Median overall survival is inferior in patients with low primary tumor expression of C14orf132.

The median OS of patients with low primary tumor expression of C14orf132 and high primary tumor expression of C14orf132 is listed in this chart, for $n=1402$ breast cancer patients.

Figure 10: Significant correlation between C14orf1 primary tumor expression and distant metastasis-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of distant metastasis-free survival for $n=1803$ total patients stratified into two groups, based on low or high expression of C14orf1 in patient primary tumors for patients in the upper survival tertile. The log rank p -value denoting statistical significance of difference in distant metastasis-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf1 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf1 expression: 87.6 months

High C14orf1 expression: 133.68 months

Chart 9: Median distant metastasis-free survival is inferior in patients with low primary tumor expression of C14orf1.

The median DMFS of patients with low primary tumor expression of C14orf1 and high primary tumor expression of C14orf1 is listed in this chart, for $n=1803$ breast cancer patients.

Figure 11: Significant correlation between C14orf2 primary tumor expression and distant metastasis-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of distant metastasis-free survival for $n=1803$ total patients stratified into two groups, based on low or high expression of C14orf2 in patient primary tumors. The log rank p -value denoting statistical significance of difference in distant metastasis-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf2 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf2 expression: 85.81 months

High C14orf2 expression: 110 months

Chart 10: Median distant metastasis-free survival is inferior in patients with low primary tumor expression of C14orf2.

The median DMFS of patients with low primary tumor expression of C14orf2 and high primary tumor expression of C14orf2 is listed in this chart, for $n=1803$ breast cancer patients.

Figure 12: Significant correlation between C14orf93 primary tumor expression and recurrence-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=3951$ total patients stratified into two groups, based on low or high expression of C14orf93 in patient primary tumors. The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on C14orf93 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low C14orf93 expression: 228.85 months

High C14orf93 expression: 216.66 months

Chart 11: Median recurrence-free survival is inferior in patients with high primary tumor expression of C14orf93.

The median RFS of patients with low primary tumor expression of C14orf93 and high primary tumor expression of C14orf93 is listed in this chart, for $n=3951$ breast cancer patients.