Impact Study of **Soil Health Card Scheme**

National Institute of Agricultural Extension Management (MANAGE)

(An organisation of Ministry of Agriculture and Farmers Welfare, Govt. of India)

Rajendranagar, Hyderabad – 500 030, Telangana State, India

www.manage.gov.in

Impact Study of Soil Health Card Scheme

NATIONAL INSTITUTE OF AGRICULTURAL EXTENSION MANAGEMENT (MANAGE)
(An organisation of the Ministry of Agriculture and Farmers Welfare, Govt. of India)
Rajendranagar, Hyderabad – 500 030, Telangana State, India
www.manage.gov.in

Citation: Reddy A Amarender (2018) Impact Study of Soil Health Card Scheme, National Institute of Agricultural Extension Management (MANAGE), Rajendranagar, Hyderabad – 500030, Telangana State, India. 106 pp.

वी. उषा रानी, मा.ए.स
सहानिदेशक

V. Usha Rani, IAS
Director General

Foreword

Government of India introduced Soil Health Card (SHC) Scheme to facilitate farmers for better understanding of soil and Integrated Nutrient Management (INM). It is a well known fact that soil is the basis for agriculture and protecting the soil is the basic responsibility of every farmer to sustain agriculture. Unfortunately, quality and yields of agriculture produce is not at expected level due to deficiency of various nutrients. At the same time, there is excessive usage of certain nutrients. All this is happening as farmer is not fully aware of various physical, chemical and microbial activities in soil and due to this, farmers are unable to apply fertilizers in balanced and required quantities.

The Department of Agriculture, Cooperation and Farmers Welfare assigned MANAGE to undertake study of implementation of Soil Health Card Scheme. The study brought out important aspects of whole gamut of activities starting from collecting soil sample, analysis and interpretation of data.

I congratulate Dr. Amarendra Reddy, Director (Monitoring & Evaluation) and Principal Investigator and his team for the hard work they have rendered by taking up this study. I hope the study will further strengthen the scheme and will have impact on the farmers knowledge on soil and application of fertilizer in a balanced manner.

(V. Usha Rani)

राष्ट्रीय कृषि विस्तार प्रबंध (मैनेज)

(कृषि एवं. किसान कल्याण मंत्रालय, भारत सरकार का संगठन, राजेन्द्रनगर, हैदराबाद-500 030, टी.एस. भारत)

NATIONAL INSTITUTE OF AGRICULTURAL EXTENSION MANAGEMENT (MANAGE)

(An organization of Ministry of Agriculture and Farmers Welfare, Government of India)

Rajendranagar, Hyderabad-500 030, T.S. India

Ph: +91(40) 24015253 (O), Fax: 040-24015388

E-mail: dgmanage@manage.gov.in, Web: manage.gov.in

Acknowledgement

The study on “Impact of Soil Health Card Scheme” has been carried out at the National Institute of Agricultural Extension Management (MANAGE), Rajendranagar, Hyderabad, as suggested and sponsored by the Ministry of Agriculture and Farmers Welfare, Government of India.

We have benefited immensely from various scholars and officials from different government departments while carrying out this study. At the outset, we would like to thank Smt. V Usha Rani, IAS, Director General of our institute as well as Smt. Rani Kumudhini, IAS then Joint Secretary, INM, Ministry of Agriculture and Farmers Welfare, Government of India for their constant encouragement and support for undertaking this impact study. We are grateful to Smt. Neerja, IAS, Joint Secretary, INM, Ministry of Agriculture and Farmers Welfare, Government of India and Dr. Chaudhary, Additional Commissioner, INM, Department of Agriculture and Cooperation and Farmers Welfare for continuous support and guidance.

We are grateful to directors and joint directors and other officials from different state department of agriculture for their cooperation during the field survey and later interactions in focus group interactions for sharing their valuable suggestions. We thank Dr. Ratna Reddy, Dr. Padma Raju, ex Vice Chancellor, PJTSAU, Rajendranagar, Dr. CP Chandrashekar, former dean, PJTSAU for their guidance and active involvement.

We thank our colleagues in MANAGE for their support and encouragement while carrying out the study.

We are also thankful to the officials and sample farmers in the study area. The study would not have reached this stage without the active co-operation of the sample households, who provided all the required data for the study without any hesitation and expectation. We thank each one of them for their invaluable support.

A Amarender Reddy
Principal Investigator

Abbreviations

ASCI	Agriculture Skill Council of India
CACP	Commission for Agricultural Costs and Prices
DSS	Decision Support System
EC	Electrical Conductivity
FAO	Food and Agriculture Organization
FC	Forward Class
FPCs	Farmer Producer Companies
GIS	Geographic Information System
GPS	Global Positioning System
ICAR	Indian Council for Agricultural Research
ICRISAT	International Crops Research Institute for the Semi-Arid Tropics
IFPRI	International Food Policy Research Institute
IISS	Indian Institute of Soil Science
IPCC	Intergovernmental Panel on Climate Change
ISRO	Indian Space Research Organization
ITPS	Intergovernmental Technical Panel on Soils
KVK	Krishi Vigyan Kendra
NABARD	National Bank for Agriculture and Rural Development
NAPCC	National Action Plan on Climate Change
NGO	Non-Governmental Organization
NMSA	National Mission for Sustainable Agriculture
OBC	Other Backward Caste
OC	Organic Carbon
PPP	Public Private Partnership
RAU	Rajendra Agricultural University
S&MF	Small and Marginal Farmers
SAU	State Agricultural University
SC	Scheduled Cast
SHC	Soil Health Card
ST	Scheduled Tribes
STCR	Soil Test Based Crop Response
STL	Soil Testing Labs
UNCCD	United Nations Convention to Combat Desertification
UNFCCC	United Nations Framework Convention on Climate Change
WS	World Soil Charter

Contents

Page no:

Executive Summary	i
Chapter 1: Introduction	1
Chapter II: Soil Health in India - Review of Experience	4
2.1: Soil Health in India	5
2.2: Soil Nutritional Status	5
2.3: Soil Organic Carbon	6
2.4: Soil Health Card to improve Soil Health	7
Chapter III: Policy interventions over the years: International and National	10
3.1: Global Policies	10
3.2: National Policies	11
Chapter IV: Objectives and Methodology	13
4.1: Objectives	13
4.2: Methodology	13
4.2.1: Selection of States, Districts & Blocks	13
4.2.2: Structure of Data from Farmers	15
4.2.3: Indicators of inputs of Soil Health Card (SHC) scheme	15
4.2.4 : Indicators of activities under SHM	15
4.2.5: Indicators of outputs of the Scheme	16
4.2.6: Indicators of outcomes	16
4.2.7: Indicators of Socio-economic impact	16
4.2.8: Information from key stakeholders & secondary data	16
Chapter V: Soil Health card Scheme: Design , Coverage & Impact	17
5.1: SHC Design and Status (inputs/activities/and outputs)	17
5.2: Status of SHC (Financial and human resource allocation)	27
5.3: Farmers feedback on the Design	33
5.4: Design in different States	37
5.5: Improvements required in the SHC Scheme design	38
5.6: Socio-economic characteristics of the sample farmers	41
5.7: Awareness, utilization and outputs under the SHC scheme	42
5.8: Farmers perception and source of information about SHC	48
5.9: Outcomes (Input use and Cost per hectare) of the SHC scheme	57
5.10: Success stories and potential benefits of the SHC scheme	62
5.11: State wise crop wise fertilizer consumption	63

Chapter VI: Policy Recommendations	68
6.1 Soil Sampling related (SHC design)	72
6.2 Soil Health Indicators (SHC design)	74
6.3 Soil Testing Infrastructure	75
6.4 Soil Extension	76
6.5 Policy related	77
References	79
Annexure-1: Feedback from agricultural officers on status of SHC and improvement needed	80
Annexure-2: Soil Health & Soil Testing: Some Practical Issues	82
Annexure-3: Field Observations compiled across states	86
Appendix-1: Review of Recent Studies	88
Appendix -2: Regression Analysis: Determinants of Performance indicators of SHC Scheme (District Level)	98
Appendix-3: Farmers Questionnaire	101

List of Tables

S.No.	Particulars	Pg.Nos.
Table 1:	Consumption of Fertilisers in million tonnes in India	2
Table 2	Soil Quality and Quantity Indicators	12
Table 3	Details of Sampling Framework	14
Table 4A	Financial Allocations and Expenditure	27
Table 4B	Zone wise analysis of funds allocation, release and utilisation	28
Table 5:	Infrastructure (number of Labs) in 2016-17	29
Table 6A:	Samples entered, farmers covered, samples tested and SHCs printed (Cycle-1)	31
Table 6B:	Samples entered, farmers covered, samples tested and SHCs printed (Cycle-2)	32
Table 7A:	Indicators mentioned in SHCs in different states of USA	35
Table 7B:	Indicators mentioned in SHCs in other countries	36
Table 8:	Basic Features of the Sample Households (HH)	41
Table 9:	Basic Features of the Sample Households (HH) by irrigation status	41
Table 10:	Zone wise awareness about SHC	42
Table 11:	Farm Size Class wise awareness about SHC	43
Table 12:	Social group wise awareness about SHC	43
Table 13:	Awareness about SHC in developed and less developed states	44
Table 14:	Awareness about SHC by Irrigation Status of the Farmers	45
Table 15:	Participation of Farmers in SHC Activities across Regions	46
Table 16:	Participation of Farmers in SHC Activities across States	46
Table 17:	Participation of Farmers in SHC Activities across Size Classes	47
Table 18:	Participation of Farmers in SHC Activities across Social Groups	47
Table 19:	Participation of Farmers in SHC Activities across irrigated area groups	47
Table 20:	Farmers perception about SHC across Regions	48
Table 21:	Farmers opinion about SHC across Size Classes	49
Table 22:	Farmers opinion about SHC in developed and less developed states	50
Table 23:	Farmers opinion about SHC across Irrigation Categories	50
Table 24:	Number of plots covered in soil testing across Zones	51
Table 25:	Number of plots covered in soil testing across Size Classes.	51
Table 26:	Number of plots covered in soil testing	52
Table 27:	Number of plots covered in Irrigation Groups	52
Table 28:	Zone wise source of information about SHC content	53
Table 29:	Source of information about SHC content across size classes	54
Table 30:	Source of information about SHC content by state groups	54
Table 31:	Source of information about SHC content by level of irrigation	54
Table 32:	Farmers Perceptions about SHC across Zones	55
Table 33:	Farmers Perceptions about SHC across Size Classes	55
Table 34:	Farmers Perceptions about SHC across Social Categories	56
Table 35:	Farmers Perceptions about SHC across States	56

Table 36:	Farmers Perceptions about SHC Irrigated Groups	57
Table 37:	Impact of SHC on fertilizer use, costs and crop yields	58
Table 38:	Impact of SHC on fertilizer use, costs and crop yields by farm size category	59
Table 39:	Status according to longevity (period of holding SHC) (cotton)	60
Table 40:	Status according to longevity of card (Paddy)	60
Table 41:	Status according to longevity of card (soybean)	60
Table 42:	Status according to longevity of card (pigeon pea)	61
Table 43:	Status according to longevity of card (Wheat)	61
Table 44:	Success stories of reduction in fertilizer use	62
Table 45:	Analysis of Fertilizer use by crop across states (CACP data) 2013-14	64
Table 46:	Fertilizer use (kg/ha) in 2014-15	66

List of Maps

S.No.	Particulars	Pg.Nos.
Map 1:	Status of Nitrogen and Zinc availability in Indian soil(Source: Economic Times)	7
Map 2:	Status of Phosphorus and Iron availability in Indian soil	8
Map 3:	Status of Potassium, Copper and Manganese availability in Indian soil	8
Map 4:	State Wise Consumption of Fertilizers	9
Map 5:	Changing chemical composition of soils	9
Map 6:	State wise samples collected per 1000 ha.	18
Map 7:	State wise samples collected per 1000 cultivators	18
Map 8:	State wise samples tested per 1000 ha	18
Map 9:	State wise samples tested per 1000 cultivators	18
Map 10:	State wise SHCs printed per 1000 ha.	20
Map 11:	State wise SHCs printed per 1000 cultivators	20
Map 12:	State wise SHCs distributed per 1000 ha	20
Map 13:	State wise SHCs distributed per 1000 cultivators	20
Map 14:	District wise samples collected per 1000 ha	21
Map 15:	District wise samples tested per 1000 ha	22
Map 16:	District wise SHCs printed per 1000 ha	23
Map 17:	State wise number of soil testing laboratories	24
Map 18:	State wise Soil Testing Labs per Million Hectares	25
Map 19:	District wise STLs per million cultivators	26

List of Figures

S.No.	Particulars	Pg.Nos.
Fig. 1:	Measure of Soil Health	1
Fig. 2:	State wise Fertilizer use (kg/ha) and N:K Ratio	2
Fig. 3:	Progressive expansion in the occurrence of nutrient deficiency	3
Fig. 4:	Fields effected by alkalinity	6
Fig.5:	Importance of including water test in the SHC	34
Fig.6	Landholding wise awareness about SHC	43
Fig.7	Method of Soil Sample Collection	44
Fig.8	Awareness about SHC Scheme	44
Fig.9	Soil sampling equipment	45
Fig. 10	Percentage of farmers in SHC activities and perception	46
Fig. 11	Methods for evaluating Soil Nutrient Status	49
Fig. 12	Number of plots covered for soil testing	51
Fig. 13	Number of plots covered in developed and less developed states	52
Fig. 14	Main Source of Information (%)	53
Fig. 15	Farmers perceptions about SHC across social categories	56
Fig. 16	Farmers perceptions about SHC across States	57
Fig. 17	Soil Test Interpretation Categories	58
Fig. 18	Azolla- a potential source of bio fertilizer	59
Fig. 19	Soil Test Crop Response Based Fertilizer Recommendation System	63
Fig. 20	Soil Test Crop Response Based Fertilizer Recommendation System	63

Executive Summary

Soil health and fertility are the basis for gaining sustainable profit through higher productivity by the farmers. Using optimal doses of fertilisers and cropping pattern as per the scientific recommendations is the first step towards sustainable farming. Soil testing is a science-based and time-tested tool for assessment of the soil fertility status and soil ailments and nutrient amendment recommendations. Soil testing, as a tool for judicious fertiliser use, works on the principle of profitability, which means if all other factors of production are optimum and none of them are limiting, there is a high probability of obtaining a more profitable response to applied nutrients based on soil testing than those applied on an ad hoc basis.

In India, the current consumption of NPK ratio is 6.7:2.4:1, which is highly skewed towards nitrogen as against the ideal ratio of 4:2:1. India is spending nearly Rupees Seventy thousand crore on fertiliser subsidy every year. According to the estimates, subsidy amount is about Rs. 5,000/ha of net cropped area and about Rs. 5,100/farmer, resulting in excessive use of fertilisers, especially NPK, at the cost of micro nutrients and manure. Hence, there is a need for balanced use of fertilisers. Keeping this in mind, the Government of India introduced Soil Health Card Scheme across India (GoI, 2017).

On December 5, 2015, the Ministry of Agriculture introduced the Soil Health Card scheme. It was approved for implementation in the remaining period of 12th plan. SHC will be provided to all farmers in the country at an interval of two years to enable the farmers to apply recommended doses of nutrients based on soil test values for improved and sustainable soil health and fertility, low costs, and higher profits.

Under the SHC scheme, cropped area was divided into grids of 10 ha for rainfed and 2.5 ha for irrigated. One soil sample from each grid was to be taken, and the test results were to be distributed to all the farmers whose lands fell under the grid. Based on the grid system, of the total 14.1 crore hectare of net cropped area, 73 lakh grid samples were to be collected to cover 7.3 crore ha in rainfed areas and 2.7 crore grid samples were to be collected to cover 6.8 crore ha irrigated land. That is, a total of 3.46 crore grid samples in two years (1.73 crore grid samples per year) and an average of 25,000 grid samples per district/year or 29 grid samples per village/year. With this, all 11 crore farmers will be covered in two years. Every year, 5.2 crore farmers need to be covered.

Under cycle 1, 2.54 crore samples were collected, 2.36 crore samples tested, 9.62 crore soil health cards printed, but only 9.33 crore SHCs distributed. This indicates that 100% target was achieved in sample collection, 93% of the target was achieved in soil testing, but only 80% of the target was achieved in SHC printing. 97% of the SHCs printed were distributed among the farmers as on September 24, 2017. Cycle 2 has already started in many states. Overall, the progress of the SHC scheme in terms of coverage is satisfactory, and now we have to focus more on the quality of soil sample collection and testing and timely distribution of SHCs to farmers. However, the progress is highly skewed. Some states, such as Karnataka, TN, Chhattisgarh, UP, Maharashtra, Telangana, and AP, were better performers compared to other states.

Objective of the impact study

As the SHC scheme has completed more than 2 years of implementation, the ministry has initiated a nationwide impact assessment with the following objectives:

- o To examine the design of the SHC scheme in terms of planning, implementation, inputs (staff, financial, and other resources), activities (trainings, lab established, and strengthened), outputs (SHCs printed and distributed to farmers).
- o To assess the modalities of delivery of the SHC scheme regarding procurement, sample collection, testing, SHC printing, and disbursal.
- o To assess the level of utilisation of SHCs by the farmers across farm size classes in irrigated and rainfed situations.
- o To assess the impacts of the SHC scheme on judicious use of fertilisers, bio-fertilisers, organic fertilisers, soil health, cropping choice, cost reduction, farm profitability, and sustainability.
- o To provide recommendations for improvement of the overall design of the programme.

Methodology

Both quantitative and qualitative approaches were adopted to achieve the objectives of the study. Qualitative information was in the form of stakeholder interviews across the states under the study, expert opinion gathering at the national and state level workshops, and interactions with the progressive farmers and agricultural officers. At the quantitative level, both secondary and primary data was collected at the national, state, and farmer levels. Secondary data mostly pertains to collection and analysis of financial and physical achievements of the SHC scheme over the years, infrastructure availability, coverage of SHCs across the states, etc. Besides, information at the international level was collected for some selected countries to see the best practices in the design of soil health cards.

The secondary data was analysed for all the states, while primary data was analysed for 16 states of India representing all agro-climatic zones. A systematic sample was drawn for the impact assessment at the farmer level. Care was taken to represent the whole country and its agro-climatic conditions. A structured questionnaire was canvassed among 3,184 sample farmers across 199 villages in 16 states. In addition, focus group discussions were conducted in each village in order to get feedback from the key informants, farmers not covered in individual surveys, and farmers who had not received their soil health card. All indicators collected from the field survey were classified as inputs (financial and physical inputs under the project), activities (different activities organised under the scheme), outputs (actual outputs of the project), outcomes (whether generated outputs were utilised by the farmers), and impacts (what are the ultimate benefits to the farmers); they're listed here. The analysis was carried out across zones/states and completed by the date of receiving SHC by the farmers, i.e., those who received it more than a year back and those who received it recently. This would help understand the long-term impacts and provide insights into whether agriculture development helps better awareness and demand for SHC.

Main Findings

It may be noted that the analysis is based on the representative, though limited, sample size across regions. This is too short a time for the scheme (only two-years old) to carry out a full-fledged impact assessment. The present analysis provides insights into the direction and cautions about any shortcomings. While the following conclusions and recommendations are based on the analysis, the weakness of the assessment needs to be kept in mind.

- ✓ Given the short duration of the scheme, awareness levels are good. . At the same time, participation of farmers in meetings and exposure visits is not high. Awareness campaigns need to be organised on the content of SHCs, use of recommended practices, reduction in fertiliser use, and costs and increase in profitability.
- ✓ There is no apparent or significant bias against socio-economically vulnerable sections. In contrast, small and marginal farmers benefit more in some cases.
- ✓ There is some reduction in fertiliser use, especially nitrogen and increase in bio-fertilisers and other micro-nutrients' use. This is a good sign as the N:P:K ratio was highly skewed towards nitrogen. Costs were reduced due to low fertiliser use. Crop yields also increased for a majority of the crops, although only moderately.
- ✓ A significant impact is the increase in the use of gypsum and other micro nutrients to some extent.
- ✓ There is a need to strengthen the SHC-related extension services to provide better advisories.
- ✓ Two-thirds of the sample farmers indicated that SHC is beneficial, which is encouraging given the short span of the programme.
- ✓ The main complaint from the farmers is the timeliness of providing the results. This, however, is linked to the infrastructure (soil testing labs) and human resources. However, after the introduction of the SHC scheme, the time lag has significantly reduced. Results need to be disseminated before the sowing season so that farmers will practice recommended crop choice and fertilisers.
- ✓ It is important to address these issues to gain confidence of the farmers in adoption of the fertilisers as per the recommendations included in the SHC.
- ✓ The scheme has poor backing from infrastructure and human resources, with significant gaps. Although some southern and western states performed better, in some states, even the allocated resources are not being spent or utilised due to lack of capacities. This should be of high priority in the immediate future.
- ✓ Proactive regions seem to be better in this regard.
- ✓ Lack of capacities with regard to skilled personnel and STLs affects the quality of services, which in turn affects the credibility of the scheme and needs immediate attention.
- ✓ Results need to be provided in time so that farmers can benefit in a better way.

Level of Utilisation and Impact of the SHC Scheme

1. About 66% of the farmers are able to understand the content of the SHC, about 57% mentioned that the recommendations are suitable for their farms, and about 53% are able to follow recommendations.
2. The SHC scheme is inclusive in nature; small and marginal farmers are proactive in the adoption of recommendations based on SHC.
3. There was a reduction in the use of urea and DAP by 20 to 30% in paddy and cotton in some states, resulting in decreased cost of cultivation. The reduction in the cost of cultivation ranged between Rs. 1,000 and Rs. 4,000 per acre.
4. The use of micro-nutrients (especially gypsum) was slightly increased after SHC distribution.
5. There was a significant increase in yield for farmers who practiced recommended practices as per the SHC.
6. With decrease in the cost of cultivation and increase in yields, net incomes of the farmers increased between 30 and 40% after the SHC scheme.

Constraints of the SHC Scheme

1. Some farmers complained that the soil test values are not representative of their fields and that the field staff did not collect the soil samples in their presence. To build trust, samples need to be collected in the presence of GRID farmers.
2. Uniform soil GRIDS of 10 ha for rainfed and 2.5 ha for irrigated should be re-examined. The grid size should be determined based on the soil variability index (soil variogram). If variability is high, GRID size should be less and vice versa.
3. Soil variogram needs to be developed at each block level to determine the GRID size at the block level.
4. Many farmers are unable to understand the content and hence are unable to follow the recommended practices.
5. Only 44% of the farmers mentioned that the extension officers explained the content.
6. SHC distribution and awareness campaigns needs to be arranged before the sowing season so that farmers can practice recommended crop choice and fertilisers.
7. Awareness campaigns need to be organised for the use of SHC in reduction of fertiliser use and costs and increase in yields.
8. Many farmers feel that SHC should also mention one or two physical and micro-biological indicators (such as soil texture, water holding capacity, and water quality and bacterial content).
9. There is a need to identify best practices in soil sample collection (predetermined GRID apps practised in Punjab and TN) and testing for scale-up.
10. Development of GIS-based soil fertility maps at the village/block level and wider publicity through wall posters and display boards in village panchayats should be promoted.
11. The Government should ensure availability of recommended fertilisers and bio-fertilisers at the village level at reasonable prices.

Soil - Sampling - related (SHC Design) Improvement

- There is a need to identify best practices in soil sample collection and testing by examining across countries and different state governments practices. There is also a need for coordination and cooperation.
- The existing uniform grid of 10 ha for drylands and 2.5 ha for irrigated lands does not take local soil variability into consideration. Grid size should be variable based on the soil variability index and should be decided at least at the block level based on soil heterogeneity, fertility maps, cropping pattern, irrigation facilities, and remote sensing maps. If the soil is more variable, grid size should be reduced and vice versa. Sampling errors need to be reduced using variogram. There should be a separate cell to monitor and recommend grid size across the country. This will also reduce cost, money, and manpower and increase the relevance of recommendations to farmers. In order to gain the credibility of the farmers, at least one sample from each farmer should be included where soil variability is high.
- Soil variograms needs to be developed at each block level. Based on the soil variability, the grid size can be determined. Variogram gives information about spatial pattern of continuous soil attributes. The variogram may be used as a critical input to decide required soil samples to be collected based on the soil variability index. More soil samples should be collected if the block-level soil variability index was high and vice versa. Variogram is a tool to investigate and quantify the spatial variability of soil properties. The geostatistical literature shows that the following soil quality indicators were found to be the most important for variogram analysis, and they can also be considered by the ministry. The available data with Indian Institute of Soil Science, Satellite maps of remote sensing agency of ISRO, and Land use planning data can be used to estimate block-level soil variograms.
- The following soil indicators may be considered to construct block-level soil variograms: (1) Soil Colour, (2) Slope, (3) Sand (%), (4) Silt (%), (5) Clay (%), (6) Nitrogen (N), (7) Phosphorus (P), (8) Potassium (K), (9) Organic Matter (OM), (10) Organic Carbon (OC), (11) pH, (12) Cation Exchange Capacity (CEC), (13) Electrical conductivity (EC), (14) C:N Ratio, (15) percentage of irrigated area, and (16) cropping pattern. The ICAR-Indian Institute of Soil Science (IISC), Bhopal; ICAR-IISWC- Indian Institute of Soil and Water Conservation, National Remote Sensing Agency (NRSA), Hyderabad; and land classification/atlas can be consulted for developing block-level soil variograms.
- Evidence shows that samples collected from 20% to 30% of the farmers in a village are enough to get reasonable soil quality for advising farmers, and hence there was no additional benefit in covering each grid of 10 ha in case of dry lands and 2.5 ha in case of irrigated land if the soil was fairly uniform. In some cases, only 20-30 samples/500 ha are sufficient, as is evident from ICRISAT experiments. This, however, needs to be explained to the farmers so that they would take the SHC recommendations seriously. One needs to be mindful of the efforts and resources gone into ICRISAT experiments.

- Although grids in some states were predetermined in the mobile app (such as in Punjab), in other cases, the procedure that is followed in dividing the cultivated village land into grids is not known to many agricultural officers and needs to be widely disseminated for accurate sampling, and this should be mentioned in the guidelines. Interestingly, Punjab is adopting GIS-tablet grid identification and GPS-based soil sample collection application, which seems to be working well and is likely to solve many field-level sample collection problems. This model should be adopted across all the states after an in-depth understanding (study) of the model.
- High density soil maps need to be developed for increasing precision at the village level.
- There is a need to give appropriate training, easy-to-use sampling tools, and transport to agricultural officers and agricultural extension officers. Incentives need to be provided for scientific sample collection.
- Coordination of agricultural extension officers and farmers needs to be enhanced and extension officers should make sure that most of the grid farmers, if not all, should be present at the time of soil sample collection. This will build farmers' confidence in SHCs.
- It was observed by the study team that in some of the block agricultural offices, soil samples were kept aside for many days and soils were exposed to moisture and weather. After soil sampling, drying should be done within 15-20 days, and grinding, machine sieving, and bottling should be done in time for proper test results. Sample test results should reach farmers before the sowing season. It might be a good idea to limit samples to the capacities. There should be more focus on the quality of soil sample collection and testing rather than coverage, as we have already covered larger areas and farmers with SHCs. This would create demand for soil testing once the credibility of the testing is established.

Improvement in Soil Health Indicators (SHC Design)

- The whole chain of soil health-plant-health-human health should be taken into account and there is a need for promotion of application of balanced application of soil (macro and micro) nutrients.
- Excess application of urea results in accumulation of nitrate in the soil and water is becoming a huge environmental problem in India. Hence, water quality information needs to be included in the SHC.
- The soil health card is more focused on chemical nutrient indicators; among physical and biological properties, only soil colour is included. Some more physical properties, such as slope of the land etc. needs to be incorporated as well.
- Microbial activity and moisture retention activity are essential but missing in SHC. Although soil organic matter is indicated, many soil testing labs are not equipped with the latest tools to measure it.
- At least one or two physical and micro-biological indicators (such as soil texture, water holding capacity, water quality, and bacterial content) need to be incorporated. Soil health index needs to be developed and incorporated into SHC, which indicates the overall health of the soil.

- Although the basic structure of SHC should be uniform, states should adopt/change as per their agro-climatic zones and needs. Some of the indicators that needs to be included in SHC are (i) cropping history, (ii) water resources (soil moisture), (iii) slope of soil, (iv) depth of soil, (v) colour of soil, (vi) soil texture (bulk density), and (vii) micro-biological activity.

Soil Testing Infrastructure (Design and Delivery)

- About 1,454 labs exist in India, of which only 700 are equipped with micro-nutrient testing facilities. Although agricultural departments procured about 7,000 mobile kits recently, they are not as good as those in full-fledged labs. Very few labs could take up micro nutrient analysis. They are neither equipped with skilled personnel nor chemicals or functional equipment. The infrastructure is grossly inadequate by any standard, given that 11 crore farmers need to be covered.
- Under the current PPP model, investments in labs are to be done by private companies with an element of subsidy. A competitive bidding process based on technical and financial bids needs to be announced, and companies that quote reasonable cost (per sample) should be selected. The Government will pay on a per-sample basis with the condition that they employ qualified and trained chemists. This model will be successful when there is no strong government presence in soil testing. However, the quality of such reports should be checked at random by authentic agencies.
- About 45% of the sample farmers are inclined to go to private STLs. At the same time, only 20 % of the farmers are willing to pay for the services. Hence, one must find ways to support farmers in this regard i.e., direct subsidy to the farmers or private STLs, etc. A competitive PPP model could be explored in this regard, while the government should take up the monitoring of the functioning of these labs more seriously. At the same time, there should be special incentive for Farmer Produce Companies (FPCs) to establish soil testing labs. There is a need for encouraging competition among private companies in setting up and running the soil testing labs so that they maintain quality at a reasonable cost. Institutional modalities could be worked out on how to equip and manage STL within the FPCs frame work.
- Some of the private soil testing labs indicated that the cost of sample collection and testing was up to Rs. 1,000 / sample. Some private companies are charging Rs. 75/element; accordingly, for 12 elements, the total cost will be Rs. 900/sample. The government should be more realistic in fixing the prices for private parties. It should focus more on the quality of the services at an acceptable (market) cost.
- There should be strengthening and upgrading of at least one soil testing lab per district as a state-of-the-art lab, equipped with world class infrastructure and accredited by internationally recognised agencies either in the public or the private sector. So, nearly 700 state-of-the-art labs are needed to act as referral labs and to give broad advice to farmers. The cost per unit will be about Rs. 4-5 crore/unit, with a total of Rs. 2,800 crore. However, if this resulted in Rs. 1,000/ha savings in fertiliser use even if we don't consider the yield increase resulting in a saving of Rs. 14,500 crore in a year for the economy. This could be a more worthwhile investment rather than spending money and providing services that have little value to farmers year after year, i.e., ending up spending more in the long run.
- State-of-the-art district level soil testing labs at a direct level should be equipped with Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES), which costs about Rs. 40-50 lakh. These labs should have a 24-hour generator for uninterrupted power supply,

computer labs with colour printing facility, and air conditioned laboratories. In addition, the lab should have the following equipment for conducting soil testing on a large scale:

KEL PLUS automatic nitrogen determination distiller
Automated Flame Photometer (for Potassium)
Automated Spectrophotometer (for Phosphorus)
Atomic Absorption Spectrophotometer (for Zn, Fe, Mn, Cu)
Water distillation still (20 lit/hour)
All glass distillation unit (5 litre/hour)
Auto analyser (N&P)
Automated pH meter
Automated EC meter
Centrifuge
Auto Burette

- Some soil scientists and agricultural officers are of the opinion that test results of mini-kits (mini-labs) are not accurate enough as that of full-fledged labs. Mini kits need to be standardised and tested for errors in calibrations. A mini kit costs about Rs. 94,800, with which the per-sample cost comes about Rs. 170-200. Some block agricultural offices received five-six mini kits, but they were not able to utilise them, as they were involved in multiple activities. A mini kit is useful for remote villages and tribal communities and should also be used to measure highly volatile elements such as Nitrogen, which needs more frequent measurement. Mini kits can be used exclusively for Nitrogen estimation in all block/mandal-level offices.
- Soil testing is a specialised and highly skill-oriented job. Frequent transfers of soil testing staff adversely affect the skill development within labs as well as the test results. There is a need to build some permanent staff in the labs who are interested and specialised in soil testing. Field observations indicate that only female officers are interested in working with soil testing labs.

Management of state-of-the-art soil testing labs could be established under the purview of FPC's federations or nodal FPCs at the district level. The governance responsibilities should be handed over to them to be run them as business models. The back of the envelop indicates that establishing a state-of-the-art lab with a Rs. 6 crore loan from NABARD in each district looks viable given that an average of 25,000 samples need to be collected and processed per year. At Rs. 1,500-2,000 per sample (at half the market price for a detailed soil analysis), the investments will be paid back in less than 2 years. The state department can have a monitoring cell created especially for this purpose, and extension services needs to be geared up to deal with soil health advisories. FPCs should be encouraged to set up demonstration plots to increase demand for soil testing by the farmers.

Soil Health Card Delivery issues

- SHC recommendations should be accompanied by block-level recommendations. There should be an intermediate solution (based on both village-level soil maps and SHC recommendations) to be able to reach the farmer's level.
- There is a need for demonstration of SHC benefits on an experimental basis in each block by adopting a comprehensive approach (systematic and scientific analysis of soil and water) and adoption of recommended doses. This would have much greater impact than the subsidised and less authentic information. General SHC schemes and model farm initiatives should go together.

- In many villages, agricultural officers are distributing SHCs in awareness campaigns through village presidents and mandals/blocks' democratically elected representatives. However, in some villages, the village revenue assistant distributes SHC and takes signatures without explaining the content. Whenever SHC is distributed in awareness campaigns and meetings directly, a greater number of farmers feel that they are convinced to use recommended practices. There is a need to follow standard protocol to inform farmers about the recommendations of the SHC when it is handed over to farmers.
- A specialised body is needed both at the central as well as state level for the management of soils. They should be given the responsibility of monitoring the quality of service by various agencies. This also provides continuity of work by the department.
- Development of GIS-based soil fertility maps at the village/block level and wider publicity through wall posters and display boards in village panchayats should be promoted. Advertisements, slogans, etc. should be developed in local languages to increase awareness. This should be taken up in a campaign mode, i.e., on the lines of anti-smoking/tobacco campaign.
- Many farmers are not aware of the SHC portal. The SHC portal should be more farmer friendly and simplified. A professional body may be employed to design the portal in more farmer friendly and effective manner.
- A simple tool to assess the quantity of urea, DAP, and MoP based on SHC needs to be displayed as wall posters in the villages.
- It should be mandatory to enter fertiliser purchases by the farmers on the soil health card by each fertiliser dealer along with the signature. This will not only increase awareness, but it will also help in adoption as per the recommendation.

Policy Recommendations for Improvement

- Index of soil health needs to be developed and incorporated into SHC, which indicates the overall health of the soil. Based on the index, soils should be classified as grade A, B, or C. The grades can be updated every SHC cycle based on soil tests. This updating of soil health index may act as an incentive to farmers to put in special efforts to upgrade soil health index, say, from Grade C to Grade A, as grade A soils may fetch higher land rental value for agricultural purposes and get higher land sale price if the land is put for sale compared to soils with grade C.
- All agricultural and rural development schemes should make it mandatory to mention soil health index based on SHC along with Aadhar in their application forms in order to get benefits. The SHC information should be linked to Aadhar and be available for download at e-seva in the public domain.
- If the SHC programme is to be successful, the high fertiliser subsidy for NPK needs to be reduced. Prices should reflect true cost to the economy; only then will farmers have an incentive to use fertilisers judiciously according to the recommendations of the SHC. At the same time, subsidy on use of recommended dose of micro-nutrients, bio-fertilisers, and organic inputs should be encouraged. However, the quality of these inputs supplied on subsidy needs to be

monitored and stringent punishment needs to be enforced if quality standards are not maintained.

- The government should set up state-of-the-art labs to test the quality of micro nutrients supplied. Accreditation of such labs to national/international standard institutes should be initiated. Supply of phosphorous soluble bacteria should be mandatory along with phosphate fertilisers and rock phosphate such as neem-coated urea. FPCs need to be encouraged to take up the SHC scheme as a business model by setting up state-of-the-art labs at the district level. Since FPCs are already involved in selling fertilisers, they are in a better position to stock all the required (according to SHC recommendations) fertiliser and micro-nutrient compositions and supply them to individual FPCs across the district.
- There should be some incentives/awards for the farmers who grow green manure and vermi-compost and whose soil fertility increased over the years based on Soil Health Card.
- Some incentives should be provided to local bodies that encourage good practices such as recycling crop residues, encouraging common lands for corporates, etc.
- Similarly, incentives can be given to villages when they adopt crop rotation with legumes.
- Innovative techniques such as neem-coated urea (for slow release of fertiliser into the soil) need to be promoted by the government. There should be 45 kg urea bags instead of 50 kg ones, which will reduce the loss/excessive use of fertilisers by about 10-20%, especially by small and marginal farmers.
- Soil sample collection, testing, and printing at the district level is significantly positively influenced by fertiliser use, number of bank accounts, net sown area, number of soil testing labs, and households with mobiles.
- Other policies, such as water exploitation, electricity, etc., should be in line so that crop diversity can protect soil health in the long run.

Chapter - I

Introduction

According to the latest estimates, 96.40 Million Hectare (Mha) area of India is undergoing the process of land degradation, i.e., 29.32% of the Total Geographic Area (TGA) of the country during 2011-13. It increased from 94.53 Mha (28.76% of the TGA) in 2003-05 (GoI, 2016). The states of Rajasthan, Maharashtra, Gujarat, Jammu & Kashmir, Karnataka, Jharkhand, Odisha, Madhya Pradesh, and Telangana contribute 24% of the degraded area in the country. Water erosion is the highest contributor (10.98% in 2011-13 and 10.83% in 2003-05), followed by vegetation degradation (8.91% in 2011-13 and 8.60% in 2003-05) and wind erosion (5.55 % in 2011-13 and 5.58 % in 2003-05). The area under desertification (arid, semi-arid, and dry sub-humid regions of the country) during 2011-13 was 82.64 Mha, whereas during 2003-05, it was 81.48 Mha. Thus, there is a cumulative increase of 1.16 Mha area under desertification. The most significant processes of desertification in the arid region have been observed to be wind erosion and, in semi-arid and dry sub-humid regions, vegetation degradation and water erosion (GoI, 2016). These estimates are much higher than the earlier assessment from the National Remote Sensing Agency (NRSA), which estimated the extent of land degradation in India at about 35.5 million hectares in 1999 (Reddy, 2003). This is mainly due to changes in definitions as well as the inclusion of forest degradation (vegetation degradation).¹

The pace of degradation has gone up during the post-green revolution period due to the intensive use of chemical inputs (Table 1). India has a long history of soil conservation interventions and programmes. From the early 20th century onwards, though, soil conservation was not among the policy priorities until the 1980s. As the extent of degradation increased over the years, soil conservation gained policy attention (refer to appendix table on Policies). However, a more focused approach to soil management has been adopted only after 2014-15. This coincided with the declaration of the international year of soils by the UN. Soil health is a holistic concept that includes the chemical, physical, and biological health of the soil (Figure 1). In line with the UN resolution, the ministry of agriculture introduced the Soil Health Card (SHC) scheme on December 5, 2015. The scheme is an improvement on the earlier National Project on Management of Soil Health and Fertility that was launched during 2008-09.

Fig1: Measure of Soil Health

¹The new estimates refer to process of degradation, which includes heavy degradation and degraded lands, while the earlier estimates included only degraded lands. Besides, the earlier estimates provided forest degradation assessment separates. If these definitional differences are taken into account, the estimates become comparable.

Table 1: Consumption of Fertilisers in million tonnes in India

Year	Million tonnes				NPK ratio		
	N	P	K	Total	N	P	K
1956	0.1	0.0	0.0	0.1	10.4	1.3	1
1981	3.7	1.2	0.6	5.5	5.9	1.9	1
1991	8.0	3.2	1.3	12.5	6.0	2.4	1
2001	10.9	4.2	1.6	16.7	7.0	2.7	
2011	16.6	8.0	3.5	28.1	4.7	2.3	1
2015	16.9	6.1	2.5	25.6	6.7	2.4	1

India is consuming about 25.6 million tonnes of fertilisers, mostly Nitrogen (17 million tonnes) followed by Phosphorous (6 million tonnes) and Potassium (2.5 million tonnes). The current NPK ratio is 6.7:2.4:1, which is highly skewed towards Nitrogen as against the ideal ratio of 4:2:1 (Table 1). There are wide variations across the states in terms of fertiliser use per acre and NPK ratios (Figure 2). India is spending about Rs. 1 lakh crore on fertiliser subsidy. It is estimated that the subsidy amount is about Rs. 6,500/ha of the net cropped area and about Rs. 7,000/farmer, resulting in excessive use of fertilisers, especially Nitrogen, at the cost of micro nutrients and manure (Figure 3). As a result of the excessive and unbalanced use of fertilisers, the amount of food grain produced per kg of fertiliser applied declined from 13 kg in the 1970s to just 4 kg by 2010. In order to promote balanced use of fertilisers, the Government of India introduced the Soil Health Card Scheme across India.

Fig. 2: State-wise Fertiliser use (kg/ha) and N:K Ratio

The SHC scheme was approved for implementation for the remaining period of the 12th plan. The scheme aims to provide SHC to all farmers in the country at a regular interval of two years in order to enable the farmers to apply recommended dosages of nutrients to realise improved and sustainable soil health and fertility status and reap higher profits. The scheme has the following objectives:

- a) To improve soil quality and profitability to farmers
- b) To generate employment for the rural youth
- c) To improve timelines in the analysis of soil samples.
- d) To introduce a single window approach from collection to the issue of SHC minimising delays and maximising convenience for farmers.
- e) To provide online delivery of SHC to farmers using the soil health card portal.
- f) To provide soil testing facilities to farmers at their doorstep.

Fig 3: Progressive expansion in the occurrence of nutrient deficiency

Under the SHC scheme, the cropped area was divided into grids of 10 ha for rainfed and 2.5 ha for irrigated and only one soil sample from each grid was taken. The test results will be distributed among all the farmers whose area was falling under the grid. Based on the grid system, the total 14.1 crore hectares of the net cropped area requires a total of 3.46 crore grid samples (with 52% of the area under rainfed [7.3 crore ha] with 73 lakh grid samples and 48% area under irrigated [6.8 crore ha] with 2.72 crore grid samples). The target is to cover 1.73 crore grid samples per year. This comes to an average of 25,000 grid samples per district/year and 29 grid samples per village/year. With this, all 10.39 crore farmers will be covered in two years. Every year, 5.2 crore farmers need to be covered.

Under cycle 1, 2.54 crore samples were collected, 2.36 crore samples tested, 9.62 crore soil health cards printed, but only 9.33 crore SHCs distributed. This indicates that 100% target was archived in sample collection, 93% of the target was achieved in soil testing, but only 80% of the target was achieved in SHC printing. 97% of the SHCs printed were distributed among the farmers as on September 24, 2017. However, the progress is highly skewed. Some states, such as Karnataka, Tamil Nadu, Chhattisgarh, Uttar Pradesh, Maharashtra, Telangana, and Andhra Pradesh, were better performers compared to other states.

Chapter –II

Soil Health in India: Review of Experience

Though there are a number of studies on the ill effects of poor soil quality across different states of India, they mostly focus on micro situations. There are no all-India studies on the impact of improved soil health. Of late, some studies have assessed the impact of soil health management programmes in Karnataka, Andhra Pradesh, Bihar, Gujarat, etc., where state governments have initiated the programmes viz. Bhoochetana in Karnataka and Krishi Mahostav in Gujarat. Here, the experiences of these initiatives are reviewed briefly (details of the studies are presented in appendix table 2).

Studies have clearly shown that farmers do adopt soil management strategies (Reddy, 2011). Most of these practices are based on their rich experience and vast knowledge of location-specific conditions. Combination of chemical fertilisers and FYM was a predominant practice, along with mixed cropping and legume cultivation. This shows that farmers understand the role of FYM and other organic manure. The analysis also showed a positive sign of the emerging organic markets. The availability of FYM is limited; promotion of fertiliser use (advertisements) and huge subsidy force farmers to use more chemical fertilisers. Farmers have indicated that recommendations provided by scientists without proper soil testing do not hold well for the conditions they are working in.

A study of impact assessment of SHC in three districts of Bihar observed that there is a large gap between recommended and actual application of fertilisers, especially in the case of urea. Despite the recommendations provided by SHC, farmers fail to adopt them (Fishman, et al., 2016). The main reasons for this include: i) farmers didn't understand the contents of the SHC; ii) farmers didn't find the soil analysis and fertiliser recommendations to be reliable or compelling; iii) other factors such as cost, liquidity, or timely availability of specific fertilisers were an added constraint.

On the other hand, studies have shown that when awareness programmes are followed up with supporting programmes such as inputs etc., soil improvements and increased crop yields are conspicuous. For instance, the Bhoochetana programme in Karnataka introduced direct benefit transfer in fertiliser subsidy to increase efficiency and strengthen fertiliser supply chain along integrated nutrient management with emphasis on organic fertilisers. Under this programme, the Karnataka government supplied micro nutrients at 50% subsidy. The study estimated that total benefits with soil health mapping and soil-test-based fertiliser recommendations along with improved practices would be Rs.4.33 lakh crore, against the estimated cost of Rs 0.254 lakh crore (ICRISAT research report IDC-6). The benefit-cost ratio would be 17:1. Besides economic benefits, there is employment generation and several environmental benefits, including enhancing the sustainability of Indian agriculture, which will be additional benefits.

Similarly, surface drainage technology for saline land reclamation was observed to be technically viable, economically feasible, and socially acceptable. The study by Raju, et al., (2015) clearly showed that land use was intensified, cropping patterns changed in favour of more remunerative crops, and crop yields increased with the use of soil health cards. There was a significant reduction in the maximum and minimum salinity. The mean yield of all crops grown significantly increased to the extent of 186%. Increase in net income was largely related to the increase in crop yield due to soil improvements.

A significant increase in net income from off-season crops was also observed. The benefit-cost ratio was more than one, and the value of land increased as well (Raju, et.al., 2015).

In a study on on-farm trails in eight districts of Andhra Pradesh, it was found that balanced nutrient treatment in the widespread multi-nutrient (including micro nutrients) deficient soils has resulted in significantly higher yields. Balanced nutrition while increasing crop yields maintained plant nutrient composition. Post-harvest soil testing in the Nalgonda district showed higher contents of soil organic carbon and available nutrients such as P, S, B, and Zn in plots with balanced nutrition treatment. In the absence of balanced nutrition, farmers were losing 8% to 102% of current yields in season 1 and 15% to 24% in each of the succeeding three to four seasons (Chander, et al., 2014)

2.1 Soil Health in India

According to the report “Degraded and Waste Lands of India” by the Indian Council for Agricultural Research (ICAR) and the National Academy for Agricultural Sciences, of the 141 million hectares of total geographical area, about 328.2 million hectares is under cultivation. Of this, about 100 million hectares-or 70%-is heading down a path where it will be incapable of supporting farming. Farmers are making the soil work more, growing two or more crops a year instead of one without proper soil health management. This unplanned intensification is exacerbating nutrient shortages and changing soils' chemical composition. Levels of organic carbon in soil are dropping across the country, making soils more vulnerable to erosion and possibly resulting in the number of earthworms falling. Not only do these excesses and imbalances reduce the productivity and life of soils, they are now resulting in poor nutritional value of our food. If the soils are deficient in some nutrients, so are the food crops grown on them. Pharma companies have consequently started adding Zinc, Copper, Selenium, Chromium, etc. to fortify their vitamin tablets.

2.2 Soil Nutritional Status

Indian Institute of Soil Science, (IISS) data shows that large parts of India are deficient in two or more critical nutrients. Regions such as the Indo-Gangetic plains-Punjab, Haryana, Uttar Pradesh, and Bihar-which produce nearly 50% of our grains and feed about 40% of our population are seeing multiple nutrient deficiencies. The reason is the imbalanced use of organic and inorganic inputs. In the past, farmers used to plough the stalks left standing on the field after the harvest, cow dung, etc. back into the soil. This ensured that nutrients taken out of the soil were replenished. The green revolution that started in the 60s changed all that. High-yielding crop varieties need more water and nutrients, spanning macro nutrients such as Nitrogen and Phosphorus to micro nutrients such as Copper and Boron. However, due to imbalanced fertiliser use, hardly any of these nutrient cycles are being completed. Farmers today use more of Urea (Nitrogen), some Potassium and Phosphorus, but not much else. Further, they choose fertilisers more by affordability and availability than what the soils need. Apart from retarding growth in yields, this unbalanced use has also damaged soils. Too much urea, for instance, turns soils acidic. The ICAR report estimates that 6.98 million hectares, or 2% of India's total geographical area, have acidic soils. These are mostly in North-East India, south Chhattisgarh, and Kerala. Another 6.7 million hectares are salt-affected. In the absence of historical data on nutrients, the degree of decline cannot be ascertained. These soils are increasingly incapable of supporting agriculture.

2.3 Soil Organic Carbon

Soils are changing from fine to hard. In healthy soils, crop residues are transformed by earthworms and other soil fauna into soft and spongy organic matter called humus (or organic carbon). This soaks up water, creates an environment where soil fauna such as earthworms thrive, and binds the soil's three constituents: sand, silt, and clay. Without humus, the soils compress and harden. Farmers, in a hurry to plant the next crop, burn their fields to clear stalks left standing after the harvest. Cow dung is scarcer. That's partly because with mechanisation, fewer households keep bullocks. The decline of grazing grounds has meant only households large enough to afford fodder can keep milch animals. The recommended amount of farmyard manure is 5-10 tonnes per hectare, whereas farmers add anywhere between zero to five tons. Poorer farmers sell manure instead of ploughing it into their fields. Or, they use dried dung as fuel. The outcomes are predictable. IISS estimated that humus depletion in the top 0-15 cm is nearly 50%, although it can occasionally be as high as 60-70% in some soil types. There is 10-20% loss in the 15-100 cm below that top layer. As humus falls, properties of the soil change. For example, its ability to absorb water reduces, resulting in erosion. The ICAR report estimates that a total of 126 M ha is suffering from various degrees of water erosion. Of this, 0.68 million hectares are seeing severe, very severe, and extremely severe erosion. Another 0.2 million hectares is seeing "moderately severe" erosion. The report estimates that very severe and severe wind erosion occurs in 16% of India's total geographical area. Both these processes contribute to desertification-soil turning into a desert. About 81 million hectares, or 25% of it, is experiencing desertification, says a 2009 ISRO paper titled 'Desertification/land degradation status mapping of India'. Anecdotal information suggests that the number of earthworms and other soil insects is falling.

Fig. 4: Fields effected by alkalinity

2.4 Soil Health Card to Improve Soil Health

The Soil Health Card programme is trying to address these problems. However, it is a crisis that we are just starting to deal with. One of the few exceptions is the government's switch to Nutrient-based Subsidy. As a response to growing micro nutrient shortages, it falls short, partly because the price signals it sends out about the nutrients to use are national while the shortages are local; partly because the government has decontrolled prices of P and K while holding on to prices of N. As a result, every time P and K prices soar, farmers dump more urea than other fertilisers; and partly because it doesn't extend the subsidy to organic manure or all 20 micronutrients: it only covers Boron and Zinc. We do not have pan-India authentic baseline data about soil nutrient concentrations. The Indian Institute of Soil Science (IISS) data, as shown in (Map 1, Map 2, Map 3 and Map 4), was derived from samples collected by soil-testing labs. The quantum of deficiency in these maps has been calculated by gauging the minimum concentration of a nutrient required (for that soil type) for above normal yields. However, macro nutrient deficiency has been mapped for only 18 odd states.

In the case of micro nutrients, we know their concentration in even fewer states with a very limited sample. There is no authentic data on pan-India soil microbial population and activity. Tamil Nadu, Gujarat, and Punjab are better in this regard than other states. India has 140 million farmers, but our labs cannot process more than 8 million soil samples in a year. It will take us 15 years, at this rate, to tell every farmer what their soil needs. The larger question is about the soils' physical condition: can they support the hydrological cycle? And for that, you need humus and soil biota. Earthworm populations, organic carbon levels, and yields are the highest when a combination of balanced fertilisation and farmyard manure is used. However, India doesn't have enough cow dung. Organic waste can be used instead of cow dung, but we need to figure out better ways to segregate organic and inorganic waste and get it across to fields. Various state governments are encouraging composting pits where agricultural and household wastes can decompose into manure that is then ploughed back into the soil.

Map 1: Status of Nitrogen and Zinc availability in Indian soil (source: Economic Times)

Map 2: Status of Phosphorus and Iron availability in Indian soil

Map 3: Status of Potassium, Copper and Manganese availability in Indian soil

Map 4: State-wise consumption of Fertilisers

Map 5: Changing chemical composition of soils

Chapter - III

Policy Interventions over the Years: International and National

Though soil management practices and policies are as old as agriculture, formal land management policies at the international level were initiated in the early 1980s with the first World Soil Charter (WSC) adopted by the FAO (Food and Agriculture Organisation) member countries in 1982 under the global soil partnership. This was followed up with a number of international policy initiatives such as the Intergovernmental Panel on Climate Change (IPCC) in 1988. Soil management received a flip with UNCCD (United Nations Convention to Combat Desertification), focusing on drought mitigation and combat desertification. This was followed by the Rio conference, which provided 27 principles to guide countries towards sustainable development. Following this, a number of conventions and protocols were initiated to help soil management at the global level. The important ones among them include:

3.1 Global policies

- a. 1997: The Kyoto Protocol is an international treaty that extends the 1992 United Nations Framework Convention on Climate Change (UNFCCC), which commits state parties to reduce greenhouse gas emissions based on the scientific consensus that (a) global warming is occurring and (b) it is extremely likely that human-made CO₂ emissions have predominantly caused it.
- b. 2000: MDG's - Soil management and prevention of desertification. Implementation of soil erosion control (by wind and water) by planting windbreaks and cover crops; improvements in soil fertility with agro-forestry systems, cover crops, and conservation of ground and surface water.
- c. 2005: The Millennium Ecosystem Assessment looked at the consequences of ecosystem change for human well-being. From 2001 to 2005, the MEA involved the work of more than 1,360 experts worldwide. Their findings provide a state-of-the-art scientific appraisal of the condition and trends in the world's ecosystems and the services they provide as well as the scientific basis for action to conserve and use them sustainability.
- d. 2008: This UNCCD policy brief "A Sustainable Development Goal for Rio+20: Zero Net Land Degradation" provides a snapshot of the world's land, explains the causes and impacts of land degradation, and suggests pathways to land-degradation neutrality. The brief reveals that sustainable land use is a prerequisite for ensuring future water, food, and energy security. Given the increasing pressure on land from agriculture, forestry, pasture, energy production, and urbanisation, urgent action is needed to halt land degradation.
- e. 2011: The Global Soil Partnership for Food Security and Climate Change Mitigation and Adaptation (GSP) brings together international, regional, and national organisations that are working in the area of soil protection and sustainable management. Launched by the Food and Agriculture Organisation of the United Nations in Rome on September 7, 2011, the partnership

aims to implement the provisions of the 1982 World Soil Charter and raise awareness and motivate action by decision-makers on the importance of soils for food security and climate change adaptation and mitigation.

- f. 2012: Sustainable Development Goal 15 of the 2030 Agenda aims to “protect, restore, and promote sustainable use of terrestrial ecosystems, sustainable forest management, combat desertification and halt and reverse land degradation and halt biodiversity loss”.
- g. 2013: The Intergovernmental Technical Panel on Soils (ITPS) was established in the first Plenary Assembly of the Global Soil Partnership held at FAO Headquarters on June 11 and 12, 2013. The ITPS is composed of 27 top soil experts representing all the regions of the world. The main function of the ITPS is to provide scientific and technical advice and guidance on global soil issues to the Global Soil Partnership primarily and to specific requests submitted by global or regional institutions. The ITPS will advocate for addressing sustainable soil management in the different sustainable development agendas.
- h. 2015: The International Year of Soils, 2015 (IYS 2015), was declared by the 68th session of the United Nations General Assembly on December 20th, 2013, after recognising December 5 as World Soil Day. The purpose of the IYS is to raise awareness worldwide about the importance of soils for food security, agriculture, as well as in mitigation of climate change, poverty alleviation, and sustainable development.

3.2 National Policies

In India, formal soil and water interventions were under way since 1900 under the British rule. Post independence, soil management has become part of five year plans from the first plan onwards (refer to the appendix table). A number of initiatives have been taken up at the national level over the years. Apart from the soil reclamation programmes during the 1970s and 80s and soil and water conservation programmes going on over the years (mainly watershed development programmes), no specific soil management programme was initiated at the national level. For the first time, during the 11th plan, National Mission for Sustainable Agriculture (NMSA) was introduced as a part of the National Action Plan on Climate Change (NAPCC). The National Project on Management of Soil Health and Fertility and the Rainfed Areas Development Programme (RADP) was also introduced. It recommended conservation agriculture, integrated nutrient management, carbon sequestration, erosion control, saline and alkaline soils management, legislation for soil protection, development of remote sensing and GPS (Global Positioning System) based on Decision Support System (DSS) and amelioration of polluted soil to rejuvenate deteriorated soils. This was followed up in the 12th plan by introducing a new scheme, 'National Project on Management of Soil Health & Fertility' (NPMSH&F). Under this scheme, soil health cards were introduced along with strengthening of soil testing labs and expanding their testing capacity. Furthermore, the Nutrient-Based Subsidy (NBS) system was introduced. In the recent years, some states, such as Karnataka, Gujarat, etc., have introduced soil management programmes, such as Bhoochetana and Krishi Mahotsav programmes. These programmes have provided insights and learnings related to the central schemes.

Table 2: Soil Quality and Quantity Indicators

Indicator	Poor	Medium	Good
Earthworms	0-1 worms in shovelful of top foot of soil. No casts or holes	2-10 in shovelful few casts, kholes or worms	10+ in top foot of soil. Lots of casts and holes in tiled clods. Birds behind tillage
Organic matter colour	Top soil colour similar to sub-soil colour	Surface colour closer to sub-soil colour	Top soil clearly defined, darker than sub-soil
Roots/residue	No visible residue or roots	Some residue, few roots	Visible residue and roots
Surface compaction	Wire breaks or bends when inserting surveyors flag	Have to push hard, need fist to push flag in	Flag goes in easily with fingers with twice the depth of the plow layer
Soil tilth mellowness friability	Looks dead. Like brick or concrete cloddy. Either blows apart or hard to pull drill through.	Somewhat cloddy. Balls up, rough pulling seedbed.	Soil crumbles well, can slice through, like cutting butter. Spongy when you walk on it.
Erosion	Large gullies over two inches deep joined to others, thin or no top soil, rapid run-off the colour of the soil	Few rills or gullies, gullies up to two inches deep. Some swift runoff, coloured water	No gullies or rills. Clear or no runoff.
Water holding capacity	Plant stress two days after a good rain	Water stress after a week	Holds water for a long period of time without puddling.
Drainage infiltration	Water lies for a long time and evaporates more than drains, always very wet aground	Water lies for a short period of the time and eventually drains.	No ponding runoff water; moves through soil steadily. Soil not too wet, not too dry.
Crop condition	Problem growing throughout season, poor growth, yellow or purple colour	Fair growth, medium green colour	Normal, healthy dark green colour, excellent growth all seasons across fields
pH nutrient holding capacity	Hard to correct for desired crop Soil test values dropping with more fertiliser applied than crops use	Little or slow change	Proper pH for crop Fair growth, spots in field different, medium green colour.

Chapter IV

Objectives and Methodology

4.1 Objectives:

As the SHC scheme has completed more than 2 years of implementation, the ministry has initiated a nationwide impact assessment with the following objectives:

- ☞ To examine the design of the SHM/ SHC scheme in terms of planning, implementation (role of state/ JDA/ADA/AO), inputs (staff, financial and other resources), activities (training, lab established and strengthened), outputs (SHCs printed and distributed to farmers).
- ☞ To assess the modalities of delivery of the SHM / SHC scheme in terms of procurement, sample collection, testing, SHC printing, and disbursal.
- ☞ To assess the level of utilisation of SHCs by the farmers across farm size classes in irrigated and rainfed situations.
- ☞ To assess the impacts of SHC scheme on judicious use of fertilisers, biofertilisers, organic fertilisers, soil health, cropping choice, cost reduction, farm profitability, and sustainability.
- ☞ To recommend improvement in the overall design of the programme.

4.2 Methodology

Both qualitative and quantitative approaches were used to assess the impact of the Soil Health Card Scheme. Qualitative information in the form of stakeholder interviews across the study states, expert opinion gathering at the national and state levels through workshops, and interactions with the progressive farmers were carried out. At the quantitative level, both secondary and primary data was collected at the national, state, and farmer levels. Secondary data mainly pertains to financial and physical achievements of the SHC scheme over the years, infrastructure availability, number and capacity of Soil Testing Laboratories, and utilisation capacity were collected and analysed.

A systematic sample has been drawn for the impact assessment at the farmer level. Care was taken to represent the whole country and different agro-climatic conditions. The sampling details are presented as follows.

4.2.1 Selection of States, Districts, and Blocks

For the secondary data, all the states were covered to understand the number of SHCs issued and farmers covered, financial and physical progress of the inputs, activities and outcomes of the SHC scheme. For primary data collection, all major states (25 major states) were divided into five zones for drawing the sample. From each zone, 2-4 states were selected, and a total of 16 states were selected for primary data collection. At the next level, two districts from each state (three districts in case of large states and one district in case of small states) were selected, i.e., a total of 29 districts. From each district, two blocks were selected randomly, i.e., a total of 65 blocks. From each block, two to three villages were selected randomly, i.e., a total of 199 villages. From each village, 16 households were selected for collecting detailed information with the help of a structured questionnaire (refer to appendix). Overall, 3,184 households were covered at the national level (Table 2).

Due care was taken to represent a few farmers who received SHC at least a year before the survey date in order to understand the impact. Hence, the sample size varies a little according to the availability of the oldest adopters in villages. Due care was also taken to cover small and marginal farmers, SC/ST farmers, educated youth, and beneficiaries under demonstrations (Table 3). The impact of SHC was assessed using the before and after methodology and also using with and without wherever possible.

We have also classified respondents based on the period of holding SHC, oldest (received SHC more than 1 year ago), old (6 months ago), new (4 months ago), and recent (less than 4 months) to assess the impacts over the period.

All the states were grouped into zones based on standard classification. Northern Zone comprised Chandigarh, Delhi, Haryana, Himachal Pradesh, Jammu and Kashmir, Punjab, and Rajasthan; North-Eastern Zone comprised Assam, Arunachal Pradesh, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim, and Tripura; Central Zone comprised the states of Chhattisgarh, Madhya Pradesh, Uttarakhand, and Uttar Pradesh; Eastern Zone comprised Bihar, Jharkhand, Odisha, and West Bengal; Western Zone comprised Dadra and Nagar Haveli, Daman and Diu, Goa, Gujarat, and Maharashtra; Southern Zone comprised Andhra Pradesh, Karnataka, Kerala, Lakshadweep, Puducherry, Tamil Nadu, and Telangana.

Table 3: Details of the sampling framework

Zone	States	Districts	Blocks	Villages	Total Sample Farmers
Central	2	3	6	20	320
East	2	3	7	15	240
North-East	4	4	10	28	448
North	3	8	11	54	864
South	3	7	24	50	800
West	2	4	7	32	512
Total	16	29	65	199	3184

For primary data collection, the following states were covered under each zone:

- **Central Zone**
 - Chhattisgarh
 - Madhya Pradesh
 - Uttar Pradesh
 - **East Zone**
 - Jharkhand
 - West Bengal
 - **North-East Zone**
 - Nagaland
 - Arunachal Pradesh
 - Assam
 - Sikkim
- **North Zone**
 - Haryana
 - Punjab
 - **South Zone**
 - Andhra Pradesh
 - Telangana
 - Karnataka
 - **West Zone**
 - Gujarat
 - Maharashtra

Furthermore, states were divided into developed and less developed states based on development indicators taken from the planning commission. The developed states include Andhra Pradesh, Telangana, West Bengal, Haryana, Maharashtra, and Gujarat. Less developed states include Uttar Pradesh, Nagaland, Sikkim, Arunachal Pradesh, and Assam. This would help in understanding whether development has any role in the effectiveness of the SHC at the farmer level.

4.2.2 Structure of Data from Farmers:

Structured questionnaires covering all the details pertaining to SHC were canvassed among the sample farmers. Besides, focus group discussions were conducted in the sample villages to get feedback from the communities. These groups include key informants and farmers not covered under individual survey and farmers who have not received the soil health card.

All the indicators collected from field survey were classified as inputs (financial and physical inputs under the project), activities (different activities organised under the scheme), outputs (actual outputs of the project), outcomes (whether generated outputs were used by the farmers), and impacts (what are the ultimate benefits to the farmers).

4.2.3 Indicators of inputs of the SHC Scheme

- i. Procurement of maps (patwari / pahani maps) and field instruments (with GPS) and deployment of staff for collection of samples from farmers' fields.
- ii. Methodology of soil sample collection (is the 10 ha grid for rainfed and 2.5 ha for irrigated adequate? Or are any improvements needed?).
- iii. Status of soil testing facilities, including soil testing labs and mobile soil testing labs (geographical distribution, adequacy, access, and reachable)
- iv. Availability of equipment (colour Photostat machine, standard paper for printing, and printing machines) and mechanisms for distribution of soil health cards to farmers.
- v. Time lag between the release of funds and actual utilisation.
- vi. Annual action plan versus implementation.
- vii. Effectiveness of partnerships (operationalisation of the Public Private Partnership mode).
- viii. Technical help from agricultural officers and allied sectors.
- ix. Number of skilled staff engaged and whether there is any shortage of staff
- x. Number of awareness camps and trainings and effectiveness of the campaign
- xi. Reaching out to remote villages (in terms of samples collected and SHCs distributed and used)
- xii. Identifying problematic soils, recommendations, and application of soil amendments

4.2.4 Indicators of Activities under SHM

- i. Number of AEOs/ ToTs trained under the scheme.
- ii. Number of meetings, trainings, exposure visits, and demonstrations conducted.
- iii. Number of soil testing laboratories strengthened (analysing capacity and utilisation).
- iv. Number of soil testing labs equipped with micro nutrient testing facilities.
- v. Number of referral labs established.
- vi. Number of STLs equipped with Soil Test-based Crop Response (STCR).
- vii. Number of farmers covered, samples collected, tested, and SHC printed.

4.2.5 Indicators of Outputs of the Scheme

- i. Number of SHCs distributed among farmers.
- ii. Number of SHCs distributed among small and marginal farmers and SC/ST farmers (whether main field was located in their farm)

4.2.6 Indicators of Outcomes

- i. Awareness about SHCs among the farmers
- ii. Sources of information about soil health card and reliability
- iii. Knowledge gain through attending demonstrations
- iv. Understanding the content and SHC's recommendations
- v. SHC portal usefulness for farmers; rating from 1 to 5 (1 is bad; 5 is excellent)
- vi. Use of fertilisers according to the SHC recommendations
- vii. Crops recommended in SHCs and actual crops sown by farmers
- viii. Use of organic manure and biofertilisers by farmers
- ix. Integrated Nutrient Management practices followed by farmers
- x. Expectations of the farmers
- xi. Overall opinion of farmers on SHC.

4.2.7 Indicators of Socio-economic Impact

- i. Perceptions of the farmers towards the soil health card
- ii. Increase in crop yields, reduction in fertiliser use and cost, and increase in profitability.
- iii. Economic returns for farmers

4.2.8 Information From Key Stakeholders and Secondary Data

- i. Scheme design (determining whether current grid of 10 ha for rainfed and 2.5 ha for irrigated is appropriate), components, activities, and outputs suitable to local implementation.
- ii. Time lag in fund allocation, release, and utilisation at different stages.
- iii. Gaps in physical and financial progress.
- iv. Comparative analysis of the best performing states and other states in actual utilisation of SHCs by farmers.
- v. Process monitoring and evaluation (institutional and administrative procedures, roles and responsibilities, maintenance of records) were analysed for learning across states.

Chapter V

Soil Health Card Scheme: Design, Coverage, and Impact

5.1 SHC Design and Status (Inputs/Activities and Outputs):

Under the SHC scheme, the cropped area was divided into grids of 10 ha for rainfed and 2.5 ha for irrigated area, and only one soil sample was taken from each grid. The test results will be distributed to all the farmers whose area was falling under the grid. Based on the grid system, out of the total 14.1 crore hectares of net cropped area, with 52% of area under rainfed (7.3 crore ha) and 73 lakh grid samples and with 48% area under irrigated (6.8 crore ha) and 2.72 crore grid samples, a total of 3.46 crore grid samples in two years will be the target. Every year, 1.72 crore grid samples need to be covered. This comes to on average 25,000 grid samples per district/year and 29 grid samples per village/year. With this, all 10.39 crore farmers will be covered in two years. Every year, 5.2 crore farmers need to be covered. Until now, the total samples entered are 1.56 crore, the number of farmers covered is 4.82 crore, samples tested are 1.23 crore and SHCs printed are 3.05 crore. This indicates that 46% of the total farmers were covered under the scheme at a national level, but only 29.3% received SHCs.

Samples Collected

State-wise samples collected were presented in map 6. It has been observed that the highest number of samples collected was in Kerala, with a maximum of 969 samples collected per 1,000 ha. Kerala is followed by Tamil Nadu, Gujarat, Uttar Pradesh, and Bihar, with 247 to 544 samples collected per 1,000 ha. Northern states such as Jammu & Kashmir, Himachal Pradesh, Haryana and southern states of Telangana, Andhra Pradesh, and Maharashtra showed a moderate number of sample collection ranging between 157 and 246 samples/1,000 ha. A semi-moderate number of samples-i.e., 27-156 samples per 1,000 ha-was collected from most of the central, eastern, and north-eastern states such as Odisha, Madhya Pradesh, Arunachal Pradesh, and Assam. The lowest number of samples-i.e., 0-26 samples per 1,000 ha-was collected in Manipur and the NCT of Delhi.

State-wise samples collected per 1,000 cultivators Map 7 show that Andhra Pradesh, Telangana, and Haryana came into green with higher samples collected per 1,000 cultivators. But Himachal Pradesh and Bihar came into red, with the lowest samples collected per 1,000 cultivators. Uttar Pradesh and West Bengal turned from light green to yellow.

Samples tested

State-wise samples tested per 1,000 ha are presented in map 8. It has been observed that the highest number of samples tested per 1,000 ha was in Kerala (727 samples tested per 1,000 ha). Kerala is followed by Tamil Nadu, Andhra Pradesh, Gujarat, and Haryana, with 206-316 samples tested per 1,000 ha. It can be seen that in most of the states (Telangana, Maharashtra, Karnataka, Madhya Pradesh, Chhattisgarh, etc.), the tested samples were in the range of 89-205 per 1,000 ha. In the rest of the states, viz. Rajasthan, Punjab, Jharkhand, and Jammu & Kashmir, only 24-88 samples were tested per 1,000 ha. The lowest number of samples tested-i.e., 0-23 samples tested per 1,000 ha-is seen in NE states Assam and Manipur.

Map 6: State-wise samples collected per 1000 ha

Map 7: State-wise samples collected per 1000 cultivators

Map 8: State-wise samples tested per 1000 ha

Map 9: State-wise samples tested per 1000 cultivators

Soil samples tested per 1,000 cultivators (Map 9) show that Gujarat turned from light green (moderately high) to green (high). Karnataka, Telangana, Madhya Pradesh turned from yellow (medium) to light green (moderately high). On the other hand, Bihar turned from low to lowest (Map 8).

SHC printed

State-wise SHCs printed per 1,000 ha is depicted in map 10. It has been observed that the highest number of SHCs printed was in Kerala and Tamil Nadu, with above 1,000 SHCs printed per 1,000 ha. These two states are followed by Andhra Pradesh, Chhattisgarh, Himachal Pradesh, Uttarakhand, and Bihar, with about 687-1002 SHCs printed in the states. Rest of the states, such as Telangana, Maharashtra, Odisha, West Bengal, Uttar Pradesh, and parts of Jammu & Kashmir displayed a moderate range of SHCs printed per 1,000 ha, i.e., in the range of 361-686. The low number of SHCs (i.e., 73-360 numbers per 1,000 ha) was seen in western states such as Rajasthan, Gujarat, Madhya Pradesh and northern states such as Punjab as well as Jharkhand. It was seen that the lowest number of SHCs printed, i.e., 0-72 SHCs printed per 1,000 ha was in North-eastern states, with the exception of Tripura and Nagaland, which showed about 361-686 SHCs printed per 1,000 ha.

A comparison of soil health cards printed per 1,000 ha and per 1,000 cultivators (Map 11) shows that there was a positive shift in Andhra Pradesh from light green (moderately high) to green (high), Telangana shifted from yellow to green (high). Gujarat and Madhya Pradesh shifted from low to medium, and Haryana moved from medium to moderately high. Bihar shifted from low to lowest (Map 10).

Soil Health Cards Distributed

State-wise SHCs distributed per 1,000 ha can be seen from map 12. It has been observed that the maximum number of SHCs distributed was in Kerala and Tamil Nadu (above 1,000 SHCs distributed per 1,000 ha). Andhra Pradesh, Chhattisgarh, Bihar, Himachal Pradesh, and Uttarakhand also showed a positive result, with 687-1001 SHCs distributed per 1,000 ha. These states are followed by Karnataka, Telangana, Odisha, West Bengal, Uttar Pradesh., and North-East states such as Sikkim, Nagaland, and Tripura, with 355-686 SHCs distributed per 1,000 ha. Northern, Western, and North-western states showed low range (74-354) of SHCs distributed per 1,000 ha. The lowest number of SHCs-i.e., 0-70-was distributed in Assam, Arunachal Pradesh, Mizoram, and Manipur.

When compared to SHCs distributed per 1,000 ha, SHCs distributed per 1,000 cultivators (Map 13) in Maharashtra, Karnataka, and Telangana shifted from yellow (medium) to light green (moderately high); Gujarat and Madhya Pradesh shifted from moderately low to medium; on the other hand, Kerala shifted from green (high) to yellow (medium), and Bihar shifted from moderately low to low. Himachal Pradesh and Uttarakhand's relative positions declined as well. Overall, the figures indicate that south India performed better, followed by central and western India. The performance was poor in the North-East (Map 12).

Map 10: State-wise SHCs printed per 1000 ha

Map 11: State-wise SHCs printed per 1000 cultivators

Map 12: State-wise SHCs distributed per 1000 ha

Map 13: State-wise SHCs distributed per 1000 cultivators

District-wise Samples Collected, Tested and Printed

District-wise data on soil samples collected (Map 14), tested (Map 15), and printed (Map 16) shows that although south-central and western Indian districts performed better, there is quite a bit of variation within the states. Even some districts in most backward states such as Bihar, Uttar Pradesh, and North-Eastern regions performed well, whereas some districts in south India performed poorly. This reflects the efficiency of district-level officers and other stakeholders. Besides, infrastructure plays an important role in the program.

Map 14: District-wise samples collected per 1000 ha

Map 15: District-wise samples tested per 1000 ha

District wise Samples Printed Per 1000 Ha

Map 16: District-wise SHCs printed per 1000 ha

Map 17: State-wise number of soil-testing laboratories

State-wise soil testing laboratories can be seen from map 17. It has been observed that the highest number of soil testing labs are found in Maharashtra and Tamil Nadu, with 156-344 labs. Gujarat and Uttar Pradesh stand second with 79-155 labs. Eastern (Chhattisgarh, Jharkhand, and West Bengal) and North-Eastern states have the lowest number (0-16) of soil testing labs. A majority of states, such as Telangana, Andhra Pradesh, Maharashtra, Karnataka, Odisha, Madhya Pradesh, Bihar, etc., have 42-78 labs (moderate range). Kerala, Himachal Pradesh, Uttarakhand, and parts of Jammu & Kashmir have 17-48 labs (semi-moderate range).

A number of soil testing laboratories per million ha are shown in map 18. This shows high density in Kerala, followed by Tamil Nadu, Himachal Pradesh, Jammu & Kashmir, and Uttarakhand. The lowest density is in Rajasthan, West Bengal, and North-East. The density in Andhra Pradesh, Telangana, Maharashtra, Gujarat, Jharkhand, Bihar, Haryana, and Punjab is medium. District-wise labs are shown in map 19. In a few districts of Tamil Nadu, Kerala, Gujarat, Andhra Pradesh, Himachal Pradesh, Punjab, and Haryana, lab density for soil testing is high. However, in many of the north, central, eastern, and north-eastern districts, lab density for soil testing is quite low.

Map 18: State-wise soil testing labs per million hectares

District wise Soil Testing Laboratories per million cultivators

Map 19: District-wise STLs per million cultivators

5.2 Status of SHCs (Financial and Human Resource Allocation)

In this section, the progress in the implementation of the SHC scheme is looked at from four important aspects, viz. financial allocations, infrastructure, human resources, and distribution of SHCs.

Allocations have been increasing over the years. In 2014-15, allocation under the project was Rs.13.68 crore, which increased to Rs.88.53 crore in 2015-16 and reached Rs.169 crore in 2016-17. But the gap between allocation, release, and expenditure is increasing as well (Table 4A). The north zone has the highest gap. This could be attributed to the low capacities of spending. During the field visits, it has been observed that the block- and district-level agricultural officers are involved in multiple tasks assigned by state governments, and hence there was a need to have separate staff for soil health card management, especially in setting up, upgradation, and maintenance of soil testing labs. Field observations also indicated that there was frequent transfer of staff in soil testing labs to the field level and vice versa. Postings in soil testing laboratories is not a priority for many agricultural officers, although a few female agricultural officers have been interested in postings at STLs. To improve the quality of soil testing, there should be a separate cadre for working in soil testing labs. It has been indicated that female agricultural officers prefer working with STLs.

Table 4A: Financial allocations and expenditure

State - wise Funds allocation, Release and Expenditure under Soil Health Cards Scheme in India (in lakhs)													
Zone	2014 -2015			2015 -2016					2016 -2017 *				
	Release	Expenditure	% Expenditure (Based on Release)	Allocation	Release	Expenditure	%Release (Based on Allocation)	%Expenditure (Based On Allocation)	Allocation	Release	Expenditure	%Release (Based on Allocation)	%Expenditure (Based on Allocation)
South	218	218	100	1250	811	811	64	64	2331	1165	697	50	29
North	389	371	95	2787	2005	492	71	17	5324	2442	0	45	0
West	440	396	90	3348	2424	1817	72	54	6235	3117	1544	50	24
East	234	219	93	1123	815	706	72	62	2274	407	88	17	3
North east	88	37	42	344	194	51	56	14	768	0	0	0	0
INDIA	1368	1241	91	8853	6248	3877	71	44	16932	7132	2329	42	14

Note: * until November 2016

Table 4B: Zone-wise analysis of funds allocation, release, and utilisation

	2015 - 16			2016-17		
Zone	Allocation (Rs. Lakh)	Released as % of allocation	UC pending as % of released	Allocation (Rs. Lakh)	Released as % of allocation	UC pending as % of released
Central	2528	43	17	4593	98	78
Eastern	1094	77	58	2348	14	100
North	1844	48	35	3303	13	59
North - Eastern	292	64	0	1161	37	23
South	19 96	56	4	3829	56	23
Western	1846	17	0	4056	42	32
Total	9600	46	23	19290	50	54

Source: Govt. of India letter dated July 27, 2017. There may be some mismatch between table 4 and 4A, as 4A is the updated one.

Table 4B shows that overall, only 46% of the funds allocated were released in the year 2015-2016 and only 50% were released in the year 2016-17. Utilisation certificate was pending for 23% of the funds released in the year 2015-16 and pending for 54% of the funds released in the year 2016-17. In the year 2015-16, fund release was better in the eastern zone, followed by the north-east and north zones, while in the year 2016-17 it was better in central and southern zones. The funds utilisation needs to be increased across all the zones.

There are 1,244 soil testing laboratories in India, of which 1,048 are static and the remaining 196 are mobile. About 90% are under state governments, and the remaining are managed by the fertiliser industry. In addition, about 7,000 mini soil testing kits were distributed to block-level agricultural offices across the states. The present analysing capacity is 1.78 crore samples per annum with a utilisation of 75.8%. The number of labs in India is 2016-17, as shown in Table 5.

In terms of analysing capacity (number of samples per annum), the South zone is at the top with about 7.0 million samples per annum, followed by the North zone (5.4 million samples) and the West zone (4.6 million samples). Among states, Tamil Nadu ranks first with 5.8 million samples, followed by Uttar Pradesh (4.1 million samples), Maharashtra (4.1 million samples), and Gujarat (1.4 million samples).

Table 5: Infrastructure (number of labs) in 2016-17

Number of Soil Testing Laboratories								Analysing	Number of	Capacity
Zone/States/UTs	State Govt.		Fertiliser Industry		Total			Capacity	Sample Sanalysed	% Utilisation
	Static	Mobile	Static	Mobile	Static	Mobile	Total	(`000 Nos.)	(`000 Nos.)	
East Zone	105	37	1	2	106	39	145	959.4	769.2	80.2
Arunachal Pradesh	5	3	-	-	5	3	8	9	7.9	87.3
Assam	7	4	-	-	7	4	11	84	60.8	72.3
Bihar	39	-	-	-	39	-	39	230	248.7	108.1
Jharkhand	8	-	-	-	8	-	8	40	10.7	26.7
Odisha	17	6	1	-	18	6	24	270	255.1	94.5
West Bengal	10	8	-	2	10	10	20	112	60.4	53.8
Manipur	4	4	-	-	4	4	8	40	1.4	3.4
Meghalaya	3	3	-	-	3	3	6	30	27.7	92.2
Nagaland	3	-	-	-	3	-	3	45	14.3	31.8
Sikkim	4	2	-	-	4	2	6	37	39.9	107.8
Tripura	2	4	-	-	2	4	6	35	17.5	50.1
Mizoram	3	3	-	-	3	3	6	27	25	92.6
North Zone	377	45	9	6	386	51	437	5444	4198	77
Haryana	35	3	2	-	37	3	40	365	247.9	67.9
Himachal Pradesh	11	4	-	-	11	4	15	125	124.4	99.5
Jammu and Kashmir	8	5	-	-	8	5	13	52	43.6	83.9
Punjab	54	12	2	3	56	15	71	631.5	282.1	44.7
Uttar Pradesh	255	18	5	3	260	21	281	4159.5	3404.6	81.9
Uttarakhand	13	3	-	-	13	3	16	106.5	95.2	89.4
Delhi	1	-	-	-	1	-	1	5	0.5	9.2
South Zone	157	32	35	4	192	36	228	6727.4	5503.2	81.8
Andhra Pradesh	55	5	27	1	82	6	88	413	345.8	83.7
Karnataka	56	-	6	2	62	2	64	295.7	194.8	65.9
Kerala	14	11	1	-	15	11	26	218	134.7	61.8
Tamil Nadu	30	16	1	1	31	17	48	5796.7	4823.5	83.2
Pondicherry	2	-	-	-	2	-	2	4	4.4	110.3
West Zone	348	59	16	11	364	70	434	4695.4	3046.6	64.9
Gujarat	132	2	4	1	136	3	139	1412	1199.1	84.9
Madhya	50	7	2	4	52	11	63	378	346.5	91.7
Pradesh Chhattisgarh	7	5	1	-	8	5	13	105	116	110.5
Maharashtra	123	23	8	4	131	27	158	2241.4	967.3	43.2
Rajasthan	34	22	1	2	35	24	59	536	402.7	75.1
Goa	2	-	-	-	2	-	2	23	15	65
India	987	173	61	23	1048	196	1244	1,78,27	1,35,17	75.8

Source: The Fertiliser Association of India. (16851); Andhra Pradesh Figures include Telangana

Capacity utilisation is more than 90% in Chhattisgarh, Pondicherry, Bihar, Sikkim, Himachal Pradesh, Odisha, Mizoram, Meghalaya, Madhya Pradesh and Uttarakhand, but less than 50% in Manipur, Jharkhand, Nagaland, Maharashtra and Punjab. Low capacity utilisation of funds and infrastructure could be due to lack of human resources. Since this is a new central scheme, state governments have taken the time to streamline and coordinate the activities, especially in north and eastern zones. The performance is better in southern and western zones, which is attributed to the efficient use of already

existing skilled workers, labs, and improvements in lab capacities, especially the setting up of mini labs, although focus group interactions reveal that the results of mini labs are not as accurate as those of full-fledged labs.

Focus group interactions with local agriculture officers indicated that there was a severe shortage of infrastructure (labs) and skilled workers. Although soil sample collection, testing, and distribution of soil test values is done by multiple agencies (including KVKs, SAUs and ICAR, and some private companies such as Nagarjuna Fertilisers), under the SHC scheme, the burden is exclusively on the State Department of Agriculture. Most of the soil sample collection work is carried out by contractual agricultural extension officers, who work under the block-level agricultural officers. In developed states such as Punjab, extension officers were provided with GPS-enabled tablets. There was no statistical database on the total staff involved in the soil sample collection and testing. As the soil sample collection was seasonal (only from April and May), all the staff below the block level was engaged in the soil sample collection. Soil testing requires specialised labs and skills that are not widely available at the block level and hence sent to district-level soil testing labs. Some estimates show that at the national level, the gap between requirement and availability is -148 %. The gap is the highest in the South zone, followed by west and north-east zones. The gap could be much higher if we include other activities as well, i.e., after sample collection drying, grinding, sieving, etc. Soil sample collection should be done in the campaign mode by involving the staff of all agricultural and allied department employees, as this activity is exclusively carried out during April-May. There is a need for developing a separate soil testing cadre and strengthening district-level soil testing labs with an estimated cost of Rs. 4-5 crore per district. Overall, it requires Rs. 2,800 to set up one state-of-the-art lab in each of 700 districts in India. Field-level observations indicated that the mini lab results are not too reliable, take a lot of time, and are also not suitable for large-scale testing. Mini labs are more suitable in remote and tribal villages as well as for the analysis of highly volatile nitrogen and other major elements. They should complement rather than substitute full-fledged labs.

Under the SHC scheme, the cropped area was divided into grids of 10 ha for rainfed and 2.5 ha for irrigated. One soil sample from each grid will be taken and test results will be distributed to all the farmers whose land falls under the grid. Based on the grid system, out of the total 14.1 crore hectare of net cropped area, 73 lakh grid samples were to be collected to cover 7.3 crore ha area under rainfed situations and 2.7 crore grid samples were to be covered under 6.8 crore ha irrigated land, with a total of 3.46 crore grid samples to be collected in two years, i.e., 1.72 crore grid samples per year. On average, this comes to 25,000 grid samples per district/year or 29 grid samples per village/year. With this, all 14 crore farmers will be covered in two years.

Under cycle 1, 2.54 crore samples were collected, 2.36 crore samples tested, 9.62 crore soil health cards printed, but only 9.33 crore SHCs distributed. This indicates that 100% target was archived in sample collection, 93% of the target was achieved in soil testing, but only 80% of the target was achieved in SHC printing. 97% of the SHCs printed have been distributed among the farmers as on September 24, 2017 (Table 6A). The progress in samples collected, tested, printed, and distributed in a short span is gigantic. However, the gap in the number of farmers covered and the number of SHCs printed and distributed needs to be bridged with proper quality checks at all stages. The progress under cycle 2 is shown in table 6B.

Table 6A: Samples entered, farmers covered, samples tested, and SHCs printed (CycleI) (in Lakhs)

Sl.No.	State	Cumulative Target for Soil Samples Collection & Testing during Cycle-I (2015-16 & 2016-17)	No. of Samples Collected (Cycle - I)	% Progress of Soil Samples Collected (Cycle - I)	No. of Samples Tested (Cycle - I)	% Progress of Soil Samples Tested (Cycle - I)	Cumulative Target for Printing & Distribution of SHCs for Cycle - I (2015-16 & 2016 -17)	No. of SHCs Printed (Cycle - I)	% Progress of SHCs Printed (Cycle - I)	No. of SHCs Distributed (Cycle - I)	% Progress of SHCs Distributed (Cycle - I)
Group - I											
1	Uttar Pradesh	47.7	47.7	100	39.34	82	233.25	106.24	46	106.24	100
2	Maharashtra	23.5	23.5	100	23.47	100	129.77	129.77	100	129.77	100
3	Madhya Pradesh	23.1	23.1	100	23.14	100	88.72	88.72	100	88.72	100
4	Rajasthan	23.1	23.1	100	23.08	100	68.86	69.56	101	68.67	99
Group - II											
1	Karnataka	16.7	16.7	100	16.66	100	78.32	78.32	100	78.32	100
2	Gujarat	15.9	15.9	100	15.89	100	51.09	48.61	95	48.61	100
3	Andhra Pradesh	13.5	13.5	100	13.48	100	74.55	74.55	100	74.55	100
4	Bihar	13.1	13.1	100	10.85	83	72.36	48.02	66	48.02	100
5	West Bengal	13.0	13.0	100	13.00	100	50.41	41.08	81	41.00	100
6	Tamil Nadu	12.7	12.7	100	12.75	100	70.00	70.00	100	70.00	100
7	Telangana	10.3	10.3	100	10.35	100	57.21	57.21	100	57.21	100
Group - III											
1	Punjab	8.4	8.4	100	4.11	49	46.20	8.59	19	8.52	99
2	Haryana	7.9	7.9	100	7.89	100	43.61	28.92	66	16.35	57
3	Chhattisgarh	7.0	7.0	100	7.04	100	38.91	41.14	106	38.91	95
4	Odisha	6.7	6.7	100	6.69	100	36.97	32.16	87	24.03	75
Group - IV											
1	Kerala	1.3	1.3	100	1.28	100	7.05	7.05	100	7.05	100
2	Goa	0.3	0.3	100	0.25	100	0.25	0.20	81	0.20	100
3	Uttarakhand	1.4	1.4	100	1.36	100	7.50	7.50	100	7.50	100
4	HP	0.7	0.7	100	0.70	100	3.85	5.43	141	3.85	71
5	J & K	1.7	1.7	100	1.40	85	9.14	5.11	56	2.50	49
6	Jharkhand	1.2	1.2	100	1.15	100	6.38	6.38	100	5.23	82
7	Arunachal Pradesh	0.2	0.2	100	0.19	93	1.14	0.14	12	0.11	80
8	Assam	2.8	2.8	100	0.85	30	15.41	1.74	11	1.58	91
9	Manipur	0.2	0.2	100	0.06	31	1.15	0.06	5	0.05	90
10	Meghalaya	0.4	0.4	100	0.39	100	2.10	2.10	100	2.10	100
11	Mizoram	0.1	0.1	100	0.10	86	0.12	0.08	69	0.08	100
12	Nagaland	0.3	0.3	100	0.33	100	1.85	1.85	100	1.85	100
13	Sikkim	0.1	0.1	100	0.13	100	0.46	0.46	100	0.46	100
14	Tripura	0.3	0.3	100	0.33	100	1.18	1.18	100	1.18	100
Union Territories											
1	Andaman & Nicobar	0.01	0.01	100	0.01	100	0.08	0.02	27	0.02	90
2	Dadra & Nagar Haveli	0.02	0.02	100	0.02	100	0.12	0.00	0	0.00	
3	Puducherry	0.04	0.04	100	0.04	100	0.20	0.20	100	0.20	100
Total		254	254	100	236	93	1198	962	80	933	97

Table 6B: Samples entered, farmers covered, samples tested, and SHCs printed (Cycle-2) (In Lakhs)

Sl.No.	State	Target for Soil Samples Collection & Testing during Cycle - II (2017 - 18)	No. of Samples Collected (Cycle - II)	% Progress of Soil Samples Collected (Cycle - II)	No. of Samples Tested (Cycle - II)	% Progress of Soil Samples Tested (Cycle - II)	Target for Printing & Distribution of SHCs for Cycle - II (2016 -18)	No. of SHCs Printed (Cycle - II)	% Progress of SHCs Printed (Cycle - II)	No. of SHCs Distributed (Cycle - II)	% Progress of SHCs Distributed (Cycle - I)
Group - I											
1	Uttar Pradesh	23.85	12.67	53	3.44	14	116.63	1.96	2	1.96	2
2	Maharashtra	11.74	14.31	100	6.87	59	64.89	3.57	6	3.57	6
3	Madhya Pradesh	11.57	6.63	57	2.93	25	44.36	8.82	20	8.80	20
4	Rajasthan	11.54	6.40	55	0.00	0	34.43	0.00	0	0.00	0
Group - II											
1	Karnataka	8.33	4.27	51	3.21	39	39.16	518	15	2.32	6
2	Gujarat	7.95	8.14	100	0.18	2	25.54	0.00	0	0.00	0
3	Andhra Pradesh	6.74	2.54	38	2.54	38	37.28	10.12	27	10.12	27
4	Bihar	6.54	2.20	34	0.94	14	36.18	2.12	6	2.12	6
5	West Bengal	6.50	2.99	46	0.90	14	25.20	2.79	11	0.00	0
6	Tamil Nadu	6.37	6.40	100	3.09	48	35.00	0.00	0	0.00	0
7	Telangana	5.17	4.01	78	0.76	15	28.60	0.00	0	0.00	0
Group - III											
1	Punjab	4.18	1.15	27	0.00	0	23.10	0.00	0	0.00	0
2	Haryana	3.94	2.77	70	0.04	1	21.80	0.04	0	0.02	0
3	Chhattisgarh	3.52	3.10	88	1.90	54	19.45	4.56	23	4.08	21
4	Odisha	3.34	1.80	54	0.83	25	18.48	0.91	5	0.76	4
Group - IV											
1	Kerala	0.64	0.00	0	0.00	0	3.53	0.00	0	0.00	0
2	Goa	0.13	0.07	54	0.04	36	0.13	0.04	34	0.04	34
3	Uttarakhand	0.68	0.44	65	0.31	46	3.75	0.69	18	01	14
4	Himachal Pradesh	0.35	0.43	100	0.42	100	1.93	2.42	100	2.42	100
5	J & K	0.83	0.05	6	0.00	1	4.57	0.00	0	0.00	0
6	Jharkhand	0.58	0.47	81	0.34	58	3.19	0.55	17	0.55	17
7	Arunachal Pradesh	0.10	0.00	0	0.00	0	0.57	0.00	0	0.00	0
8	Assam	1.39	0.00	0	0	0	7.70	0.00	0	0.00	0
9	Manipur	0.10	0.00	0	0.00	0	0.57	0.00	0	0.00	0
10	Meghalaya	0.20	0.12	61	0.09	45	1.05	0.30	29	0.25	24
11	Mizoram	0.06	0.00	7	0.00	0	0.06	0.00	0	0.00	0
12	Nagaland	0.17	0.01	4	0.01	3	0.92	0.00	0	0.00	0
13	Sikkim	0.07	0.00	0	000	0	0.23	0.00	0	0.00	0
14	Tripura	0.16	0.07	43	0.05	31	0.59	0.02	3	0.02	3
Union Territories											
1	Andaman & Nicobar	0.01	0.01	81	0.00	47	0.04	0.00	0	0.00	0
2	Dadra & Nagar Haveli	0.01	0.00	0	0.00	0	0.06	0.00	0	0.00	0
3	Puducherry	0.02	0.03	100	0.01	79	0.10	0.00	0	0.00	0
Total		126.8	81.1	64	28.9	23	599.1	44.7	7	37.5	6

5.3 Farmers' Feedback on the Design

The design of SHCs needs to be looked at from the point of view of coverage and intensity of soil testing. Coverage means the scale at which the samples are taken and how representative they are in identifying the variations in soil quality. In the case of intensity, what are the parameters covered while testing the soils? While farmers would benefit more from a comprehensive SHC that is more representative of their plots and covers all the important aspects of soils, it could be time consuming and expensive on the administrative side. More soil indicators mean more sophisticated STLs and better equipment and skilled personnel.

Soil experts and farmers feel that the present scale (2.5 ha grid for irrigated areas and 10 ha grid for the rainfed area) is not representative of all the soil types. Given the high fragmentation in India, each farmer may have a number of fragments, that too not at one place. Farmers want more soil samples to be collected and tested so that they can rely on the recommendations. The number of soil samples per unit area should be based on soil variability. In most of the developed countries, soils are tested as per the requirement by crop.

It has also been observed in the field that the norms prescribed in the SHC scheme are not followed due to various reasons. The timing of soil sampling and the methods of sampling are not adopted in a number of cases. Moreover, the time taken for sample collection and its testing in the lab (although it improved after the SHC scheme) is much longer than usually prescribed. This results in samples losing their characters due to exposure. Hence, the accuracy of the tests and the recommendations prescribed may not be reliable. In some areas, SHCs are not provided to all the farmers. Besides, it has been observed that SHCs are not provided in time. Lack of accuracy, along with the absence of timely provision of recommendations, has affected the credibility of SHCs among farmers. Lack of skilled staff, equipped laboratories, and other infrastructure is the main reason for these shortcomings in the scheme (for details, refer to the appendix on soil health and soil testing and field observations as well as opinion expressed by different stakeholders).

A comparison of indicators included in SHCs in India with other countries such as the USA (across its states) indicates that only nutrient availability, along with some physical soil characters such as PH and EC, are included in India, while a number of states in USA use many other indicators, such as physical indicators, topography, water, soil indicators, biological indicators and nutrient holding capacity (Table 7). Some of these indicators, such as water quality, soil erosion, slope, crop residue, etc., are very important for proper soil management. Experts feel that soil texture, direction of slope, water quality, irrigation source, cracks on soil surface, depth of the soil and crop residues can be included in the Soil Health Card. However, cost may be an issue, but in the case of some aspects such as the irrigation source, depth of soil etc. could be collected from general observation and through the farmer's knowledge about his farm. The importance of including water test in the SHC can be seen in Fig. 5.

Fig. 5: Importance of including water test in the SHC
Salt-affected Irrigation Water

Normal Irrigation Water

Table 7A: Indicators mentioned in SHCs in different states of USA

	Indicators in soil health card		Countries/states/companies													
			India		USA states											
				Connecticut	Georgia	Idaho		Palouse & Nezpere prairies	North - eastern Illinois	Maryland	Montak	Nebraska	Pennsylvania	Northern rivers area	Cornell University's comprehensive assessment of soil health	(private company)
1	Nutrients availability															
A	Macro	N:P:K	Yes	-	-	Yes	Yes	-	-	-	-	-	-	-	Yes	Yes
B	Secondary	Sulphur	Yes	-	-	-	-	-	-	-	-	-	-	-	-	-
C	Micro	Zn, Fe, Cu, Mg, B	Yes	-	-	-	-	-	-	-	-	-	-	-	Yes	Yes
D	Physical characteristics	pH	Yes	-	Yes	-	Yes	-	-	Yes	-	-	-	Yes	Yes	Yes
		Ec	Yes	-	-	-	-	-	-	-	-	-	-	-	-	-
		Soil structure	-	-	Yes	Yes	Yes	Yes	Yes	Yes	Yes	-	Yes	Yes	Yes	-
		Soil slacking	-	-	-	Yes	-	-	-	-	-	-	-	Yes	-	-
		Soil erosion	-	-	Yes	Yes	-	Yes	Yes	Yes	Yes	Yes	Yes	-	-	-
		Organic carbon	Yes	-	Yes	-	-	Yes	-	Yes	-	Yes	Yes	-	Yes	Yes
2	Recommended crop - wise	NPK ratio	Yes	-	-	-	-	-	-	-	-	-	-	-	-	Yes
3	General recommendations	Organic manure	Yes	-	-	-	Yes	-	-	-	-	-	-	-	-	Yes
		Bio - fertiliser	Yes	-	-	-	-	-	-	-	-	-	-	-	-	-
		Lime gypsum	Yes	-	-	-	-	-	-	-	-	-	-	-	-	-
4	Topography	Direction of slope	-	Yes	-	-	-	-	-	-	-	-	-	-	-	-
5	Water	Water quality	-	Yes	-	-	-	-	-	-	-	-	-	-	Yes	-
		Irrigation source	-	Yes	-	-	-	-	-	-	-	-	-	-	-	-
		Water holding capacity	-	Yes	-	-	yes	-	-	Yes	-	Yes	Yes	-	Yes	-
		Infiltration	-	-	-	-	-	Yes	-	-	Yes	Yes	-	Yes	-	-
6	Soil indicators	Cracks on soil surface	-	Yes	-	-	-	-	-	-	-	-	-	-	yes	-
		Depth of soil	-	Yes	-	-	-	-	-	-	-	-	-	-	yes	-
		Depth and colour of topsoil layer	-	Yes	-	-	-	-	-	-	-	-	-	-	-	-
		Moisture of soil two days	-	Yes	-	-	-	-	-	-	-	-	-	-	-	Yes
7	Biological indicators	Biological activity	-	Yes	Yes	Yes	-	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	-
		Crop residue	-	Yes	-	Yes	-	Yes	-	Yes	Yes	Yes	-	-	-	-
8	Nutrient holding capacity		-	-	-	-	-	-	-	Yes	-	-	-	-	-	Yes
	Autoclave - Citrate Extractable ACE) Protein Test		-	-	-	-	-	-	-	-	-	-	-	-	Yes	-
	Cost		free				-								\$110/ sample	\$50/ sample

Table 7B: Indicators mentioned in SHCs in other countries

1	Nutrients Availability		Yes (only N)	Yes (only N)	Yes (only N)	Yes (only N)	Yes	Yes	Yes	Yes	Yes
A	Macro	N:P:K	-	-	-	-	Yes	-	-	Yes	
B	Secondary	Sulphur	Yes	Yes	Yes	Yes	-	-	-	-	
C	Micro	Zn, Fe, Cu, Mg, B					Yes (only B)	-	Yes	-	
			Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
D	Physical characteristics	pH	Yes	Yes	Yes	Yes	Yes	Yes	-	-	Yes
		Ec	-	-	-	-	-	-	-	Yes	
		Soil structure	-	-	-	-	-	-	-	-	
		Soil slacking	-	-	-	-	-	-	-	Yes	
		Soil erosion	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
		Organic carbon	-		-	-	-	-	-	-	
2	Recommended NPK ratio crop wise		Yes	Yes	Yes	Yes	-	-	-	-	
			Yes	Yes	Yes	Yes	Yes	-	-	Yes	
3	General recommendations	Organic manure	-	-	-	-	-	-	-	-	
		Bio fertiliser	-	-	-	-	-	-	-	-	
		Lime/gypsum	-	-	-	-	-	-	-	-	
4	Topography	Direction of slope	-	-	-	-	-	-	-	-	
5	Water	Water quality	-	-	-	-	-	-	-	-	
		Irrigation source	Yes	Yes	Yes	Yes	-	-	-	Yes	
		Water holding capacity					-	-	-	-	
		Infiltration	-	-	-	-	-	-	-	-	
6	Soil indicators	Cracks on soil surface	-	-	-	-	-	-	-	-	
		Depth of soil	-	-	-	-	-	-	-	-	
		Depth and colour of topsoil layer	-	-	-	-	-	-	-	Yes	
		Moisture of soil two days after heavy rain	-	-	-	-	-	-	-	-	
		How moist soil particles hold together	-	-	-	-	-	-	-	-	Yes
		Rooting system	-	-	-	-	-	-	-	-	
		Soil respiration	yes	yes	yes	yes	-	-	-	Yes	
7	Biological indicators	Biological activity	-	-	-	-	-	-	-	-	
		Crop residue	-	-	-	-	-	-	Yes	-	
8	Nutrient holding capacity		-	-	-	-	-	-	-	-	
	Autoclave-Citrate Extractable (ACE) Protein Test						-	-	-	-	
	Cost										

5.4 Designs in Different States

***Bhoochetana* in Karnataka**

The Government of Karnataka initiated a novel mission with the project 'Bhoochetana' from the year 2009-10 with the mission goal of increasing the average productivity of soil by 20%. The consortium partners of the project are Karnataka State Department of Agriculture, Watershed Development Department, UAS, Bengaluru/Dharwad/Raichur, while ICRISAT Hyderabad is the technical consultant.

Main strategies: Soil-test-based nutrient management with a major emphasis on micro nutrients, distribution of inputs at 50% subsidy at the cluster village level, services of farmer facilitators for transfer of technology, farmer field schools, wide publicity through wall writings, posters, village meetings and mass media, effective project monitoring, and feedback.

Under *Bhoochetana*, stratified soil sampling method was used. Districts were divided into three topo-sequences. In each topo-sequence location, samples were taken proportionately from small, medium, and large farmholding farmers' fields to represent different soil colour, texture, cropping system, and agronomic management.

Soil sampling was not done for all the farmers in the village but was collected from only about 20% of the farmers, and the soil sample size was based on the soil variability index with a larger number of samples from higher variability index and fewer number from lower variability index to economise on costs without losing relevance of soil test results for farmers. Test procedures were carried out accurately and in a timely manner. In addition, micro nutrients such as S, Mg, and B were supplied at 50% subsidy, if there was deficiency in soils.

Project implementation started during *Kharif* 2009-10 in six districts, covering 2.25 lakh hectares, 1440 villages, and 2 lakh farmers. The *Rabi* area coverage during 2009-10 was 0.59 lakh Ha. An enhancement in yields of 33-45% was observed in the treated areas. The project was extended to 16 districts during 2010-11, covering 12.0 lakh hectares during the *Kharif* season, including 5,030 villages and 8.50 lakh farmers. The *rabi* area coverage during 2010-11 was 3.32 lakh Ha. An enhancement in yields of 21-41% was observed in the treated areas. During *Kharif* 2011-12, the *Bhoochetana* programme was implemented in all 30 districts, covering 25.4 lakh hectares in 13,800 villages and 20 lakh farmers. The *rabi* area coverage during 2011-12 was 5.40 lakh Ha. An enhancement in yields of 29-41% was observed in the treated areas. The programme was extended to 50 lakh ha of a dryland area and 5 lakh ha of irrigated area during 2012-13. By the fourth year, the project reach was to 26,000 villages covering 42 lakh farmers. There was a considerable area coverage during *Rabi* season as well, i.e., 27 lakh ha during 2012-13. An enhancement in yields of 11-37% was observed in the treated areas.

IFPRI Bihar randomised an experiment of the SHC design

International Food Policy Research Institute (IFPRI) conducted a randomised controlled trial in Bihar to understand the effectiveness of the soil health cards on farmers' incomes on an experimental basis before the SHC scheme of the Government of India. They found no evidence of any impact of soil testing and customized fertiliser recommendations on actual fertiliser use or the willingness to pay for the soil testing services. The study was conducted by the Department of Soil Science of Rajendra Agricultural University (RAU), Bihar, and IFPRI. The three possible explanations for the lack of response are as follows:

- a) Farmers simply did not understand the content of the SHC; we should not expect farmers to change their behaviour on the basis of recommendations that they do not understand.

- b) Farmers understood the contents of the SHC but did not find the soil analysis and fertiliser recommendations to be reliable or compelling.
- c) Farmers did in fact internalise recommendations, and the information did alter their preferred fertiliser mix, but other factors (such as cost, liquidity, or timely availability of specific fertilisers) prevented them from acting on these preferences by shifting their actual application.

5.5 Improvements Required in the SHC Scheme Design

The Scheme

Soil health card is field-specific detailed report of soil fertility status and other important soil parameters that affect crop productivity. Besides soil health, it provides an advisory on soil-test-based use of fertilisers and amendments. As per the design, soil health assessment of farmer fields will be taken up once in two years and soil health cards will be issued. Districts, blocks, and villages within them will be selected in such a way that an action plan is in place to cover them every two years. The nodal soil test lab shall prepare timelines for scheduling the soil health cards in the district in phases. The year-wise coverage of the number of talukas / blocks can be prepared in such a manner that a continuous soil analysis takes place every two years.

Selection Process: The State Governments will prepare yearly action plans indicating the number of districts to be covered, the number of irrigated holdings and rainfed holdings in the selected districts, the number of soil samples to be drawn from irrigated holdings, the number of samples to be drawn from rainfed holdings, and finally, the total number of samples to be drawn and tested. Soil sample collection should be done during the summer months, when there was no standing crop. In most of the cases, this window was open only in April and May, just for 60 days. Hence, all the sample collection should be completed within these two to three months and the sample test results should reach farmers by July. But, generally it takes six to nine months to issue soil health cards. And one season will be over by the time the test results come in. This needs to be rectified and the soil test results should reach farmers before the onset of monsoons so that they can plan the cropping pattern and fertiliser use.

Agencies: a) Agriculture Department staff; b) Science Colleges and students and soil testing/chemistry laboratory staff; c) State Agricultural Universities and their soil testing staff. Soil testing is to be outsourced to private agencies through tenders wherever feasible (based on the technical and financial bids, price quoted for one sample test). Science colleges could be nominated with a provision of providing equipment. Direct aid for setting up of soil testing laboratories is another option. Many farmers expressed faith in private soil testing laboratories as they maintained quality and timely distribution of test results. But only a few farmers are willing to pay for soil tests. Hence, there was a need for promoting the Public-Private Partnership mode of soil testing labs at different locations. In this, investments may be through private sectors with a subsidy component from the government towards payments for testing (per sample basis). The rate per sample is determined in a competitive bidding process from the technically qualified labs that employ qualified and trained chemists. Though this system is currently working, there is a need for increasing competitiveness among private institutions to maintain quality at a reasonable cost. Farmer Producer Organisations need to be encouraged to participate in the bidding process through special incentives.

Sampling norms: The quality of soil testing results and fertiliser recommendations depend upon soil sampling. The following scientific norms are prescribed for this:

- i. In the irrigated areas, samples will be drawn in a grid of 2.5 ha. In rainfed areas, sampling will be drawn in a 10 ha grid.

- ii. In irrigated areas, large, medium, and semi-medium holdings will be sampled and tested holding-wise. In case of marginal and small holdings, sampling in a 2.5 ha grid will be followed. However, field observations indicate that only one sample is taken and tested per grid and the same results are printed on SHCs of all grid farmers.
- iii. In rainfed areas, all the large holdings will be sampled and tested. In case of medium or semi-medium, small and marginal holdings will be sampled and tested in a 10ha grid. Here too, field observations indicate that only one sample per grid is taken and tested. The sample was collected from 3-4 locations in the grid and pooled for testing.
- iv. The ideal time for the collection of soil samples is between sowing/planting of the other crop, i.e., when fields are vacant. This happens mostly in the months of March to June. In many locations, this window was open just for two months, April and May. Often, samples were taken on buds, and only road side plots were covered without following proper sampling framework. However, to meet the target, sample collection should be done in the campaign mode by involving agriculture and allied departments, science collages, and village youth. The agricultural department should monitor the programme closely in order to maintain the quality of samples collected.
- v. The sampling depth for field crops should be 0 to 15 cm.
- vi. Samples have to be collected preferably with the help of stainless steel tube augur, or with a khurpi / spade or kassi.
- vii. A brief training for the soil sampling staff/students/farmers/field extension machinery would be necessary to ensure the collection of representative soil sample and their labelling and transportation to the STLs. Field observations indicated that samples collected were kept idle for months. They need to be properly dried, sieved, packed, and labelled in time to avoid exposure to atmosphere.
- viii. GPS coordinates have to be essentially recorded at the time of soil sampling, which will be downloaded in the STL computer. It is also important to adopt a computerised pre-determined grid system with randomised sampling points, as adopted by Punjab's agricultural department. Punjab's agricultural department distributed tablets with a software application, in which predetermined grids were mapped in GPS-enabled software for the identification of grids in the fields and taking random samples within the grid. This will reduce human error in sampling and collection.
- ix. The target villages will use stratified sampling techniques. Samples will be collected from marginal, small, and large farms to address variations that arise due to different management practices. Within each farm size class, samples will be chosen to represent all possible soil fertility variations.
- x. The focus group discussions and key informant interviews indicated that the uniform grid size does not take into account differences in variation in soil types and cropping pattern. The grid size should be larger if there is less variability in soils and vice versa. This will not only represent soil types based on statistical principles, but it will also reduce costs and workforce. Some studies show that covering 20-30% of the farmers or 20-30 samples per 500 ha is enough for reasonable soil test values if the sample is collected scientifically.
- xi. There is a need for introducing flexible mechanisms for grid sample sizes according to the

variability in the soil fertility index. In this regard, soil fertility variability maps should be developed for each village. Based on this, the grid size should be reduced in low variability villages and increased in high variability villages. In this way, the collected samples will represent the farmer's actual soil conditions with little sampling error, and work burden on the local staff can be reduced to some extent and representativeness of soil samples can be enhanced, which is crucial for increasing the confidence of farmers in the SHC.

Soil analysis: Soil samples should be processed following standard procedures and analysed for various parameters, namely pH; electrical conductivity (EC); Organic Carbon (OC); and available N, P, K, S and micronutrients (Zn, Cu, Fe, Mn & B). However, only a few soil testing laboratories had facilities for micro nutrient tests. In some laboratories, instruments are outdated and take a lot of time to analyse just N, P, and K. There were frequent power cuts, and some were not in a working condition. In some of the laboratories, there was a shortage of chemical reagents to test for basic elements such as NPK. In the recent years, state departments procured mini soil testing kits. In some of the block agricultural offices/labs, five to six mini kits were procured, but agricultural extension officers were too busy with multiple tasks and hardly had any time to test soil samples. It was observed that soil testing through mini kits was time consuming, and the test results were not as accurate as those of full-fledged labs. Hence, in addition to the department soil testing labs, science colleges with soil testing laboratories were involved and assigned the task of soil testing. This was a good decision for timely and accurate dispatch of soil test results to farmers. The students did the work of soil testing under the guidance and supervision of professors. To overcome staff shortage with the state government agricultural department, private agencies were engaged through a competitive bidding process. Field observations indicated that although some private companies maintained quality, some did not. There was a need for proper monitoring and evaluation of the work of private companies. Alternatively, contractual staff can be employed for the testing of soil samples. The soil analysis has to be completed within three weeks of receipt of soil samples in the STL. In the field visits, it was observed that the process of drying, grinding of soil and bottling, etc. was not followed properly by many soil testing labs. There was an inordinate delay in sample testing up to six to nine months. This needed to be rectified to increase the confidence of the farmers in SHC.

There was a significant gap between farmers covered and SHCs distributed. It was observed that awareness camps, campaigns, and Kisan Melas should be the main channel for the distribution of SHCs as that would achieve two goals of handing over the SHCs and explaining how to use SHC simultaneously. It was noted that SHCs were distributed by agricultural officers through democratically elected president or block-level elected representatives in campaigns and through village revenue assistant or by post/online. It was also observed that whenever the SHC was distributed by elected representatives after explaining the content, farmers felt that they understood the utility of the cards and followed recommendations.

Soil Health Card: SHC contains information regarding soil fertility and provides recommendations on fertiliser application for crops and soil amendments required in the case of saline or alkaline soils. Suggestions are made regarding integrated nutrient management. Based on the soil analysis, these are the observations:

- i. Fertiliser recommendations will be developed after considering the available infrastructure/financial resources for the small holders.
- ii. The critical values for delineating deficiency levels will be tested and doses will be recommended for applications.
- iii. Soil-test-based applications for removing deficiencies will be standardised as a component of agronomic practice for the selected crops.

- iv. Timelines will be determined by nutrient status mapping based on soil fertility analysis and productivity enhancement through the application of deficient nutrients. Data will be developed for diagnostic soil analysis and deficient fields in each district. Nutrient recommendations will be prepared separately for Kharif and rabi crops. Block-wise fertiliser dosage adjusted for soil test nutrient status for various crops will be developed.

Cross Checking: Mechanisms are in place for random checking of 1% of the total samples that will be analysed by external agencies in order to ensure the quality of soil analysis. A team of Deputy Director of Agriculture (Ext.), District Agriculture Officer, Assistant Director of Agriculture (STL), and a Soil Scientist from SAU/ ICAR would jointly monitor, inspect, and evaluate the functioning of STLs / other concerned agencies pertaining to the issue of soil health cards to the respective districts. The joint certificate will be issued by the District Agriculture Officer, Sarpanch, and Gram Sevak regarding satisfactory sample collection work. Also, a list of farmers whose soil sample has been collected will be displayed at the Gram Panchayat Office.

At the district level, a state-of-the-art soil testing lab needs to be developed either by a public or private agency, which will be accredited by an international agency. This lab should act as referral lab and develop block-wise soil fertility maps to determine the grid size in each block.

5.6 Socio-economic Characteristics of the Sample Farmers

The basic socio-economic background of the sample farmers is given in Table 8. Overall, the share of small and marginal farmers is 38.2% and SC/ST farmers is 22.1%. The region-wise differences in the share of small and marginal framers is given in the table.

Table 8: Basic Features of the Sample Households (HH)

Region / State	Sample Size (no.)	Coverage of social groups (% SC/ST)	Coverage of Economic Groups			Average Farm Size (acres)
			% of S&MF	% of Medium Farmers	% of Large Farmers	
Central	320	12.7	20.9	43.9	35.2	7.5
East	240	51.4	78.7	21.3	0.0	1.7
North East	448	55.4	47.0	51.0	2.0	3.1
North	864	15.1	25.9	27.1	47.0	8.3
South	800	12.7	28.1	56.7	15.2	4.7
West	512	8.0	51.0	42.5	6.5	3.0
Total	3184	22.1	38.2	41.3	20.4	5.1

Note: S&MF= Small and Marginal farmers (<2.5 acres); Medium Farmers= 2.5 to 7.5 Acres; Large Farmers= > 7.5 acres

Table 9: Basic Features of the Sample Households (HH) by irrigation status

% irrigated group	Coverage of social groups (% SC/ST)	Coverage of Economic Groups			Average Farm Size (acres)
		% of S&MF	% of Medium Farmers	% of Large Farmers	
Less irrigated	27.0	26.8	58.5	28.0	4.6
Medium irrigated	54.4	14.0	66.7	18.7	4.8
Mostly irrigated	17.9	56.8	34.4	42.8	6.0
All	22.1	49.0	41.3	38.2	5.1

Note: Less irrigated (less than 50% of the total area irrigated); medium irrigated (50 to 80%); mostly irrigated (more than 80% of total area is irrigated)

The study grouped farmers based on the share of irrigated area in the total area (Table 9). Farmers who possess less than 50% of area under irrigation were classified as “less irrigated”, farmers with 50% to 80% irrigated area were classified as “medium irrigated”, and farmers with more than 80% irrigated area were classified as “mostly irrigated”. The average area of farmers was higher in “mostly irrigated” than “less irrigated” and “medium irrigated” group of farmers. Table 9 shows that among SC/ST farmers, about 27.0% had irrigation facilities for less than 50% of their total cultivated area, about 54.4% of the farmers had irrigated area between 50% and 80% of their total land, and the remaining 17.9% farmers had land with more than 80% irrigated. The percentage of irrigated area was higher among small and marginal farmers compared to other farmers.

5.7 Awareness, Utilisation, and Outputs under the SHC Scheme

Farm-level impacts are assessed in terms of awareness, participation in the activities, fertiliser use, and impact on crop yields, cost reduction, and profitability. Analysis is carried out across farm size classes and social groups. The analysis is presented by region (zone), development, and time of card issued (old, recent, new, etc).

Awareness

Awareness levels are quite good in South, West, Central, and Eastern zones, with about 80% to 90% awareness. North-East has the lowest awareness at 31.8%, followed by North (78.4%). At the national level, 82.2 % of the sample farmers are aware of SHC (Table 10). The high awareness in some zones could be attributed to the proactive nature of the state governments in the soil health card initiative. Very poor awareness in the North-East may be due to prolonged attachment to traditional ways of farming and less use of fertilisers. It should also be noted that organic farming is practiced on a large scale here. So, the priority for SHC appears to be low at the official as well as farmer level.

Table 10: Zone-wise awareness about SHC

Region/State	% of Sample Farmers			
	Awareness about SHC	Discussion of results	Awareness about portal	Portal info useful
Central	84.9	71.5	47.1	35.1
East	84.4	70.2	33.6	24.2
North-East	31.8	10.9	3.5	2.1
North	78.4	30.4	4.5	1.2
South	89.8	72.8	33.4	23.1
West	88.5	69.4	15.2	12.2
All	82.1	58.6	18.1	16.8

Across the size classes (Table 11), variations in awareness are not much, though higher proportion of medium and large farmers are aware that the SHC is higher among large farmers, reflecting their better access to officials and other community members. Interestingly, awareness about the SHC portal is higher among small and marginal farmers, and it may be the reason that these farmers don't get cards directly from the extension workers and depend on the portal for printing their cards instead. Cards were issued to large farmers at their doorsteps or through campaigns.

Table 11: Farm size class-wise awareness about SHC

Farm size category	% of Sample Farmers are / saying			
	Awareness about SHC	Discussion of results	Awareness about portal	Portal info useful
Small & Marginal	79.4	57.3	21.6	21.4
Medium	81.1	61.5	12.6	11.1
Large	89.1	54.9	21.1	19.8
All	82.1	58.6	18.1	16.8

Fig. 6: Landholding-wise awareness about SHC

The social class-wise analysis indicates that Other Backward Caste (OBC) farmers are more aware of SHC (Table 12). This could be due to the reason that they own poor quality lands when compared to FC farmers. Moreover, of late, these communities are showing more interest in and attachment towards farming (Reddy, 2016).

Table 12: Social group-wise awareness about SHC

Region / State	% of Sample Farmers are / saying			
	Awareness about SHC	Discussion of results	Awareness about portal	Portal info useful
FCs	80.5	56.6	15.7	14.0
BCs	95.2	73.6	19.5	19.2
SC/ STs	67.4	41.9	19.8	19.4
All	82.1	58.6	18.1	16.8

Fig. 7: Method of Soil Sample Collection

Table 13: Awareness about SHC in developed and less developed states

State Group	% of Sample Farmers are / saying			
	Awareness about SHC	Discussion of results	Awareness about portal	Portal info useful
Developed	92.4	74.2	24.2	22.9
Less Developed	55.1	17.2	0.8	0.7
All	82.1	58.6	18.1	16.8

Fig. 8: Awareness about the SHC scheme

The most significant difference in awareness was between developed and less developed states (Table 13) rather than zone-wise. 92.4% of the farmers in developed states are aware about SHC compared to only 55.1% of farmers in less developed states. Awareness about the SHC portal was about 22.9% among developed state farmers, compared to negligible farmers (0.7%) in case of less developed states.

Table 14. Awareness about SHC by irrigation status of the farmers

% Irrigated Group	% of Sample Farmers / saying yes			
	Awareness about SHC	Discussion of results	Awareness about portal	Portal info useful
Less Irrigated	84.6	59.1	4.5	3.4
Medium Irrigated	76.1	49.2	5.8	4.2
Mostly Irrigated	80.1	57.4	23.7	22.4
All	82.1	58.6	18.1	16.8

Awareness about SHC was high among “less irrigated” farmers (84.6 %), but awareness about the SHC portal and the usefulness of the portal was higher among “mostly irrigated” farmers (22.4 %). Again, discussion about the results was higher among “less irrigated” farmers (Table 14).

Fig 9: Soil sampling equipment

♦ **Participation**

Overall, it was observed that 36.3% of the sample farmers participate in meetings, 25.9% of the farmers participate in exposure visits, and 26.3% of the farmers participate in the trainings conducted by agricultural officers. About 25% and 22% of the farmers said that they have benefited from the trainings and exposure visits, respectively. The participation of farmers from the Central and Eastern zones in SHC activities is the highest, whereas North-East and the Northern region were less than 10% (Table 15).

Table 15: Participation of farmers in SHC activities across regions

Region/State	% of Sample Farmers Participating in			% of Farmers Benefiting from		
	Meetings	Exposure visits	Trainings	Meetings	Exposure visits	Trainings
Central	65.9	63.4	67.5	63.4	61.7	61.5
East	63.9	62.0	75.5	61.2	57.3	70.5
North-East	10.3	9.4	5.1	5.8	2.8	2.0
North	16.3	10.2	5.6	5.0	3.4	5.1
South	51.5	27.6	27.5	47.8	26.4	26.7
West	35.1	13.4	3.7	29.9	6.2	0.7
All	36.3	25.9	26.3	33.6	21.8	24.8

Fig 10 : Percentage of Farmers in SHC Activities and Perception

Table 16: Participation of farmers in SHC activities across states

State Group	% of Sample Farmers Participating in			% of Farmers Benefitting from		
	Meetings	Exposure visits	Trainings	Meetings	Exposure visits	Trainings
Developed	46.3	33.4	32.6	43.6	29.4	31.1
Less developed	9.7	6.1	9.8	7.1	1.8	8.2
All	36.3	25.9	26.3	33.6	21.8	24.8

As in the case of awareness, participation in different meetings, exposure visits, and trainings is also significantly higher in developed states (46.3%) compared to less-developed states (9.7%) (Table 16). Significant differences exist even in the percentage of farmers benefiting from exposure visits and trainings.

Although there was no significant difference in the participation of different size class farmers, generally, participation by small and marginal farmers is higher among all the size classes (Table 17). It is interesting to see that small and marginal farmers are more interested and get benefits from government training programmes and exposure visits. Large farmers' participation may be less due to other sources of information.

Table 17: Participation of farmers in SHC activities across size classes

Region / State	% of Sample Farmers Participating in			% of Farmers Benefitting from		
	Meetings	Exposure visits	Trainings	Meetings	Exposure visits	Trainings
Large	33.4	26.1	26.9	30.9	25.7	26.6
Medium	38.7	23.4	23.2	35.6	19.8	21.9
Small & Marginal	35.1	28.4	29.4	32.9	22.0	27.0
All	36.3	25.9	26.3	33.6	21.8	24.8

Participation of farmers from the OBC group is the highest across all activities (Table 18). On the other hand, the percentage of farmers benefiting from these activities was the highest among SC/ST farmers. As far as participation and related benefits are concerned, lower socioeconomic groups (both small and marginal farmers and ST/SC farmers) benefit more when compared to others.

Table 18: Participation of farmers in SHC activities across social groups

Social Group	% of Sample Farmers Participating in				% of Farmers Benefitting from	
	Meetings	Exposure visits	Trainings	Meetings	Exposure visits	Trainings
FCs	30.1	21.9	19.9	28.7	17.1	18.0
OBCs	50.3	32.4	34.1	46.0	29.0	33.5
SC/STs	29.9	25.2	29.3	26.6	22.1	27.1
All	36.3	25.9	26.3	33.6	21.8	24.8

Again, Table 19 indicates that the less irrigated farmers participate more in meetings, while most irrigated farmers participate in exposure visits and trainings when compared to less and medium irrigated farmers. In the case of benefits, a high proportion of most irrigated category farmers get the most in all the three activities. This is in line with our overall observations that the small and marginal farmers and resource-poor farmers are trying to participate in more promotional government programmes, though the benefits are skewed in favour of the resource-rich within the groups. The reason is that their livelihoods are mostly dependent on agriculture compared to large farmers, who can also get income from other sources.

Table 19: Participation of farmers in SHC activities across irrigated area groups

% of Irrigated Group	% of Sample Farmers Participating in			% of Farmers Benefitting from		
	Meetings	Exposure visits	Trainings	Meetings	Exposure visits	Trainings
Less irrigated	38.0	21.4	20.9	33.8	20.5	20.3
Medium irrigated	26.9	21.6	21.1	25.7	14.0	17.0
Mostly irrigated	36.6	27.5	28.3	34.2	22.9	26.7
All	36.3	25.9	26.3	33.6	21.8	24.8

5.8 Farmers' Perception and Source of Information about SHC

Overall, 57% of the sample farmers understand the SHC information. Farmers opine that easy language and colours (green for sufficiency; yellow/brown for deficit; red for severely deficit) should be used to indicate the necessary information to the farmers (Table 20). About 62.8% of the farmers use fertilisers according to the recommendations made by the SHC. This should be improved by imparting awareness about the need for soil management. 38.7% of the sample farmers received assistance from local agricultural extension staff regarding the adoption of recommendations as per the SHC. About 73.1% of the sample farmers said that recommendations are suitable for all plots of the land. About 78% of the sample farmers indicated that the samples represent all the soil types. However, it should be noted that the sample is slightly biased towards better performing districts within states. Proper sampling methods need to be used to increase coverage and relevance of the soil samples collected and tested among all plots. About 68% of the sample farmers said that the results are provided in time, indicating 32% of the sample farmers did not get SHC in time. It seems that there was a delay in sample testing and disbursal of SHCs after sample collection. Agricultural officers should take care that the SHC should reach the farmers in time. It is recommended that the SHCs be distributed during the campaigns and kisan melas through elected representatives in the presence of local agricultural officers. Agricultural officers should display and demonstrate the utility of the content of SHC and show how to adopt the recommended doses of fertilisers as per the SHC.

About 60% of the sample farmers reported that agricultural officers/agricultural extension officers explained the content of the SHC. About 67 % of the sample farmers found the recommendations practical. 46% of sample farmers reported difficulties in adopting practices. Most of the reasons cited are (i) costly to purchase fertilisers, (ii) non-availability of micro nutrients, (iii) perception that traditional high doses of fertilisers will provide more yields than the recommended lower doses, (iv) little difference in existing practice and SHC recommended practice, (v) there was little reduction in fertiliser doses for small farmers and farmers generally prefer more fertilisers as a result, and (vi) agricultural officers are too busy to give proper advice. Agricultural officers are an important link between the farmers and the information they need. At the block level, agricultural officers are burdened with multiple activities. Information dissemination and soil extension has become a low priority compared to distribution of seed and financial assistance under different government programmes. There is a need for pro-active PPP models, under which rural educated youth and entrepreneurs can actively participate in soil sample collection, testing, and dissemination of SHCs for wider adoption of recommended practices by the farmers.

Table 20: Farmers' perception about SHC across regions

Region	% of sample farmers saying yes								
	1	2	3	4	5	6	7	8	9
Central	79.0	77.2	50.4	80.4	85.51	82.6	88.4	77.5	52.2
East	84.4	83.6	30.5	89.1	92.97	89.8	82.0	83.2	50.8
North - East	32.3	21.0	14.9	57.6	76.52	80.3	20.2	39.9	59.6
North	49.0	63.1	0.8	71.3	78.90	39.0	58.8	63.5	36.5
South	71.0	68.3	51.2	73.1	77.21	76.7	78.1	76.4	31.9
West	75.6	90.0	96.3	90.3	92.54	81.5	91.0	77.6	73.6
All	62.8	65.7	38.7	75.3	82.34	71.2	68.2	68.8	47.8

1= Understanding; 2= Using fertilisers and micronutrients according to SHC; 3= Received financial assistance from govt.; 4 = Recommendations suitable to all plots; 5= Do samples represent all the soil types; 6= Results provided in time; 7= Do extension worker explain content; 8= Recommendations practical; 9= Difficulties in adopting practices.

Table 21: Farmers' perception about SHC across size classes

Size class	% of sample farmers saying yes								
	1	2	3	4	5	6	7	8	9
Large	78.7	78.3	27.6	81.0	87.6	78.5	84.0	76.6	42.7
Medium	55.8	56.9	42.8	67.7	72.5	66.4	61.8	58.7	46.0
Small & Marginal	61.9	68.4	40.3	80.6	90.2	72.4	66.7	75.4	52.5
All	62.8	65.7	38.7	75.3	82.3	71.2	68.2	68.8	47.8

1= Understanding; 2= Using fertilisers and micronutrients according to SHC; 3= Received financial assistance from govt.; 4 = Recommendations suitable to all plots; 5= Do samples represent all the soil types; 6= Results provided in time; 7= Do extension worker explain content; 8= Recommendations practical; 9= Difficulties in adopting practices.

Zone-wise farmers' opinion about the implementation of the SHC scheme was shown in Table 20. Overall, the perception index was higher in Eastern India, Western India, and Southern India. It was low in North and North-Eastern India. Hence, there was a need for more intensive efforts in North and North-Eastern zones. In many indicators of adoption of the SHC scheme, small and marginal farmers seem to be better compared to large farmers (Table 21). Farmers in more developed states and districts had better knowledge about the soil health card, and they practised recommended practices (Table 22). In the developed states, farmers face fewer difficulties in adopting recommended practices.

Methods for evaluating soil nutrient status

- **Determination of available nutrients by extraction solution**
- **Determination of exchangeable cations**
- **Soil test correlations with crop response to the applied fertilizer**
- **Rapid soil tests**

Fig. 11: Methods for Evaluating Soil Nutrient Status

Table 22: Farmers opinion about SHC in developed and less developed states

State Group	% of sample farmers saying yes								
	1	2	3	4	5	6	7	8	9
Developed	74.8	76.2	49.8	80.3	84.8	77.5	83.6	76.3	47.0
Less developed	31.1	37.9	9.3	62.2	76.0	54.3	27.4	48.9	49.9
All	62.8	65.7	38.7	75.3	82.3	71.2	68.2	68.8	47.8

1= Understanding; 2= Using fertilisers and micronutrients according to SHC; 3= Financial assistance from govt.; 4 = Recommendations suitable to all plots; 5= Do samples represent all the soil types; 6= Results provided in time; 7= Do extension worker explain content; 8= Recommendations practical; 9= Difficulties in adopting practices.

Table 23: Farmers' opinion about SHC across Irrigation Categories

% irrigated group	% of sample farmers saying yes								
	1	2	3	4	5	6	7	8	9
Less irrigated	57.9	57.4	47.7	76.4	88.6	70.8	76.3	66.0	45.4
Medium irrigated	49.7	37.4	15.2	66.7	76.6	78.4	41.5	54.4	54.4
Mostly irrigated	65.4	70.5	38.4	75.8	81.1	70.6	68.4	70.8	47.9
All	62.8	65.7	38.7	75.3	82.3	71.2	68.2	68.8	47.8

1= Understanding; 2= Using fertilisers and micronutrients according to SHC; 3= Received financial assistance from govt.; 4 = Recommendations suitable to all plots; 5= Do samples represent all the soil types; 6= Results provided in time; 7= Do extension workers explain content; 8= Recommendations practical; 9= Difficulties in adopting practices.

Table 23 reveals that a higher percentage of the “less irrigated” farmers opine that they understand the SHC scheme, using fertilisers and micro nutrients as per the SHC, getting assistance from agricultural officers, recommendations suitable to their soils, sample test values representing their soils, results provided in time, etc. This indicates that the SHC scheme is more pro-poor compared to other government programmes.

It is prescribed that in a grid of 10 hectare of dry land, the soil sample should be collected from the land of all large and medium farmers falling in the grid and also should cover some plots of small farmers through random selection. However, in practice, most of the soil samplers collect sub-samples from 3-4 locations of grid farmers and don't covering small farmers' plots (Table 24). Field surveys show that most of the sub-samples are collected from only one plot of the grid (large) farmers. About 51% of the farmers said that sub-samples were taken from only one plot of the farmers, whereas about 28% of the farmers responded that sub-samples were taken from a few plots of the farmers, and about 18% of the farmers said that the sub-sample was taken from all the plots. However, if the soil is homogenous, representativeness may not be lost with small a number of sub-samples, but if the soil variability is higher, samplers must collect samples from each plot and test them separately. Because of this reason, the study suggests that the grid size should vary with soil variability so that soil test values represent soils correctly without any sampling error. In North-East and South India, there is better coverage compared to other states. A few farmers also reported (2.5%) that they received SHCs without any sample taken from their fields. Many of the field extension officers don't know the exact procedure to be followed in soil sample collection, and they said that they don't see the field manual on soil sample collection. Hence, every block-level office should have the soil sampling manual, and this needs to be properly explained to the soil sample collectors. There is a need for standard protocols for soil sample collection. The soil sample collection of different soil conditions and seasons needs to be standardised. The procedures should vary according to the variability of soil, i.e. the slope of the land, the water sources, etc.

Table 24: Number of plots covered in soil testing across zones

Zone	All (%)	Some (%)	One plot (%)	None (%)
Central	63.4	19.2	12.7	4.7
East	6.6	5.1	84.0	4.3
North-East	18.9	55.6	23.0	2.5
North	1.5	54.3	41.0	3.2
South	42.8	20.3	36.3	0.6
West	25.6	21.6	45.5	7.2
All	25.4	32.2	39.0	3.3

According to farm size groups, 68% among small and marginal farmers indicated that only one plot of their land was covered for soil sample collection, about 19% mentioned that some of their plots were covered, and only 14.5% mentioned that all their plots were covered for soil sampling (Table 25). While 17.5% of large farmers said that all their plots were covered for sampling, 26% said that only one plot was considered for sample collection, and the remaining 55% said that some of their plots were considered for sample collection. Farmers who have land with less irrigated area also mentioned that their soils are covered in soil sampling, which is a good sign. There was no significant difference between developed and less developed states in the coverage of the samples (Tables 25 to 27).

Fig. 12: Number of Plots Covered for Soil Testing

Table 25: Number of plots covered in soil testing across size classes

Size class	All (%)	Some (%)	One	(%)None (%)
Large	18.9	37.1	42.4	1.6
Medium	35.6	21.8	36.6	6.1
Small and marginal	25.1	36.5	35.1	3.3
All	25.4	32.2	39.0	3.3

Table 26: Number of plots Covered in Soil Testing

State group	All (%)	Some (%)	One (%)	None (%)
Developed	29.8	32.5	35.0	2.7
Less developed	13.9	31.5	49.5	5.1
Total	25.4	32.2	39.0	3.3

Fig. 13: Number of plots covered in developed and less developed states

Table 27: Number of plots covered in irrigated groups

% irrigated group	All (%)	Some (%)	One (%)	None (%)
Less irrigated	48.5	24.7	23.2	3.7
Medium irrigated	19.9	44.4	28.1	7.6
Mostly irrigated	19.6	33.2	44.3	2.9
All	25.4	32.2	39.0	3.3

Source of Information

Most of the information was provided by agricultural extension officers and agricultural officers. KVKs and NGOs are active in North-East and Western India. The role of KVKs and NGOs should be highlighted and encouraged by the government for easy dissemination of information (Table 28).

Table 28: Zone-wise sources of information about the SHC content

Region	% of the farmers received information from			
	KVK	AEO/AOs	Scientists of SAU/ICAR	NGOs
Central	39.3	58.8	1.5	0.4
East	5.4	87.4	4.8	2.4
North-East	10.6	77.5	2.8	9.1
North	5.8	89.8	4.2	0.2
South	6.9	85.6	3	4.5
West	1.0	90.1	3.4	5.5
All	10.3	84.6	2.6	2.5

Fig. 14: Main source of information (%)

KVKs and NGOs are active in providing information in less developed states and also cover most of the small and marginal farmers (Table 29) as the extension system in the less developed states are not able to cover all the farmers, given the shortage of skilled workforce and infrastructure (Table 30 & 31). “Less irrigated” farmers mostly got information from AEOs/AOs, while KVKs and NGOs mostly provide information to medium and mostly irrigated farmers (Table 30).

Table 29: Source of information about SHC content across size classes

Size class	% of the farmers received information from			
	KVK	AEO/AOs	Scientists of SAU/ICAR	NGOs
Large	18.4	76.1	3.2	2.3
Medium	12.1	83.1	2.9	1.9
Small & Marginal	5.0	90.2	1.2	3.6
All	10.3	84.6	2.6	2.5

Table 30: Source of information about SHC content by state groups

State group	KVK	AEO/AOs	Scientists of SAU/ICAR	NGOs
Developed	8.8	86.6	3.1	1.5
Less developed	10.9	81.9	2.1	5.1
Total	10.3	84.6	2.6	2.5

Table 31: Source of information about SHC content by level of irrigation

% irrigated group	% of the farmers received information from			
	KVK	AEO/AOs	Scientist of SAU/ICAR	NGO's
Less irrigated	5.3	89.7	1.5	3.5
Medium irrigated	4.5	88.9	2.6	4.0
Mostly irrigated	12.8	80.8	4.1	2.3
All	10.3	84.6	2.6	2.5

Perceptions about SHC benefits

On an average, 70 % of the farmers indicated that they benefited from SHC, while the remaining suggested improvement (Table 32). In most regions, except North-East, farmers feel that the benefits from the SHC scheme are huge in terms of reduction in fertiliser use and cost of cultivation and increased yields. About 45.5% of the farmers are inclined to go to private STLs. About 23.3 % of the sample farmers are willing to pay for the services. However, farmers are willing to pay only in the North and the East as government machinery does not meet the expectations of the farmers in these zones. The zone-wise, farm-size-category-wise, and social-group-wise farmers' perceptions were given in Table 32, Table 33, and Table 34, respectively. It was observed that the farmers in less developed states did not benefit as much as farmers in developed states (Table 35). A majority of the farmers mentioned that they benefited through:

- ✓ Reduction in fertiliser use
- ✓ Identification and application of micro nutrients
- ✓ Increase in yield
- ✓ Knowledge gain

Farmers suggest the following:

- ✓ They should get information in time
- ✓ There is a need for more awareness camps
- ✓ Recommended inputs should be provided free/on a subsidy basis by the government

Table 32: Farmers perceptions about SHC across Zones

Zone	% of Sample Farmers				Type of benefits (by priority)	Nature of improvement
	Benefitting	Need improvement	Willing to go to private STL	Willing to pay		
Central	80.8	73.6	7.6	16.7	Reduce fertiliser use	Timely distribution of SHCs
East	84.8	83.7	47.8	13.6	Reduce fertiliser use	Sample to be taken from all fields
North - East	15.9	13.6	42.9	13.1	Reduce cost	Subsidised micro - nutrient supply
North	63.3	12.5	31.9	22.4	Awareness about soil nutrition	Make farmers aware about SHC
South	86.6	24.7	12.1	10.7	Save fertiliser doses	Information about SHC in time
West	86.8	10.9	43.0	41.3	Increase of production	Recommended practices in more understandable ways
All	73.6	29.1	32.4	19.5		

Table 33: Farmers perceptions about SHC across Size classes

Region/ State	% of sample farmers				Type of benefits (by priority)	Nature of improvement
	Benefitting	Need improvement	Willing to go to private STL	Willing to pay		
Large	82.9	26.4	19.2	17.1	Reduce fertiliser use	Give the SHC reports in time
Medium	73.8	26.7	27.5	19.4	Knowledge about SHC	Free nutrients from govt
Small and marginal	68.4	33.1	44.6	20.9	Get knowledge	Free/subsidized micro nutrients from govt
All	73.6	29.1	32.4	19.5

Fig. 15: Farmers perceptions about SHC across social categories

Table 34: Farmers perceptions about SHC across Social Categories

Region/State	% of sample farmers				Type of benefits (by priority)*	Nature of improvement*
	Benefitting	Need improvement	Willing to go to private STL	Willing to pay		
OCs	78.3	23.5	37.1	21.3	Reduces fertilisers use	Give the SHC reports in time
BCs	82.8	33.5	18.5	19.5	Reduces cost	More trainings needed
SC/STs	50.8	34.9	41.3	15.8	Awareness about soils	Free soil nutrients by govt
All	73.6	29.1	32.4	19.5	Reduces fertiliser use	More trainings needed

Table 35: Farmers perceptions about SHC across states

State Group	% of sample farmers				Type of benefits (by priority)*	Nature of improvement*
	Benefitting	Need improvement	Willing to go to private STL	Willing to pay		
Developed	91.6	32.9	32.2	21.8	Optimal dose of fertilisers	Give information of SHC in time
Less developed	26.0	18.9	32.6	13.6	Awareness about SHC	Free inputs from agricultural dept
All	73.6	29.1	32.4	19.5

Fig. 16: Farmers' perceptions about SHC across states

Table 36 shows that the “mostly irrigated” farmers mentioned that there is a need for improvement in the SHC scheme. About 50% of them are willing to go for private soil testing labs and about 25% of them are willing to pay.

Table 36: Farmers' perceptions about SHC Irrigated Groups

% irrigated groups	% of sample farmers				Type of benefits (by priority)	Nature of improvement
	Benefitting	Need improvement	Go to private STL	Willing to pay		
Less irrigated	78.6	17.8	13.7	16.8	Awareness about SHC	Timely Information about SHC
Medium irrigated	44.4	21.1	39.2	24.0	Save fertiliser doses	Free fertilisers from the dept.
Mostly irrigated	74.9	32.9	36.8	19.9	Reduces fertiliser use	Make farmers aware about SHC
All	73.6	29.1	32.4	19.5	Reduces fertiliser use	Timely Information about SHC

5.9 Outcomes (Input Use and Cost Per Hectare) of the SHC Scheme

The impact of SHC was assessed for three important crops, viz paddy, cotton, and soybean, by comparing fertiliser use and yield before possessing SHC and after following the recommendations of SHC. The study has assessed change in the area under the crops, use of major fertilisers (urea, DAP/SSP and MoP), cost of cultivation, and yields. For these three crops farmers, reported a decline in the area after the SHC scheme, indicating that most of the farmers have diversified to less input-intensive crops from more input-intensive crops, such as paddy and cotton, after the introduction of SHCs. Overall, paddy farmers reduced urea use by about 13%, DAP/SSP by about 12 %, and

Potassium by about 4 %. This is a healthy sign of moving towards a balanced use of fertilisers. All the three crops showed a decline in costs per acre by about 8 to 10%. There was a substantial increase in the use of manures, especially in paddy, which resulted in decreased cost of cultivation per unit area. On the other hand, soybean farmers indicated an increase in fertiliser use, although marginally (Table 37). Crop yields have increased slightly after introducing SHC to all three crops. Among different farm size classes, small and marginal as well as large farmers experienced decline in costs in all the selected crops, while medium farmers reported decline in the case of paddy and soybean crops (Table 38).

Table 37. Impact of SHC on fertiliser use, costs, and crop yields (per ha)

Crop/re gion	S H C	Cost C2	Gross return	Net Return	Yield (qtl)	N (kg)	P (kg)	K (kg)	Fertili ser (kg)	Fert (Rs)	Man ure (qtl)	Manure (Rs.)	Cost A2+FL	Returns over variable cost A2+FL
Cotton														
South	B	66613	80500	13887	17.5	124	64	33	221	6904	9	1004	49136	31364
	A	63283	82800	19517	18	105	60	28	193	5912	12	1256	44106	38694
West	B	73327	96600	23273	21	131	62	22	215	7172	22	2553	52730	43870
	A	70807	101200	30393	22	106	55	20	181	5976	28	3273	47736	53464
Total	B	69970	88550	18580	19.25	127.5	63	27.5	218	7038	15.5	1778.5	50933	37617
	A	67045	92000	24955	20	105.5	57.5	24	187	5944	20	2264.5	45921	46079
% change		-4.2	3.9	34.3	3.9	-17.3	-8.7	-12.7	-14.2	-15.5	29.0	27.3	-9.8	22.5
Paddy														
North	B	54895	81675	26780	45	78	34	18	130	4373	9.5	399	38839	42836
	A	52968	81675	28707	45	68	29	17	114	3786	10.5	473	35103	46572
South	B	73886	92565	18680	51	79	42	22	143	7441	15.5	651	49854	42711
	A	70397	96195	25798	53	67	41	20.5	128.5	6290	19	855	45218	50977
Total	B	64390.5	87120	22730	48	78.5	38	20	136.5	5907	12.5	525	44347	42773
	A	61682.5	88935	27252.5	49	67.5	35	18.75	121.25	5038	14.75	664	40161	48774
% change		-4.2	2.1	19.9	2.1	-14.0	-7.9	-6.3	-11.2	-14.7	18.0	26.4	-9.4	14
Soybean														
South	B	24459	30800	6341	11	31	46	6	83	2074	4	572	21910	8890
	A	23529	30800	10271	11	26	42	6	74	1850	5	762	19500	11300
West	B	35590	42000	6410	15	30	45	6.5	81.5	3597	4	848	28525	13475
	A	33646	44800	11154	16	26.5	38	7	71.5	2976	5	1087	26552	18248
Total	B	30024.5	36400	6375.5	13	30.5	45.5	6.25	82.25	2835.5	4	710	25618	11182
	A	28587.5	37800	10712.5	13.5	26.25	40	6.5	72.75	2413	5	924.5	23026	14774
% change		-4.8	3.8	68.0	3.8	-13.9	-12.1	4.0	-11.6	-14.9	25.0	30.2	-8.7	32.1

Note: B= Before soil health card; A= After soil health card scheme *because of the lack of availability of bio-fertilisers farmers are not able to adopt.

Fig 17: Soil test interpretation categories

Fig.18: Azolla- a potential source of bio-fertiliser

Table 38. Impact of SHC on fertiliser use, costs, and crop yields by farm size category

Crop/Farm size	SH C	Cost C2	Gross return	Net Return	Yield (qtl)	N (kg)	P (kg)	K (kg)	Fertiliser (kg)	Fert. (Rs)	Manure (qtl)	Manure (Rs.)	Cost A2+FL	Returns over variable cost A2+FL
Cotton														
Large	B	53173	87400	34227	19	128.5	66	30	224.5	7144	14	1580	48990	38410
	A	49891	92000	46709	20	108.5	59	26	193.5	6170	16	2045	44150	47850
% change		-6.2	5.3	36.5	5.3	-15.6	-10.6	-13.3	-13.8	-13.6	14.3	29.4	-9.9	15
S&MF	B	63136	87400	24264	18	126.5	60	25	211.5	6934	17	1973	52877	34523
	A	61701	92000	30299	18.5	102.75	56	22	180.75	5725	24.45	2485	47717	44283
% change		-2.3	5.3	24.9	2.8	-18.8	-6.7	-12.0	-14.5	-17.4	43.8	26.0	-9.8	15
Paddy														
Large	B	71075	99825	28750	55	85	43.25	21	149.25	6195	11.75	485	48024	51801
	A	68995	103455	34460	57	69	38	19.5	126.5	5115	13.75	580	44077	59378
% change		-2.9	3.6	19.9	3.6	-18.8	-12.1	-7.1	-15.2	-17.4	17.0	19.6	-8.2	12
S&MF	B	57690	73508	15818	40.5	72	33	19.2	124.2	5617	13	659	40670	32838
	A	54360	73961	19601	40.75	66	31.7	18.2	115.9	4956	15.5	878	36244	37717
% change		-5.8	0.6	23.9	0.6	-8.3	-3.9	-5.2	-6.7	-11.8	19.2	33.2	-10.9	11
Soybean														
Large	B	31082	36400	5318	13	33	47	8	88	2811	4	595	25811	10589
	A	29620	36400	6780	13	28	42	8.7	78.7	2500	5	850	23472	12928
% change		-4.7	0.0	27.5	0.0	-15.2	-10.6	8.7	-10.6	-11.1	25.0	42.9	-9.1	9
S&MF	B	28984	36400	7416	13	28	44	7	79	2524	4.5	790	24625	11775
	A	27560	39200	11640	14	24.55	38	7	69.55	2050	5.65	933	22582	16618
% change		-4.9	7.7	57.0	7.7	-12.3	-13.6	0.0	-12.0	-18.8	25.6	18.1	-8.3	16

Note: B=before SHC scheme; A=after SHC scheme

Table 39: Status according to longevity of card (cotton/ha)

Before SHC scheme	Cost C2	Return over Cost C2	Net Return	Yield (q/ha)	N(kg /ha)	p (kg /ha)	k (kg)	Fertiliser (kg)	Fert (Rs.)	Manure (Qtl)	Manure (Rs.)	Cost A2+FL	Returns over cost A2+FL
Old	70905	87400	16495	19	114	57	21	192	6381	24	2550	49030	38370
New	69793	82800	13007	18	121	54	20	195	6472	22	2447	50083	32717
% change after SHC scheme													
Old	- 5.1	4.5	35	4.3	-20	-10	-15	-17	-18	31	28	11	17
New	- 3.4	3.4	33	3.5	-15	-8	-11	-12	-14	27	27	9.2	14
	- 4.25	3.95	34	3.9	-17.5	-9	-13	-14.5	-16	29	27.5	10.1	15.5
	- 4.2	3.9	34.3	3.9	-17.3	-8.7	-12.7	-14.2	-15.5	29.0	27.3	-10.0	15.7

Table 39 shows that in case of the cotton crop, farmers who possessed SHC for one year reduced fertiliser use significantly compared to farmers who possessed SHC just three months earlier. Yields increased by near about 4% in both old and new cluster groups, while cost of cultivation was significantly lower (3.4%) compared to the farmers who possessed SHC for one year (5.1%).

Table 40: Status according to longevity of card (Paddy/ha)

Percentile Group based on period of holding SHC	Cost C2	Return over Cost C2	Net Return	Yield (q/ha)	N (kg)	p (kg /ha)	k (kg)	Fertiliser (kg)	Fert (Rs.)	Manure re(q)	Manure (Rs.)	Cost A2+FL	Returns over cost A2+FL
Old	54668	68970	14302	38	79	35	22	136	4482	14	1237	39646	29324
New	55223	72600	17377	40	82	65	25	172	4570	13	1250	44595	28005
% change after SHC scheme													
Old	-5.2	2.3	22.9	2.6	-16	-9	-5.5	-12.2	-16	16.5	25	-11	12
New	-3.2	2.1	16.85	1.5	-12	-7	-7.2	-10	-13.5	20	28	-8.5	10.2

In case of Paddy (Table 40), yields have gone up among farmers who possessed SHC for one year (2.6 %). NPK use has been consistently less after the SHC scheme. There has been a consistent decrease in fertiliser use, which is the case across all farmer categories. Net returns of the farmers who possessed the SHC for one year (22.9 %) increased compared to the farmers who received SHC just three months earlier (16.85 %).

Table 41: Status according to longevity of card (Soybean/ha)

Percentile Group based on period of holding SHC	Cost C2	Return over Cost C2	Net Return	Yield (q/ha)	N (kg)	p (kg /ha)	k (kg)	Fertiliser (kg)	Fert (Rs.)	Manure re(q)	Manure (Rs.)	Cost A2+FL	Returns over cost A2+FL
Old	28496	39200	1304	14	20	37	3	60	2687	5	937	23959	15241
New	26760	33600	3840	12	22	35	4	61	2590	4	1,000	23052	10548
% change after SHC scheme													
Old	-5.8	4.5	72	5	-16	-13.8	5	-13	-16	27	33.5	9.8	23
New	-3.7	3.2	64	2.8	-11.5	-10.5	3	-10.3	-13.6	23	28	7.2	20.8

Interestingly, fertiliser use has reduced to nearly 10-13% in soybean after SHC (Table 41). Among the farmers who possessed SHC for the last one year, crop yields increased by 5% with the reduced cost of 6%. NPK use has been consistently less after the SHC scheme was observed for soybean.

Table 42: Status according to longevity of card (Pigeon pea/ha)

Percentile Group of month, Year SHC	Cost C2	Return over Cost C2	Net Return	Yield (q/ha)	N (kg)	p (k)	k (kg)	Fertiliser (kg)	Fert (Rs.)	Manure (q)	Manure (Rs.)	Cost A2+FL	Returns over cost A2+FL
Before SHC scheme													
Old	62437	92398	29960	19	48	41	9	98	4030	2	209	43362	49036
New	61950	91800	29850	18.5	47	40	10	97	4150	2	200	44431	47369
% change after SHC scheme													
Old	-4.5	-9.9	-21.1	6.7	-16.9	-13.9	-10	-18.9	-7	4.9	5.3	112	24
New	-4.5	-9.7	-20.4	5.4	-12.8	-10	-7	-16.7	-7.2	5	7.5	10.5	21

The use of fertilisers has reduced to 19% in case of pigeon pea, which is appreciable. This is a good sign, as farmers generally don't apply fertilisers for pulse crops, as indicated in Table 42. In line with this, the yields have increased slightly among the oldest SHC users (6.7%) although from a very high base for Pigeon pea as the crop is mostly grown under rainfed conditions and farmers give lower priority to these crops. This also highlights that the design of the SHC scheme (10 ha unit) under rainfed conditions needs to be further improved.

Table 43: Status according to longevity of card (Wheat/ha)

Percentile Group of month, Year SHC	Cost C2	Return over Cost C2	Net Return	Yield (q/ha)	N (kg)	p (kg/ ha)	k (kg)	Fertiliser (kg)	Fert (Rs.)	Manure (q)	Manure (Rs.)	Cost A2+FL	Returns over cost A2+FL
Before SHC scheme													
Old	48202	70606	22404	34	106	49	5	160	4716	4	253	33457	37149
New	48525	70514	21989	35	110	48	6	164	4800	4	270	34345	36169
% change after SHC scheme													
Old	-7	4	28	4	-10	-18	-11	-7	-6	11	7	10.5	13
New	-6	3	25	3	-9	-15	-9	-4	-4	8	4	8	10

In the case of wheat, farmers who possessed SHC for the last one year reduced fertiliser use as well as the total cost of cultivation (Table 43). At the same time, their average yield went up by 4%. Urea use reduced by 10%, phosphorous use reduced by 18%, and potassium use reduced by 11%. As a result, the total cost came down by 7%. Although yields increased among other farmers after the SHC scheme, there was no consistency in fertiliser use.

5.10 Success stories and potential benefits of the SHC scheme

More intensive data was collected from 157 farmers across the states during the study (Table 44). This section illustrates the changes observed among these farmers. Out of 157 farmers, 149 farmers reduced nitrogen use and only eight farmers increased after getting the SHC. Average reduction is 30 kg/acre (reduction ranged between 0 and 210 kg per hectare). Out of 157 farmers, 119 farmers decreased phosphorous use and only 38 farmers increased their phosphorus use after SHC. On average, farmers reduced phosphorus use by 11.8 kg/acre. 60 farmers increased their potassium use after SHC, with an average increase of 12 kg/acre. About 50 % of the farmers said SHC made them aware about soil health and helped them reduce the fertiliser use, which ultimately leads to decrease in the cost of cultivation. Out of 157 farmers, 143 farmers experienced increase in productivity after applying the recommended doses as per SHC information. Overall, after getting SHC, farmers have reduced N, P, and K use, especially nitrogen and increased micro nutrients.

Table 44: Success stories of reduction in fertiliser use

Crop cultivated	Before SHC (kg/ha)				After SHC (kg/ha)				% change after SHC				Production Before SHC (kg/ha)	increase in production (kg/ha)	Decreased in Value (Rs/ha)				Increased production value	Total gain (Rs/ha)
	N	P	K	MN	N	P	K	MN	N	P	K	MN			N	P	K	MN		
Cotton	316	119	56	0	237	43	30	0	-25	-64	-46	0	1850	300	632	1371	464	0	10500	12967
Groun dnut	92	106	28	0	32	51	44	0	-65	-52	57	0	1680	466	478	992	-287	0	16310	17493
Maize	283	70	99	0	173	51	55	125	-39	-27	-44	125	2750	863	883	340	784	-1250	9493	10250
Paddy	182	81	53	3	109	59	42	18	-40	-27	-21	18	3280	1141	582	394	200	-150	15974	17000
Paddy, gram	20	14	2	10	12	6	2	10	-40	-57	0	10	3165	927	64	144	0	0	12978	13186
Ragi	109	132	66	0	75	54	35	18	-31	-59	-47	18	1940	1317	270	1402	558	-180	15804	17855
Soybean gram	28	73	16	0	46	111	13	0	64	52	-19	0	1350	630	-143	-683	55	0	15750	14978
Sunflower	119	79	0	0	96	62	0	0	-19	-22		0	2030	1519	181	313	0	0	45570	46064
Total	174	88	49	0	106	59	41	20	-39	-33	-16	20	2268	1010	543	523	141	-198	17797	18807

Estimates from table 44 show that the gains from the practices of the recommended doses of fertilisers as per the SHC show that up to Rs.1,000 can be saved from the reduction in fertiliser expenditure per hectare, and major gain arises from the increase in yields due to adoption of balanced fertilisers to the extent of Rs.17, 000 per hectare.

Fig. 19: Soil-test-crop-response-based fertiliser recommendation system

Figure 20: Soil-test-crop-response-based fertiliser recommendation system

5.11 State-wise and crop-wise fertiliser consumption

State-wise consumption of fertilisers:

As per the Commission for Agricultural Costs and Prices (CACP) data, there was a wide variation in fertiliser application among different states (Table 46). For example, in the case of cotton fertiliser, consumption was the highest in Maharashtra (256 kg/ha) and lowest in Madhya Pradesh (107 kg/ha). Similarly, in case of maize fertiliser, consumption was the highest in Andhra Pradesh (244 kg/ha) and lowest in Chhattisgarh (only 21 kg/ha). In case of paddy, highest fertiliser use was in Karnataka (313 kg/ha) and lowest in Assam (17 kg/ha). These types of anomalies can be eliminated through the adoption of SHCs, and this will also reduce regional yield gaps to a significant extent.

Table 45: Analysis of fertiliser use by crop across states (CACP data) 2013

Crop	State	Yield (qtl/ha)	Fertiliser use (kg/ha)	Gross return (Rs/ha)
Cotton	Maharashtra	19	256	86947
	AP	17	221	70240
	Gujarat	22	216	106209
	Punjab	17	197	94563
	TN	15	180	70542
	Odisha	12	167	47586
	Karnataka	15	158	74865
	Haryana	16	139	85466
	Rajasthan	20	114	105814
	MP	17	107	79010
	Total	19	201	90385
Maize	AP	61	244	76205
	TN	49	212	65552
	Karnataka	40	206	50524
	Gujarat	17	147	33291
	Rajasthan	20	111	31409
	Bihar	23	100	31992
	MP	19	71	24940
	UP	20	70	28512
	HP	14	46	22486
	Chhattisgarh	15	21	18356
	Total	29	130	40194
Paddy	Karnataka	57	313	93307
	TN	51	235	77414
	AP	54	230	79154
	Haryana	48	209	124345
	Punjab	58	199	97222
	Kerala	47	176	97428
	Gujarat	34	173	54250
	UP	39	160	62707
	Uttarakhand	41	157	59953
	WB	42	141	60384
	Chhattisgarh	31	131	42507
	MP	34	114	61441
	Bihar	25	106	32557
	Maharashtra	31	102	53874
	Odisha	28	85	37616
	Jharkhand	19	53	25634
	HP	24	30	43420
	Assam	30	17	34753
	Total	38	138	58543

Sugarcane	Maharashtra	1072	665	244612
	TN	951	415	237486
	Karnataka	848	383	180535
	Haryana	572	263	175556
	UP	497	202	140784
	AP	723	180	168031
	Uttarakhand	463	149	127016
	Total	814	418	199182
Wheat	Punjab	49	242	76828
	Haryana	46	205	82292
	WB	32	191	46693
	Gujarat	32	188	56414
	UP	36	179	62352
	Maharashtra	24	157	45308
	Uttarakhand	28	154	51012
	Bihar	27	144	49135
	Rajasthan	41	127	78468
	Chhattisgarh	16	123	24359
	Jharkhand	23	114	38478
	MP	32	108	54649
	HP	16	57	31029
	Total	34	161	59333

Table 46: Fertiliser use (kg/ha) in 2014-15

State/Zone	2014-15				NPK ratio		
	N	P	K	Total	N	P	K
Pondicherry	196	43	46	285	4.3	0.9	1
AP	147	61	30	237	4.9	2.0	1
Telangana	163	53	16	231	10.0	3.3	1
Karnataka	96	48	32	176	3.0	1.5	1
TN	97	36	31	164	3.2	1.2	1
Kerala	20	8	12	41	1.7	0.7	1
A&N Islands	13	10	7	29	1.9	1.5	1
South Zone	106	43	26	175	4.1	1.7	1
Gujarat	96	28	9	133	10.6	3.1	1
Maharashtra	69	34	23	126	3.0	1.5	1
Chhattisgarh	57	27	9	92	6.6	3.1	1
MP	49	26	3	78	15.5	8.4	1
Rajasthan	40	14	1	55	62.5	21.7	1
West Zone	59	25	9	93	6.7	2.9	1
Punjab	179	43	5	227	36.0	8.7	1
Haryana	172	43	6	221	28.0	7.0	1
Uttarakhand	129	24	7	160	18.7	3.5	1
UP	112	32	7	150	16.9	4.9	1
HP	36	9	10	54	3.7	0.9	1
North Zone	131	36	7	174	19.7	5.3	1
Bihar	133	31	15	179	9.0	2.1	1
WB	78	40	33	151	2.4	1.2	1
Odisha	36	15	7	57	5.3	2.1	1
Jharkhand	31	7	1	38	28.2	6.0	1
East Zone	75	27	17	118	4.4	1.6	1
Assam	68	22	35	125	1.9	0.6	1
Manipur	45	9	8	62	5.7	1.1	1
Tripura	18	14	8	41	2.2	1.7	1
Nagaland	3	2	1	6	2.8	1.6	1
Meghalaya							
Arunachal Pradesh							
Mizoram							
Sikkim							
NE zone	45	15	22	83	2.0	0.7	1
All-India	85	31	13	128	6.7	2.4	1

Source: Fertiliser Statistics

The state-wise fertiliser use is given in table 46. It shows that fertiliser use per hectare was the highest in Puducherry (285 kg/ha), followed by Andhra Pradesh (237 kg/ha), Telangana (231 kg/ha), Karnataka (176 kg/ha), and Tamil Nadu (164 kg/ha). The NPK ratio was close to the recommended practice for states such as Pondicherry, Andhra Pradesh, Karnataka, Tamil Nadu, Himachal Pradesh, and Maharashtra. In general, NPK ratio for southern and eastern zones is closer to the recommendation. But in case of northern states, Jharkhand, Rajasthan, Punjab, Haryana, Uttarakhand, and Uttar Pradesh are much different from the recommended doses, and this needs to be changed. SHC-based recommendations will help in reducing these imbalances in fertiliser application, which will help soil health in turn.

Chapter VI

Policy Recommendations

Soil health and fertility form the basis for sustainable profitability of the farmers. Using optimal doses of fertilisers and cropping pattern as per the scientific recommendation are the first step towards sustainable farming. Soil testing is a science-based and time-tested tool for assessment of soil fertility status and soil ailments and for nutrient amendment recommendations. Soil testing, as a tool for judicious fertiliser use, works on the principle of profitability, which means if all other factors of production are optimum and none of them are limiting, there is a probability of obtaining a more profitable response to applied nutrients based on soil testing than those applied on an ad hoc basis.

In India, the current consumption of NPK ratio is 6.7:2.4:1, which is highly skewed towards nitrogen as against the ideal ratio of 4:2:1. India spends nearly Rs. 70,000 crore on fertiliser subsidy every year. According to the estimates, subsidy amount is about Rs. 5,000/ha of net cropped area and about Rs. 5,100/farmer, resulting in excessive use of fertilisers, especially NPK, at the cost of micro nutrients and manure. Hence, there is a need for balanced use of fertilisers. Keeping this in mind, the Government of India introduced Soil Health Card Scheme across India (GoI, 2017).

On December 5, 2015, the Ministry of Agriculture introduced the Soil Health Card scheme. It was approved for implementation in the remaining period of 12th plan. SHC will be provided to all farmers in the country at an interval of two years to enable the farmers to apply recommended doses of nutrients based on soil test values for improved and sustainable soil health and fertility, low costs, and higher profits.

Under the SHC scheme, cropped area was divided into grids of 10 ha for rainfed and 2.5 ha for irrigated. One soil sample from each grid was to be taken, and the test results were to be distributed to all the farmers whose lands fell under the grid. Based on the grid system, of the total 14.1 crore hectare of net cropped area, 73 lakh grid samples were to be collected to cover 7.3 crore ha in rainfed areas and 2.7 crore grid samples were to be collected to cover 6.8 crore ha irrigated land. That is, a total of 3.46 crore grid samples in two years (1.73 crore grid samples per year) and an average of 25,000 grid samples per district/year or 29 grid samples per village/year. With this, all 10.39 crore farmers will be covered in two years. Every year, 5.2 crore farmers need to be covered.

Under cycle 1, 2.54 crore samples were collected, 2.36 crore samples tested, 9.62 crore soil health cards printed, but only 9.33 crore SHCs distributed. This indicates that 100% target was archived in sample collection, 93% of the target was achieved in soil testing, but only 80% of the target was achieved in SHC printing. 97% of the SHCs printed were distributed among the farmers as on September 24, 2017. Cycle 2 has already started in many states.

Objectives of the Impact Study

As the SHC scheme has completed more than 2 years of implementation, the ministry has initiated a nationwide impact assessment with the following objectives:

- o To examine the design of the SHC scheme in terms of planning, implementation, inputs (staff, financial, and other resources), activities (trainings, lab established, and strengthened), outputs (SHCs printed and distributed to farmers).
- o To assess the modalities of delivery of the SHC scheme regarding procurement, sample collection, testing, SHC printing, and disbursal.
- o To assess the level of utilisation of SHCs by the farmers across farm size classes in irrigated and rainfed situations.
- o To assess the impacts of the SHC scheme on judicious use of fertilisers, bio-fertilisers, organic fertilisers, soil health, cropping choice, cost reduction, farm profitability, and sustainability.
- o To provide recommendations for improvement of the overall design of the programme.

❖ Methodology

Both quantitative and qualitative approaches were adopted to achieve the objectives of the study. Qualitative information was in the form of stakeholder interviews across the states under the study, expert opinion gathering at the national and state level workshops, and interactions with the progressive farmers and agricultural officers. At the quantitative level, both secondary and primary data was collected at the national, state, and farmer levels. Secondary data mostly pertains collection and analysis of financial and physical achievements of the SHC scheme over the years, infrastructure availability, coverage of SHCs across the states, etc. Besides, information at the international level was collected for some selected countries to see the best practices in the design of soil health cards.

The secondary data was analysed for all the states, while primary data was analysed for 16 states of India representing all agro-climatic zones. A systematic sample was drawn for the impact assessment at the farmer level. Care was taken to represent the whole country and its agro-climatic conditions. A structured questionnaire was canvassed among 3,184 sample farmers across 199 villages in 16 states. In addition, focus group discussions were conducted in each village to get feedback from the key informants, farmers not covered for individual surveys and farmers who had not received soil health card. All indicators collected from the field survey were classified as inputs (financial and physical inputs under the project), activities (different activities organised under the scheme), outputs (actual outputs of the project), outcomes (whether generated outputs were utilised by the farmers), and impacts (what are the ultimate benefits to the farmers); they're listed here. The analyses were carried out across zones/states and completed by the date of receiving SHC by the farmers, i.e., those who received SHC more than a year back and those who received it recently. This would help understand the long-term impacts as well as provide insights into whether agriculture development helps better awareness and demand for SHC.

Cost Concepts Used in Calculating the Net Returns and Returns over Variable Costs

1. Costs are generated through certain cost concepts.

These cost concepts and the items of costs included under each concept are as follows:

Cost A1 (all paid out costs incurred by owner-cultivator):

- i. Value of hired human labour
- ii. Value of hired bullock labour
- iii. Value of owned bullock labour
- iv. Value of hired machinery labour
- v. Hired machinery charges
- vi. Value of seed (both farm produced and purchased)
- vii. Value of insecticides and pesticides
- viii. Value of manure (owned and purchased)
- ix. Value of fertiliser
- x. Depreciation on implements and farm buildings
- xi. Irrigation charges
- xii. Land revenue, cesses, and other taxes
- xiii. Interest on working capital
- xiv. Miscellaneous expenses (artisans etc.)

Cost A2: Cost A1 + rent paid for leased-in land

Cost B1: Cost A1 + interest on value of owned fixed capital assets(excluding land).

Cost B2: Cost B1 + rental value of owned land (net of land revenue) and rent paid for leased-in land

Cost C1: Cost B1 + imputed value of family labour

Cost C2: Cost B2 + imputed value of family labour

Cost C2*: Cost C2 adjusted to take into account valuation of human labour at market rate or statutory minimum wage rate, whichever is higher.

Cost C3: Cost C2* + value of management input at 10% of total cost (C2*).

In this study, only cost A1 and cost C2 were used to calculate returns over variable costs and net returns, respectively, by deducting costs from gross returns.

2. Imputation Methods

Some of the inputs used in the production process are provided by family sources. The criteria adopted for deriving imputed values of these inputs are as follows:

Sl. No.	Items	Criteria
(1)	(2)	(3)
i.	Family Labour	On the basis of statutory wage rate or the actual market rate, whichever is higher
ii.	Owned animal labour	On the basis of cost of maintenance, which includes cost of green and dry fodder and concentrates, depreciation on animal and cattle shed, upkeep labour charges and other expenses.
iii.	Owned machinery charges	On the basis of cost of maintenance of farm machinery, which includes diesel, electricity, lubricants, depreciation, repairs and other maintenance expenses.
iv.	Implements	Depreciation and charges on account of minor repairs.
v.	Farm produced manure	Evaluated at rates prevailing in the village.
vi.	Rent of owned land	Estimated on the basis of prevailing rents in the village for identical type of land or as reported by the sample farmers subject to the ceiling of fair rents given in the land legislation of the concerned state.
vii.	Interest on onward fixed capital	Interest on present value of fixed assets charged at the rate of 10% per annum.

3. Allocation/Apportion of Joint Costs:

The expenditure incurred on, or imputed for, some of the cost items relate to the farm as a whole. Such joint costs are allocated to individual enterprises, among different categories of livestock, and so on. Depreciation on farm buildings and implements, land rents, land revenue, cesses and taxes, interest on owned fixed capital are such costs that are allocated to each category of crops in proportion to their areas. The cost for livestock is allocated to each category of animals in proportion of their numbers to the total number of animals owned by the farmer.

The apportionment of total costs incurred jointly on different crops grown in mixture crops is done in proportion to the total value of output contributed by individual crops in the crop mixtures. The apportionment of total costs of cultivation between the main product and the by product(s) is done in proportion to their contribution to the total value of output.

❖ Results

It should be noted that the analysis is based on the representative, though limited, sample size across regions. It is too short a time for the scheme (only two-years old) to carry out a full-fledged impact assessment. The present analysis provides insights about the direction and cautions about any shortcomings. While the following conclusions and recommendations are based on the analysis, the weakness of the assessment needs to be kept in mind.

- ✓ Given the short duration of the scheme, awareness levels are good. At the same time, participation of farmers in meetings and exposure visits is not high. Awareness campaigns need to be organised on the content of SHCs, use of recommended practices, reduction in fertiliser use, and costs and increase in profitability.
- ✓ There is no apparent or significant bias against socio-economically vulnerable sections. In contrast, small and marginal farmers benefit more in some cases.
- ✓ There is some reduction in fertiliser use, especially nitrogen and increase in bio-fertilisers and other micro nutrients use. This is a good sign as the N: P: K ratio was highly skewed towards nitrogen. Costs were reduced due to low fertiliser use. Crop yields also increased for a majority of the crops, although only moderately.
- ✓ A significant impact is the increase in the use of gypsum and other micro nutrients to some extent.
- ✓ There is a need to strengthen the SHC-related extension services to provide better advisories.
- ✓ Two-thirds of the sample farmers indicated that SHC is beneficial, which is encouraging given the short span of the programme.
- ✓ The main complaint from the farmers is the timeliness of providing the results. This, however, is linked to the infrastructure (soil testing labs) and human resources. However, after the introduction of the SHC scheme, the time lag has significantly reduced. Results need to be disseminated before the sowing season so that farmers will practice the recommended crop choice and fertilisers.
- ✓ It is important to address these issues to gain confidence of the farmers in adoption of the fertilisers as per the recommendations included in the SHC.
- ✓ The scheme has poor backing from infrastructure and human resources, with significant gaps. Although some southern and western states performed better, in some states, even the allocated resources are not being spent or utilised due to lack of capacities. This should be of high priority in the immediate future.
- ✓ Proactive regions seem to be better in this regard.
- ✓ Lack of capacities with regard to skilled personnel and STLs affects the quality of services, which in turn affects the credibility of the scheme and needs immediate attention.
- ✓ Results need to be provided in time so that farmers can benefit better.

6.1 Soil Sampling (SHC Design)

- There is a need to identify best practices in soil sample collection and testing by examining across countries and different state governments practices. There is also a need for coordination and cooperation.

- The existing uniform grid of 10 ha for dry lands and 2.5 ha for irrigated lands does not take local soil variability into consideration. Grid size should be variable based on the soil variability index. It should be decided at least at the block level based on soil heterogeneity, fertility maps, cropping pattern, irrigation facilities, and remote sensing maps. If soil is more variable, grid size should be reduced, and vice versa. Sampling errors need to be reduced by using variogram. There should be a separate cell to monitor and recommend grid size across the country. This will also reduce cost, money, and manpower and increase the relevance of recommendations to farmers. In order to gain credibility of the farmers, at least one sample from each farmer should be included where soil variability is high.
- Soil variograms need to be developed at each block level. Grid size can be determined based on the soil variability. Variogram gives information about the spatial pattern of continuous soil attributes. The variogram may be used as a critical input to decide required soil samples to be collected based on the soil variability index. More soil samples should be collected if the block-level soil variability index was high and vice versa. Variogram is a tool to investigate and quantify the spatial variability of soil properties. The geostatistical literature shows that the following soil quality indicators were found to be the most important for variogram analysis, which can also be considered by the ministry. The available data with Indian Institute of Soil Science, Satellite maps of remote sensing agency of ISRO, and Land use planning data can be used to estimate block-level soil variograms.
- The following soil indicators may be considered to construct block-level soil variograms: (1) Soil Colour, (2) Slope, (3) Sand (%), (4) Silt (%), (5) Clay (%), (6) Nitrogen (N), (7) Phosphorus (P), (8) Potassium (K), (9) Organic Matter (OM), (10) Organic Carbon (OC), (11) pH, (12) Cation Exchange Capacity (CEC), (13) Electrical conductivity (EC), (14) C:N Ratio, (15) Cropping Pattern. The ICAR-Indian Institute of Soil Science (IISC), Bhopal; ICAR-IISWC-Indian Institute of Soil and Water Conservation, National Remote Sensing Agency (NRSA), Hyderabad; and land classification/atlas can be consulted for developing block-level soil variograms.
- Evidence shows that samples collected from 20% to 30% of the farmers in a village are enough to get reasonable soil quality for advising farmers, and hence there was no additional benefit in covering each grid of 10 ha in case of dry lands and 2.5 ha in case of irrigated land if the soil was fairly uniform. In some cases, only 20-30 samples/500 ha are sufficient, as is evident from ICRISAT experiments. This, however, needs to be explained to the farmers so that they would take the SHC recommendations seriously. One needs to be mindful of the efforts and resources gone into ICRISAT experiments.
- Although grids in some states were predetermined in the mobile app (such as in Punjab), in other cases, the procedure that's followed in dividing the cultivated village land into grids is not known to many agricultural officers and needs to be widely disseminated for accurate sampling, and this should be mentioned in the guidelines. Interestingly, Punjab is adopting a GIS-tablet grid identification and GPS-based soil sample collection application, which seems to be working well and is likely to solve many field-level sample collection problems. This model should be adopted across all the states after an in-depth understanding (study) of the model.

- High density soil maps need to be developed for increasing precision at the village level.
- Agricultural officers and agricultural extension officers need to be given appropriate training, transport, and easy-to-use sampling tools and incentives for scientific sample collection.
- Coordination of agricultural extension officers and farmers needs to be enhanced, and extension officers should make sure that most of the grid farmers, if not all, should be present at the time of soil sample collection. This will build farmers' confidence in the soil health cards.
- It was observed by the study team that in some of the block agricultural offices, soil samples were kept aside for many days and soils were exposed to moisture and weather. After soil sampling, drying should be done within 15-20 days, and grinding, machine sieving, and bottling should be done in time for proper test results. Sample test results should reach farmers before the sowing season. It might be a good idea to limit samples to the capacities. The target should be to provide more accurate results rather than coverage. This would create demand for soil testing once the credibility of the testing is established.

6.2 Soil Health Indicators (SHC Design)

- The whole chain of soil health-plant-health-human health should be taken into account, and there is a need for promotion of balanced application of soil (macro and micro) nutrients.
- Excess application of urea results in accumulation of nitrate in the soil, and water is becoming a huge environmental problem in India. Hence, water quality information needs to be included in the SHC.
- The soil health card is more focused on chemical nutrient indicators; among physical and biological properties, only soil colour is included. Some more physical properties, such as slope of the land etc. needs to be incorporated as well.
- Microbial activity and moisture retention activity are essential but missing in SHC. Although soil organic matter is indicated, many soil testing labs are not equipped with the latest tools to measure it.
- At least one or two physical and micro-biological indicators (such as soil texture, water holding capacity, water quality and bacterial content) need to be incorporated. Soil health index needs to be developed and incorporated into SHC, which indicates the overall health of the soil.
- Although the basic structure of SHC should be uniform, states should adopt/change as per their agro-climatic zones and needs. Some of the indicators that need to be included in SHC are (i) cropping history, (ii) water resources (soil moisture), (iii) slope of soil, (iv) depth of soil, (v) colour of soil, (vi) soil texture (bulk density), and (vii) micro-biological activity.

6.3 Soil Testing Infrastructure

- About 1,454 labs exist in India, of which only 700 are equipped with micro nutrient testing facilities. Although agricultural departments procured about 7,000 mobile kits recently, they are not as good as full-fledged labs. Very few labs could take up micro nutrient analysis. They are neither equipped with skilled personnel nor chemicals or functional equipment. The infrastructure is grossly inadequate by any standard, given that 11 crore farmers need to be covered.
- Under the current PPP model, investments in labs are to be done by private companies with an element of subsidy. A competitive bidding process based on technical and financial bids needs to be announced, and companies that quote reasonable cost (per sample) should be selected. The government will pay on a per-sample basis with the condition that they employ qualified and trained chemists. This model will be successful when there is no strong government presence in soil testing. However, the quality of such reports should be checked at random by authentic agencies.
- About 45% of the sample farmers are inclined to go to private STLs. At the same time, only 20 % of the farmers are willing to pay for the services. Hence, one must find ways to support farmers in this regard, i.e., direct subsidy to the farmers or private STLs, etc. A competitive PPP model could be explored in this regard, while the government should take up the monitoring of the functioning of these labs more seriously. At the same time, there should be special incentive for Farmer Produce Companies (FPCs) to establish soil testing labs. There is a need for encouraging competition among private companies in setting up and running the soil testing labs so that they maintain quality at a reasonable cost. Institutional modalities could be worked out regarding how to equip and manage STL within the FPCs framework.
- Some of the private soil testing labs indicated that the cost of sample collection and testing was up to Rs. 1,000/sample. Some private companies are charging Rs. 75/element; accordingly, for 12 elements, the total cost will be Rs. 900/sample. The government should be more realistic in fixing the prices for private parties. It should focus more on the quality of services at an acceptable (market) cost.
- There should be strengthening and upgrading of at least one soil testing lab per district as a state-of-the-art lab, equipped with world class infrastructure and accredited by internationally recognised agencies either in the public or the private sector. So, nearly 700 state-of-the-art labs are needed to act as referral labs and to give broad advice to farmers. The cost per unit will be about Rs.4-5 crore/unit, with a total of Rs. 2,800 crore. However, if this resulted in Rs. 1,000/ha savings in fertiliser use even if we don't consider the yield increase resulting in a saving of Rs. 14,500 crore in a year for the economy. This could be a more worthwhile investment than spending money and providing services that have little value to farmers year after year, i.e., spending more in the long run.
- State-of-the-art district-level soil testing labs at a direct level should be equipped with Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES), which costs about Rs.

40-50 lakh. These labs should have a 24-hour generator for uninterrupted power supply, computer labs with colour printing facility, and air conditioned laboratories. In addition, the lab should have the following equipment for conducting soil testing on a large scale:

- o KEL PLUS automatic nitrogen determination distiller
 - o Automated Flame Photometer (for Potassium)
 - o Automated Spectrophotometer (for Phosphorus)
 - o Atomic Absorption Spectrophotometer (for Zn, Fe, Mn, Cu)
 - o Water distillation still (20 lit/hour)
 - o All glass distillation unit (5 litre/hour)
 - o Auto analyser (N&P)
 - o Automated pH meter
 - o Automated EC meter
 - o Centrifuge
- Some soil scientists and agricultural officers are of the opinion that test results of mini kits (mini labs) are not as accurate enough as those of full-fledged labs. Mini kits need to be standardised and tested for errors in calibrations. A mini kit costs about Rs. 94,800. With this, the per-sample cost comes about Rs. 170-200. A mini kit is useful for remote villages and tribal communities and should also be used to measure highly volatile elements such as Nitrogen, which needs more frequent measurement. Some block agricultural offices received five-six mini kits, but they were not able to utilise them as they are involved in multiple activities.
 - Soil testing is a specialised and highly skill-oriented job. Frequent transfers of soil testing staff adversely affect the skill development within labs as well as the test results. There is a need to build some permanent staff in the labs who are interested and specialised in soil testing. Field observations indicate that only female officers are interested in working with soil testing labs.
 - Management of state-of-the-art soil testing labs could be established under the purview of FPC's federations or nodal FPCs at the district level. The governance responsibilities should be handed over to them to be run them as business models. The back of the envelop indicates that establishing a state-of-the-art lab with a Rs. 6 crore loan from NABARD in each district looks viable given that an average of 25,000 samples need to be collected and processed per year. At Rs. 1,500-2,000 per sample (at half the market price for a detailed soil analysis), the investments will be paid back in less than 2 years. The state department can have a monitoring cell created especially for this purpose, and extension services need to gear up to deal with soil health advisories. FPCs should be encouraged to set up demonstration plots.

6.4 Soil Extension

- SHC recommendations should be accompanied by block-level recommendations. There should be an intermediate solution (based on both village-level soil maps and SHC recommendations) to be able to reach the farmer's level.

- There is a need for demonstration of SHC benefits on an experimental basis in each block by adopting a comprehensive approach (systematic and scientific analysis of soil and water) and recommended doses. This would have much greater impact than subsidised and less authentic information. General SHG schemes and model farm initiatives should go together.
- In many villages, agricultural officers are distributing SHCs in awareness campaigns through village presidents and mandals/blocks' democratically elected representatives. However, in some villages, the village revenue assistant distributes SHC and takes signatures without explaining the content. Whenever SHC is distributed in awareness campaigns and meetings directly, a greater number of farmers feel that they are convinced to use recommended practices. There is a need to follow standard protocol to inform farmers about the recommendations of the SHC when it is handed over to farmers.
- A specialised body is needed both at the central as well as the state level for the management of soils. They should be given the responsibility of monitoring the quality of service by various agencies. This also provides continuity in the workings of the department.
- Development of GIS-based soil fertility maps at the village/block level and wider publicity through wall posters and display boards in village panchayats should be promoted. Advertisements, slogans, etc. should be developed in local languages to increase awareness. This should be taken up in a campaign mode, i.e., on the lines of the anti-smoking/tobacco campaign.
- Many farmers are not aware of the SHC portal. The SHC portal should be more farmer-friendly and simplified. A professional body may be employed to design the portal in more farmer-friendly and effective manner.
- A simple tool to assess the quantity of urea, DAP, and MoP based on SHC needs to be displayed as wall posters in the villages.
- It should be mandatory to enter fertiliser purchases by the farmers on the soil health card by each fertiliser dealer along with the signature. This will not only increase awareness, but will also help in adoption as per the recommendation.

6.5 Policy-related

- If the SHC programme is to be successful, the high fertiliser subsidy for NPK needs to be reduced. Prices should reflect true cost to the economy; only then will farmers have incentives to use fertilisers judiciously according to the recommendations of the SHC. On the contrary, subsidy on micro nutrients should be increased. However, the quality of micro nutrients supplied on a subsidy basis in some states is highly doubtful.
- The government should set up state-of-the-art labs to test the quality of micro nutrients supplied. Accreditation of such labs to national/international standard institutes should be initiated as well. Supply of phosphorous soluble bacteria should be mandatory along with phosphate fertilisers and

rock phosphate-like neem-coated urea. FPCs need to be encouraged to take up the SHC scheme as a business model by setting up state-of-the-art labs at the district level. Since FPCs are already involved in selling fertilisers, they are in a better position to stock all the required (according to SHC recommendations) fertiliser and micro-nutrient compositions and supply them to individual FPCs across the district.

- State-of-the-art district-level soil testing labs should be equipped with Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES), which costs about Rs. 40-50 lakh. These labs should have a 24-hour generator for uninterrupted power supply, computer labs with colour printing facility, and air conditioned laboratories. In addition, the lab should have the following equipment for conducting soil testing on a large scale:
 - o KEL PLUS automatic nitrogen determination distiller
 - o Automated Flame Photometer (for Potassium)
 - o Automated Spectrophotometer (for phosphorus)
 - o Atomic Absorption Spectrophotometer (for Zn, Fe, Mn, Cu)
 - o Water distillation still (20 lit/hour)
 - o All glass distillation unit (5 litre/hour)
 - o Auto-analyser (N&P)
 - o Automated Ph meter
 - o Automated EC meter
 - o Centrifuge
- There should be some incentives/awards for the farmers who grow green manure and vermi-compost and whose soil fertility increased over the years based on Soil Health Card.
- Some incentives should be provided to local bodies that encourage good practices such as recycling crop residues, encouraging common lands for corporates, etc.
- Similarly, incentives can be given to villages when they adopt crop rotation with legumes.
- Innovative techniques such as neem-coated urea (for slow release of fertiliser into the soil) need to be promoted by the government. There should be 45 kg urea bags instead of 50 kg ones, which will reduce the loss/excessive use of fertilisers by about 10-20%, especially by small and marginal farmers.
- Soil sample collection, testing, and printing at the district level is significantly positively influenced by fertiliser use, number of bank accounts, net sown area, number of soil testing labs, and households with mobiles.
- Other policies, such as water exploitation, electricity, etc., should be in line so that crop diversity that can protect soil health in the long run.

References

1. GoI (2016); *Desertification and Land Degradation Atlas of India (Based on IRS AWiFS data of 2011-13 and 2003-05)*; Space Applications Centre Indian Space Research Organisation Department of Space, Government of India, Ahmedabad, June.
http://www.indiaenvironmentportal.org.in/files/file/Desertification_Atlas_2016.pdf
2. Reddy, B. Suresh. "Dynamics of soil fertility management practices in semi-arid regions: a case study of AP." *Economic and Political Weekly* (2011): 56-63.
3. Fishman, R., Kishore, A., Rothler, Y. and Ward, P., 2016. Can Information Help Reduce Imbalanced Application of Fertilisers in India? Experimental Evidence from Bihar (No. 235705). Agricultural and Applied Economics Association.
4. Chander, G., Wani, S.P., Sahrawat, K.L., Dixit, S., Venkateswarlu, B., Rajesh, C., Rao, P.N. and Pardhasaradhi, G., 2014. Soil test-based nutrient balancing improved crop productivity and rural livelihoods: case study from rainfed semi-arid tropics in Andhra Pradesh, India. *Archives of Agronomy and Soil Science*, 60(8), pp.1051-1066.
5. Raju, K.V., Gaur, A. and Wani, S.P., 2015. Poverty reduction approach in South Asia: Rejuvenating centuries old water bodies to improve rural livelihoods, A case of Karnataka state, India.
6. Rajashekar (2011) India's Soil Crisis, *The Economic Times* Mumbai; Date: Jul 12, 2011.
7. Reddy V. Ratna (2003); "Land Degradation in India: Extents, Costs and Determinants", *Economic and Political Weekly*, Vol. XXXVIII, No. 44, November 1-7, 2003

Feedback from agricultural officers on status of soil health card and improvement needed

Questions	Feedback from 27 officers
Status of soil health card scheme	
Status of procurement of maps (patawari maps), field instruments (GPS), and deployment of staff for collection of samples from farmers' fields	Majority of officers mentioned that they don't have patawari maps; instead, they use remote sensing data. They use the GPS system on their mobile and face shortage of manpower and funds.
Is the methodology of soil sample collection (10 ha grid for rained and 2.5 for irrigated) adequate?	Only nine officers said sampling methodology is adequate. Others said it is okay for homogenous land but inadequate in a majority of cases because of variability, topography, and because 10 ha is too big for precise analysis.
Status of soil testing facilities including soil testing labs and mobile labs	Quality of almost all labs is poor. Many of them are capable of measuring only NPK, not micronutrients. Most of them are outdated and face shortage of expert manpower.
Availability of colour Photostat machine, standard paper for printing, printing machines, and mechanism for distribution of soil health cards to farmers	16 officers said laboratory infrastructure is good.
Release of funds, time lags between release of funds to actual utilisation	13 officers said there was a time lag, with insufficient funds.
SHC portal usefulness to farmers: rating from 1 to 5	All officers said portal is useful but farmers are not aware of it and they don't have required facilities.
Overall targets versus achievements as per the annual action plan	23 officers said they achieved their annual targets but faced manpower shortage.
Soil health management (indicators)	
Number of AEOs/TOTs trained under the scheme	Very few out of the total district-level officers are trained.
Number of soil testing laboratories (analysing capacity and utilisation)	20 officers said there is at least one good lab at the district level but its capacity is low, and hence there is a need for more labs.

Are referral labs sufficient?	12 officers said referral labs are not sufficient, and 11 officers said they are sufficient.
How many STLs are equipped with soil-test-based crop response (STCR)?	Only 11 labs at the district level are equipped with STCR.
Number of demonstrations conducted and number of participants involved?	12 officers said they conducted several demonstrations, reaching many farmers.
Socio-economic impact on farmers (indicators)	
Number of SHCs among small and marginal farmers and SC/ST farmers (whether main field located in their farm)	15 officers stated that up to some level, small marginal, and SC/ST farmers get SHC. In some cases, the main field is not located on their farm.
Is there any difference in crops identified in SHCs and actual crops sown by farmers?	Yes, farmers are not taking advice on SHC crops.
Is there any difference in recommended fertilisers based on SHC and actual application?	20 officers said there is significant difference; many believe that recommended doses are not practical.
Status of use of organic manure and bio-fertilisers by farmers	A majority of the officers said farmers are aware about organic manure, and many are using it but they're not using bio-fertilisers. Many said there is no bio-fertiliser available in the market. Organic manure use is greater in case of small-scale farmers, while large-scale farmers don't use it.
Integrated nutrient management practices followed by farmers	A majority of the officers said few progressive farmers are practising it but only in case of horticulture crops (high value crops).
What is the attitude of farmers towards soil health card?	22 officers said farmers are aware and have a positive attitude, but many are sceptical. It will take time earn the trust of farmers.
Soil productivity and economic returns to farmers	A majority of the officers agree that SHC helps farmers in improving the soil health and ultimately increasing productivity, but economic benefits are uncertain because of marketing problems.

Soil Health and Soil Testing: Some Practical Issues (Expert Opinions)

Soil testing is a science-based and time-tested tool for assessment of soil fertility status and soil ailments and nutrient and amendment recommendations. Soil testing as a tool for judicious fertiliser use works on the principle of profitability, which means if all other factors of productivity are optimum and none of them are limiting, there is a high probability to obtain a more profitable response to applied nutrients based on soil testing than those applied on an ad hoc basis. Unfortunately, soil testing in India continues to be a government-driven programme rather than a farmer-driven one. The number of STLs is inadequate vis-à-vis the number of operational holdings, but it is equally true that the demand for soil testing is extremely low. Farmers do not opt for soil testing as a basis for their fertiliser use decisions.

In my opinion, the qualification peaks developed by ASCI hold true when you have a large number of private soil testing laboratories that are self-financed and can gain sufficient income. Unfortunately, the soil testing and soil health care issue is a skill development activity with little income generation. It serves as a decision tool for making fertiliser recommendations.

One of the reasons for the failure of soil testing in making a breakthrough year all these be grouped as follows:

1. Lack of awareness among the farmers
2. Soil sampling techniques being followed
3. Inordinate delay in the analysis of soil samples; sometimes, the results are given 3-4 months after the sample is analysed.
4. Lack of proper laboratory facilities, including frequent power breakdown, availability of chemicals, glassware, water facilities, upkeep of instruments and equipment.

In this context, the ASCI has suggested training for three QPS, namely:

- Soil samplers/collectors
- Soil and water testing lab analysts
- Soil and water testing lab assistants

5. It is worth mentioning that the soil sample collection work is normally taken up during the months of May-June of every year, i.e., 1-2 months in a year, and the rest of the time, the soil samplers do not have any work. Once the sample is collected, where should the sample be sent? Who has to carry out the analysis? Where are the laboratory facilities? Who are the people to man these laboratories? What are the standard procedures being followed? Do we have standard practical manuals for use in STLs?
6. Unfortunately, soil testing is a government driven initiative in India with very little farmer participation.

Some of the reasons behind the failure of STLs indicate that lack of qualification and manpower capability in STL is a major constraint as most of the staff in STL is not qualified and not competent enough to do the analysis.

Another major constraint is the lack of or weak linkage between Dept. of Agricultural and other research organisations such as SAUs, NGOs, etc.

- As per critical analysis, lack of qualified and capable manpower in STL is the major constraint as most of the STL staff is not qualified and not competent enough to perform analysis.
- Training support from research organisations for STL personnel is at a poor level. Consequently, the scientific fertiliser recommendations don't get transferred from SAU and research organisations to extension agencies.
- Lack of knowledge about improved analytical methods, lack of latest and sophisticated equipment, inadequate and lack of automation are the major technical and financial constraints.
- Lack of computers and adequate software in STL also results in non-optimal utilisation of STL capacity.
- To focus on soil testing, there needs to be a separate directorate on soil testing at the state level on the lines of Dept. of Agriculture.
- There is need to appoint a continuous monitoring committee on soil testing at the state, district, *mandal* level. The monitoring committee at the state level may involve Dept. of Agriculture, SAU, fertiliser firms, and farmer representatives.
- Qualified and capable personnel should be appointed.
- Capacity building should be provided through practical methods to all the staff involved in soil testing up to the grassroot level, which includes the method of sampling, laboratory methodology, and interpretation of STCR recommendations for a reasonable period.
- Cross-checking of soil sample analysis should be done at the central laboratory.
- High targets should be reduced and be made reasonable
- A nominal fee should be charged instead of free-of-cost soil sample analysis.
- Farmer awareness and training programs on representative soil sample collection and analysis and application of fertilisers as per STCR technology should be organised as well. A few select farmers from each *mandal* may be sent to other research organisations such as IISS, Bhopal, to learn about the latest technology.
- Motivation techniques for soil testing personnel, such as performance-based promotions, special increments, exposure visits, deputation for related seminars, and workshops should be followed as well.
- Publicity programmes should be conducted on soil testing and information should be sent through pamphlets, newspapers, mass media, even up to the village level, along with fertility status maps and recommendations.
- Financial powers such as contingency amount to a minimal extent to the in-charge of soil testing should meet day-to-day constraints at the laboratory level.
- Soil testing should be made a central subject.
- Soil testing programmes should be brought administratively under the Dept. of Agriculture and technically under the State Agriculture University.
- Transfer of STL staff must be from one lab to the other lab but not to other depts. No additional unrelated works should be assigned to them, such as election duty, etc.

Efficiency Factors

- Collection of soil samples will be done mostly in the summer season, when there is no crop.
- Thousands of samples will reach laboratories for analysis during that period.
- A minimum of 30-35 samples a day (all parameters included) could be a possible target.
- Running the labs at 10,000 samples/year capacity would need very efficient, effective systems and processes in place for 100% capacity utilisation.
- Time saving without compromising on quality and reliability is a challenge.

Practical Issues and Steps to Improve the Facilities

- Most of the laboratories in India are still adopting single extractants for individual estimation of nutrients, such as P, K, and S, etc.
- There is no problem with them, but they consume more time and resources and strain the workers a lot.
- Availability of qualified passionate and dedicated manpower to work in labs is a challenge.
- Manpower with specific qualification, aptitude, and passion should be allotted this job to obtain desirable results.
- They should be trained often to enhance operational efficiency.
- Operations should be incentivised, building motivation and enhancing work efficiency.
- There is inconsistent power supply and unfavourable working environment in rural and semi-urban areas where most of the laboratories are situated.
- In order to have uninterrupted operations, alternative power sources such as generator and battery backup will be useful.
- Adopt those instruments that can work with a chargeable battery as well as power sources.
- Instruments shall be more rugged, reliable, and be able to perform under abnormal working conditions.
- Use of semi-automatic/manually operable equipment (which can be serviced locally, with reduced power dependency) is a more practical option than using fully automated equipment that will have no control over the operator.
- Induction of a new system is important.
- There is a need to update recommendations with the current knowledge based on sound agronomic and research. There should also be a Decision Support System (DSS). Models shall be widely used to make appropriate fertiliser recommendations.
- Networking and frequent upgradation of soil test information in the public domain will help implement new ways and means of fertilisation practices and site-specific nutrient management options on a large scale.
- Delays in reporting of test scale.
- Automation in operation from sample registrations to test results data entry until generation of test reports will save considerable time for routine operations.
- It's great that some states have implemented online or web-based report generations.
- There should be linking of soil health cards and the issue of soil passbooks, linking them to fertiliser subsidies, green manure seeds, organic manure. Only the fertilisers needed for field crops should be provided, and excess use of fertilisers should be avoided.

In order to make soil testing a viable tool for sustainable productivity, the following may be attempted:

- Soil testing is a great tool to assess soil fertility and nutrient supplying capacity
- Timely reporting of soil test results to farmers is crucial in the whole programme.
- Speed and reliability of the operation is the most important.
- Appropriate systems and processes should be in place to effectively implement the programme to get desired results.

Instead of planning the training for soil samplers/collectors, soil and water testing lab analysis, and soil and water testing lab assistants, an alternative suggestion could be that since most of the states offer a two-year diploma in agriculture, such training may be given to these diploma holders, as they already have a preliminary idea about the subject and its importance. During the slack season, their services could be utilised for other agriculture-related activities. Furthermore, in many states, there are VDOs/AEOs (below the rank of AOs) who could drafted the soil and water analysis and the AOs could be trained on advanced aspects such as GPS, induction systems, soil health cards, fertiliser recommendations, etc., as their services could be utilised in a more efficient way.

Field Observations (Compiled across States)

- In some states, the progress is not satisfactory.
- A majority of AEOs and farmers do not have awareness about the process of collection of soil in the fields.
- A majority of farmers did not get SHCs even six months after sample collection as there was a shortage of soil testing laboratories at the local/district level.
- In most cases, farmers were not involved in the collection of the soil samples and their fields were often not covered. Hence, farmers did not take this seriously.
- Farmers expressed that they have more than two parcels of land with varied soil quality but samples were taken from one parcel only. SHC based on one parcel is not suitable to another parcel of land.
- They wanted to take soil samples from each plot.
- Some farmers did not have awareness about the utility of SHC results and lost their cards.
- Even AO/AEO did not explain the content and use of SHCs to a majority of farmers while distributing them.
- Some farmers said that even AEOs were not too aware about the results and need sensitisation about the scheme for proper dissemination of knowledge related to SHC.
- Some illiterate and marginal farmers said that AEOs did not contact them and met only big and progressive farmers.
- Progressive farmers or big farmers followed the recommendations of SHC.
- In most mandals, micro nutrients in the soil are very poor. Marginal and small farmers were unable to adapt these practices due to high cost and also non-availability of nutrients. Only some big farmers are able to use them.
- Some farmers did not follow SHC recommendations because (i) they don't understand the content, (ii) they are costly, (iii) non-availability, (iv) they believe in high doses of fertilisers, (v) they followed traditional application of fertilisers.
- Sensitisation meetings, trainings, distribution of SHCs through elected representatives and follow-ups by the agriculture department are factors that help increase the utility of SHC.
- Farmers expressed that results of soil sample were given for a specific crop. Hence, they could not understand the application of fertilisers for the rest of the crops. SHCs also did not mention the alternative cropping pattern suitable for the local conditions.
- Farmers were not convinced about the soil test because earlier, the agriculture department took soil samples four to five times from some of the farmers, but results were not communicated even after 2-3 years. They got SHC only from this scheme.
- Farmers feel that if they follow SHC recommendations, they face some problems. For example, as per SHC, in tribal areas, if they use more 'N', crop would be attacked by more pests and diseases. Although some farmers adopted the recommended doses of fertilisers, they went back to their traditional methods.
- Some farmers got (one-two farmers for GP) soil tests from NGOs, private fertiliser companies, and KVKs etc. from 2008 to 2013. KVKs and private labs charged for this, but they explained the content to farmers, and most of the farmers practised the recommendations promptly.

- Overall, some farmers reduce fertiliser application by getting advice from progressive, large-scale, or educated farmers.
- After soil test, a majority of the farmers started using micro nutrients that they were not aware of earlier.

Study Limitations

- GPS not captured in some villages.
- In Telangana, new AEOs were appointed in February, 2017. So, they were unable to mobilise farmers who got SHC last year.
- In some mandals/villages, SHCs were not distributed to farmers due to shortage of soil testing labs.

Recommendations:

- This scheme is very useful for farmers. Some constraints and gaps were observed in implementing the scheme.
- Farmers need to be made aware about SHC and its results, importance, etc.
- To build capacity of AOs or AEOs for promotion of fertiliser/nutrient management practices,
- More meetings and trainings are needed for farmers and follow-ups are necessary for the success of the scheme.
- Separate staff is required for SHC to more efficiently promote recommendations to all farmers.
- Establishment of state-of-the-art labs in all district headquarters with large capacities to meet the target for districts is the need of the hour. Mini labs need to be further upgraded for accuracy and speed.
- Micro nutrients may be provided at subsidised prices to the farmers for 2-3 years
- AEOs are not able to actively participate in the SHC scheme as they have multiple tasks and are under pressure
- KVKs, SAUs, ICAR institutions, and private companies should participate in the programme by involving educated local youth in soil sample collection, testing, and distribution.

Review of Recent Studies

No	Reference	Context	Methodology	Analysis
1	Can information help reduce imbalanced use of fertilisers in India? Evidence from Bihar			
	Ram Fishman, Avinash Kishore, Yoav Rothler, Patrick S. Ward, Shankar Jha, R. K. P. Singh.	Three districts from Bihar. Period of study: April 2014	Samples from 509 households (treatment) and 294 households (controlled) were collected. A simplified becker- degroot-marschak method was used to calculate the WTP of farmers for zinc deficiency.	Data from sample fields shows that the gap between recommended fertiliser application and actual application is huge and especially big in the case of urea. The main reasons for farmers' i g n o r a n c e o f recommendations of soil health card were as follows: 1. Farmers didn't understand the contents of the SHC 2. Farmers didn't find the soil analysis and fertiliser recommendations to be reliable or compelling. 3. Other factors such as cost, liquidity, or timely availability of specific fertilisers were constraints.

No	Reference	Context	Methodology	Analysis
2 Soil health mapping and direct benefit: transfer of fertiliser subsidy				
	ICRISAT research report IDC-6		Authors collected data from various government authorities and ground-level farm results after implementation of the bhoomichetana project.	<p>Introduced direct benefit transfer in fertiliser subsidy to increase efficiency and strengthen fertiliser supply chain.</p> <p>Apply integrated nutrient management with emphasis on organic fertilisers.</p> <p>Authors calculated with some assumptions that total benefits with soil health mapping and soil test based fertiliser recommendations along with improved practices will be Rs. 4.33 lakh, while the cost was estimated at Rs 0.254 lakh crore. The benefit-cost ratio would be 17:1. In addition to economic benefits, there would be several environment benefits, employment generation, and enhanced sustainability of Indian agriculture.</p>

No	Reference	Context	Methodology	Analysis
3	Dynamics of soil fertility management practices in semi-arid regions: A case study of AP			
	B Suresh Reddy - EPW	Andhra Pradesh	A total of 360 farmers were gathered from three district of AP and were personally interviewed using a structured schedule and ex post facto research design coupled with case studies and PRA methods. Secondary data was also collected, Fertiliser recommendations were obtained. The data gathered was analysed using average frequency and percentages. The Baletese & Coelli model was used to calculate the average technical efficiency. After regression analysis using frontier production function, the average TE was calculated.	The study was designed for in-depth knowledge on SFM strategies adopted by farmers in dryland areas, examining the significance of SFM practices, and contributing to the overall policy discourse on SFM in semi-arid areas. Farmer's practices are based on their vast experience and rich knowledge of specific local conditions. Combination of chemical fertilisers and FYM was a predominate practice, followed by mixed cropping and legume cultivation. This shows that farmers understand the role of FYM and other organic manure. Analysis also showed a positive sign or the emerging organic markets. The availability of FYM is limited and advertisements cause farmers to use more of chemical fertilisers. Recommendations given by the scientists do not hold true in conditions under which the farmers work. Several recommendations were given based on the analysis.

No	Reference	Context	Methodology	Analysis
4	Impact of saline soil reclamation on enhancing farm productivity and farmers' income in Karnataka – an economic analysis			
	Raju R. Thimmappa K. A.L. Pathan Siddayya - NAAS	Karnataka	A study was conducted in the Ugar Budruk village in Karnataka. About 70% of cultivable land was affected by salinity and waterlogging. SSD was installed between 2009-10 and 2011-12 in 925 ha, covering 644 farmers. The study was based on both primary as well as secondary data collected from 120 sample farmers by area random sampling. The costs and returns were estimated using inputs and outputs and multiplying them by the current year prices as well as using cost concepts. These were then used to compare the pre and post SSD effects.	Soil salinity was major cause of land degradation. Surface drainage technology for saline land reclamation is technically viable, economically feasible, and socially acceptable. Land use was intensified, cropping patterns changed in favour of more remunerative crops, and crop yields increased. Cropping intensity increased, showing a positive effect. There was a significant reduction in the max and min salinity. There was a wide gap in the salinity of drainage water after installation. Mean yield of all crops grown significantly increased. yield increase by 186%. Cost of cultivation increased due to better performance of crops due to demands for more inputs. Increase in net income was largely related to the increase in crop yield due to SSD intervention. Significant increase in net income from off-season crops was also observed. Benefit-cost ratio increased by more than one. Value of land increased. Thus, we can say that to overcome the problem of waterlogging and soil salinity, installation of SSD technology is required.

No	Reference	Context	Methodology	Analysis
5	Soil test-based nutrient balancing improved crop productivity and rural livelihoods: a case study from rainfed semi-arid tropics in Andhra Pradesh			
	<p>Girish Chander, Suhas Wani, Kanwar Sahrawat, Sreenath Dixit, B Venkateswarlu, C Rajesh, P Narsimha Rao, and G Parthasarathi.</p> <p>Archives of agronomy and soil science</p>	Andhra Pradesh	<p>Three to nine villages in eight rainfed districts selected for study. Soil samples were collected from 826 farmer fields in the targeted clusters following the farmer participatory stratified sampling technique. After recommending the ideal fertiliser dose, on-farm observations for the 2008-2011 period were used for the ANOVA technique to test whether balanced nutrient changed farms yields significantly.</p>	<p>Fields of eight districts of Andhra Pradesh showed widespread multi-nutrient deficiencies, including secondary and micro nutrients.</p> <p>In on-farm trials' important rainy and post rainy season crops during 2008-11, significantly higher yield was recorded under the balanced nutrition treatment compared to traditional farmer practice (more NPK, none micro nutrients).</p> <p>Balanced nutrition while increasing crop yields maintained plant nutrient composition.</p> <p>Harvest in Nalgonda district showed higher contents of soil organic C and available nutrients such as P, S, B, and Zn in plots with balanced nutrition, which are apparently responsible for residual benefits in the succeeding seasons.</p> <p>In the absence of balanced nutrition, farmers are apparently losing 8% to 102% of current yield levels in season 1 and 15% to 24% in each of the succeeding three to four seasons.</p>

National Policies on Soil Management			
Plan	Years	Policies	Progress
1st FYP	1951 - 1956	Soil conservation associatioans	1953 - Central soil conservation board set up
		Establishment of a Soil Conservation Branch at the Forest Research Institute, Dehradun	Sum of about Rs. 1.6 crore was spent
		Soil and land utilisation survey	Contour bunding and terracing of about 700,000 acres of agricultural lands
		Soil conservation in community development projects	Desert Afforestation and Research Station was set up at Jodhpur
		Soil conservation in river valley project areas	Eight regional research-cum-demonstration centres were established
		Central Land Utilisation and Soil Conservation Organisation at the Centre	
		Land Utilisation and Soil Conservation Board in every State.	
2nd FYP	1956 - 1961	Training centres have been established by the Government of India at the research stations	Rs. 18 crore spent
		Pilot project demonstration centres to be established in different parts of the country for giving demonstrations of soil conservation practices to farmers	An area of 2 million acres covered under contour bunding and terracing
		Erosion control programmes should therefore be accompanied by appropriate programmes of education and resettlement	An integrated all-India Soil Conservation and Land Use Survey was initiated
		Agency of the national extension service	70 officers and 900 assistants have been trained
		Village panchayat should become responsible for soil conservation measures and ensuring minimum standards of land management by individual cultivators. They should also receive such technical and financial assistance as may be needed.	40 demonstration projects, each covering about 1,000 acres, were undertaken on a catchment basis
		Central fertiliser pool	The Desert Afforestation and Research Station at Jodhpur was reorganised as a Central Arid Zone Research Institute in collaboration with UNESCO

3rd FYP	1961- 1966	Intensive Agriculture Development program (IADP) was the first major experiment of the Indian government in the field of agriculture, and it was also known as a “package programme” as it was based on the package approach	An area of 1.9 million hectares was reclaimed
		River valley projects	The work was initiated in 13 river valley projects
		Reclamation of alkaline and usar lands	
		The Central Arid Zone Research Institute For desert reclamation	
		Empowering State Governments to frame soil conservation schemes for the basin of a river or a stream or for groups of villages has been recommended	
		The Fertiliser Distribution Enquiry Committee	
		A Central Fertiliser Marketing Corporation	
		Set up a Central Institute of Pedology and Soil Mechanics.	
4th FYP	1969 - 1974	Adopt an "area saturation" approach so as to treat all types of land on a complete watershed basis.	The chemical fertilisers in terms of nitrogenous and phosphatic recorded 10.58 lakh and 4.1 lakh tones, respectively, ending in 1973-74
		A resource inventory unit was established at the centre during 1966-67	At the end of 1968-69, there were 65 soil-testing laboratories all over the country with a total annual capacity of handling 1.08 million samples
		The scheme of soil conservation in the catchments of river valley projects	
5th FYP	1974 - 1979	Sharp increase in the prices of food, fertilisers, and oil seriously upset the assumption on which draft had been framed	The total coverage under soil and water conservation was 21.7 million hectare in 1977-78
		Soil and water conservation programmes have also been taken up with institutional credit support, and the targets of physical performance are likely to be achieved.	The area covered by soil conservation measures is only 23.40 million hectares so far.
		A centrally sponsored scheme of soil conservation in the catchments of river valley project (RVP) was started	
		Soil and water conservation programmes are being taken up on the watershed approach	

6th FYP	1980 - 1985	Flood- Prone Rivers (FPR) was started	Under soil and water conservation measures, the achievement is about 6 million hectare
		Small watersheds with an area of 1,000-2,000 hectares, treatment of which is practicable and manageable, will have to be taken up increasingly	The area treated under soil water conservation measures aggregates to about 29.4 million hectares
		Integrated watershed management in the catchments of eight flood prone rivers of the Indo-Gangetic Basin will also be taken up.	8.4 million tonnes of fertilisers are reported to have been achieved by the end of 1984-85.
			It consists of 5.6 million tonnes of N, 1.9 million tonnes of P, and 0.9 million tonnes of K
7th FYP	1980 - 1985	The Centrally Sponsored Scheme of soil conservation in the catchments of river valley projects and Integrated Water Management in the catchments of eight flood-prone rivers in the Indo-Gangetic basin will be continued and intensified.	The consumption of fertilisers will be increased from 8.4 million tonnes in 1984-85 to 13.5-14.0 million tonnes in 1989-90.
		National Watershed Development Programme for Rainfed Agriculture.	The consumption of fertilisers (N+.P+K) rose to a level of 11.3 million tonnes in the final year of the Seventh plan.
		Programmes have been taken up to encourage the use of bio-fertilisers.	Forty blue green algae sub-centres were established for the production of algae culture through field multiplication programme under the national project for bio-fertiliser development.
		Five central sector schemes, including National Project on Development and Use of Bio-Fertiliser and National Project on Quality Control, were implemented.	Soil and water conservation activity in 27 catchments were taken up in 17 states, covering 2.4 million hectares by the end of the Seventh Plan with a reported expenditure of Rs. 307 crore.
		Schemes for balanced and integrated use of fertilisers and a National Project on Development of Fertiliser Use in Rainfed Areas were introduced.	
		Schemes on soil conservation in the catchments of interstate river valley projects and the flood-prone rivers, reclamation of alkaline (usar) soil, control of shifting cultivation, and development of ravine areas were taken up.	

8th FYP	1992 - 1997	Efficient use of chemical fertilisers, recycling of organic wastes, and use of bio-fertilisers have an important place in the sustainable agricultural development process	An amount of Rs 40.826 crore was released to the States of the North-East region and 0.67 lakh ha. The area was treated through treatment packages.
		A centrally sponsored scheme of reclamation of alkali soil was continued from the seventh plan.	The level of 14.3 million tonnes was reached in 1996-97.
		The scheme of Watershed Development Project in Shifting Cultivation Areas (WDPSCA) was launched in seven north-eastern states.	Phosphatic and potassic fertilisers were decontrolled in August 1992. Only urea (nitrogenous fertiliser) continued to be under the price control system and involves a heavy subsidy for keeping the farm gate prices low.
		To make soil treatment cost-effective, the guidelines emphasised on the vegetative conservation measures with active involvement of the beneficiaries and the non-governmental organisations (NGOs).	The All India Soil and Land Use Survey has covered 1155.74 lakh ha so far under priority delineation survey and 85.65 lakh ha under detailed soil survey.
		The "Application of Remote Sensing Technology for Soil Survey and Land Use Planning" scheme.	Up to the end of the Eighth Plan, only 17.96 m ha had been covered/treated.
		Seventh Plan Centrally Sponsored Schemes of Soil Conservation in the catchments of River Valley Projects (RVP) and Integrated Watershed Management in the catchments of Flood Prone Rivers (FPR) were continued.	
9th FYP	1997 - 2002	Community and village institutions will be encouraged to participate in protecting natural resources from degradation.	1.5 lakh ha. has been treated with an expenditure of Rs 82 crore under watershed management.
		Programmes for regeneration of land and water resources will be strengthened	An area of 0.60 m.ha. Out of 3.5 m.ha. of alkali, land has been reclaimed until the end of 2002-03.
		Macro Management Mode brings together 27 centrally sponsored schemes under one umbrella - Reclamation & Development of Alkali & Acid Soil (R&DAAS)	
		Centrally sponsored scheme of reclamation of alkali soil - extension of the scheme to all other states of India was approved where alkali soil problems exist as per scientific parameter.	
10th FYP	2002 - 2007	Unsustainable practices such as excessive use of water together with imbalanced use of fertilisers were seen.	Around 22 million hectares of degraded land was reportedly treated under these various schemes at a cost of Rs. 8,810 crore.

		A condition that a certain percentage of allotted land (say 40 or 50%) must be utilised for tree cover can be stipulated so as to increase the crown area for improving the environmental and ecological conditions	
		Soil survey and land degradation mapping of the entire country would be conducted from the Tenth Plan onwards in a mission mode approach.	
		Adoption of Integrated Nutrient Management (INM)/Integrated Plant Nutrient Supply (IPNS)	
11th FYP	2007 - 2012	The Centre introduced a new scheme, the 'National Project on Management of Soil Health & Fertility' (NPMSH&F).	Fertiliser consumption rose over 30% during the Eleventh Plan, the main reason being that the world prices of all fertilisers and feedstock doubled since 2006
		Soil health cards and strengthening of soil testing labs and expanding their testing capacity	By 2010–11, there were 1,049 soil test labs in the country with a soil analysis capacity of 106 lakh soil samples per annum
		Minimum investment of Rs 36,000 crore in Natural Resources Management (NRM)	
		Nutrient-based subsidy (NBS) system was adopted	
12th FYP	2012 - 2017	National Mission for Sustainable Agriculture (NMSA). Conceived originally as part of the National Action Plan on Climate Change (NAPCC)	
		The National Project on Management of Soil Health and Fertility and the Rainfed Areas Development Programme (RADP)	
		Conservation agriculture (CA), integrated nutrient management, carbon sequestration, erosion control, saline and alkaline soils management, legislation for soil protection, development of remote sensing and GPS-based Decision Support System (DSS), and amelioration of polluted soil are required to rejuvenate deteriorated soils.	

Regression analysis: Determinants of performance indicators of the SHC scheme (District level)

Variables included in the district-level regression analysis

Variables	Source
1. Number of bank accounts	RBI databank
2. Rural households with computer/laptop and internet	Census 2011
3. Rural households with mobile	Census 2011
4. Samples entered (collected)	Soil health card portal
5. Number of farmers covered	Soil health card portal
6. Samples tested	Soil health card portal
7. Soil health cards printed	Soil health card portal
8. Net area sown (ha)	Data provided by CRIDA
9. Net irrigated area (ha)	Data provided by CRIDA
10. N, P, K consumption in both <i>rabi</i> and <i>kharif</i>	Fertilisers statistics, 2010-11
11. Number of cultivators	Census 2011
12. Total rural bank branches (both PSU and commercial)	RBI bank branch statistics, 2010
13. Road density	Road statistics taken from Pradhan mantri gram sadak yojana and divided by geographical area.
14. Literacy	Census 2011
15. Number of SC and ST population	Socio-economic caste census
16. Number of soil testing labs and magnesium, boron, zinc testing capacity.	Soil health card portal
17. Yield	District-wise data collected from Ramesh Chand's working paper 'Instability and regional variation in Indian agriculture' published by NCAP.

Basic regression model:

- 1) Y (samples entered[collected])= f (number of bank accounts, households with computer and internet, households with mobile, net irrigated area, net sown area, total fertiliser consumption, rural bank branches, road density, literate population, number of SC and ST population, number of labs)

Determinants of samples collected (district level)

Dependent variable: samples collected	B	Significance level
(Constant)	-1963	.587
Total fertiliser use (Tonnes)	.233	.000
No. of bank accounts	.032	.000
Net sown area (ha)	.050	.000
Net irrigated area (ha)	-.096	.000
Households with computer/laptop an internet	-3.481	.000
Households with mobile	.071	.006
No. of labs	1098.157	.021
Total rural bank branches (both PSU and commercial)	78.164	.041

- 2) Y (samples tested)= f (number of bank accounts, households with a computer and internet, households with mobile, net irrigated area, net sown area, total fertiliser consumption, rural bank branches, road density, literate population, number of SC and ST population, number of labs)

Determinants of samples tested (district level)

Dependent variable: samples tested	B	Sig.
(Constant)	-2577	.438
Total fertiliser use (Tonnes)	.241	.000
No. of bank accounts	.040	.000
Net sown area (ha)	.039	.000
Net irrigated area (ha)	-.102	.000
Households with computer/laptop and internet	-3.286	.000
Total rural bank branches (both PSU and commercial)	88.392	.013
Households with mobile	.052	.030

- 3) Y (Soil health card printed)= f (number of bank accounts, households with computer and internet, households with mobile, net irrigated area, net sown area, total fertiliser consumption, rural bank branches, road density, literate population, number of SC and ST population, number of labs)

Determinants of SHCs printed (district level)

Y = SHCs printed	B	Sig.
(Constant)	16000	.133
No. of bank accounts	.126	.000
Net sown area (ha)	.101	.000
Total rural bank branches (both PSU and commercial)	474.2	.000
Net irrigated area (ha)	-.217	.000
Total fertiliser use (Tonnes)	.288	.021

1. Farmer Questionnaire

Farmer questionnaire

Soil Health Card Scheme 2017

**National Institute of Agricultural Extension Management (MANAGE),
Hyderabad**

Section I: Household Details

1. State
2. District
3. Block
4. Village
5. Name of the farmer: Mobile No.
6. Social group: FC/OBC/SC/ST
7. Age:
8. Sex:
9. Family size: M- ; F- ; C-
10. Number of family members working on the farm: M- ; F-
11. Education (no. of years of head of family):
12. Landholding
 - a. Irrigated area.....number of plots
 - b. Rainfed area.....number of plots.....
13. Annual net income after deducting costs (Rs.)
 - a. Agriculture:
 - b. Livestock:
 - c. Labour:
 - d. Others:

Section II: About soil health card

14. Are you aware about soil health card? Y/N
15. Are you aware of the soil health card portal? Y/N
16. How many soil samples have been taken from your village?
17. Are the results discussed among farmers in the village? Y/N
18. Does the soil health card portal give useful information?
19. Details of soil health card scheme operation in your village.

Section III: Initial activities under the SHC scheme

Purpose	Participation Number (Y/N)	Distance from village	Number of days	Nature of Inputs	Benefits if any
Mobilisation (farmers)					
Meetings conducted					
Exposure visits					
Trainings conducted					

Section-IV Cropping Pattern (for all)

Crop name	Season K/R/ annual	Area (acre)	Yield (per care)	Fertiliser use Urea (bags)	DAP (bags)	MoP (bags)	Gypsum (Kg)	Micro Nutrients (Kg)	Bio- fertili sers (Kg)	Liming material (Kg)	Others	Total Cost
Before SHC scheme												
Crop 1												
Crop 2												
Crop 3												
Last year 2016-17												
Crop 1												
Crop 2												
Crop 3												

Section V: Soil Health Card holders

20. Are you having a soil health card? Y/N
- If yes
21. In which month and year was soil testing done?
22. In which year and month did you get the SHC?
23. From where did you get the SHC?
24. Are you able to understand the information in the soil health card? Y/N
25. Are you using fertilisers and micro nutrients as per the SHC? Y/N
26. Have you received any financial assistance from the government? Y/N
27. If you are not following recommendation, why is that?
28. How many of your plots are covered for soil testing?
- One / none /some/all
29. Are the recommendations suitable for all your plots?
- Y/N
30. How many soil samples were taken from your farm?
31. Do you think the samples represent all the soil types in your farm? Y/N
32. If no, how many samples are required?

33. Are the results provided in time? Y/N
34. Did the agricultural extension worker explain the content of SHC? Y/N
35. Who informed you about the content of SHC?
 - a. KVK
 - b. Agricultural Extension Officer/Agricultural Officer
 - c. Scientist of SAU/ICAR
 - d. NGOs
36. Are the recommendations practical? Y / N
37. Are the recommended inputs easily available in the market?
38. Did you face any difficulties in adopting the practices? Y/N

Section 4: Perception of the SHC scheme

39. What is the main benefit (s) from SHC?
40. Would you like to suggest any improvement to the SHC system? Y/N
41. If yes, provide details.
42. Would you like to take your soils to private testing labs? Y/N
43. Are you willing to pay for the soil health card?
44. Observations from the investigators.
45. Name of the investigator.

State-wise fund released and unspent balance under the Soil Health Card scheme
(As on 30.01.2017)
(Rs. In lakh)

State	2014-15			2015-16				2016-17				Total		
	Released	UC	UC	Allocation	Released	UC	UC	Allocation	Released	UC	UC	Released	UC	UC
		Received	Pending			Received	Pending			Received	Pending		Received	Pending
Andhra Pr.	124	124	0	706	458	458	0	1394	697	0	697	1279	582	697
Arunachal Pr.	12	12	0	30	17	17	0	50	0	0	0	29	29	0
Assam	51	0	51	253	142	3	139	619	0	0	0	193	3	190
Bihar	253	150	103	674	488	240	248	1289	0	0	0	742	390	351
Chhattisgarh	66	66	0	383	281	195	86	685	343	0	343	690	261	429
Goa	12	0	12	17	12	0	12	19	0	0	0	24	0	24
Gujarat	81	81	0	836	603	420	184	1057	529	0	529	1213	501	712
Haryana	34	17	18	401	288	210	78	440	0	0	0	322	227	96
Himachal Pr.	24	24	0	71	47	47	0	59	29	0	29	101	71	29
J & K	32	0	32	141	77	49	28	326	0	0	0	109	49	60
Jharkhand	51	36	15	73	54	31	23	128	64	0	64	169	66	103
Karnataka	127	101	26	894	650	650	0	2060	1030	649	381	1807	1400	407
Kerala	113	0	113	83	61	0	61	132	0	0	0	174	0	174
Madhya Pr.	143	99	44	1228	888	465	423	2747	1374	329	1044	2404	894	1511
Maharashtra	216	216	0	1285	932	932	0	2430	1215	1215	0	2363	2363	0
Manipur	13	13	0	31	18	12	6	63	0	0	0	30	25	6
Meghalaya	14	0	14	36	21	0	21	43	0	0	0	34	0	34
Mizoram	12	0	12	24	14	14	0	36	0	0	0	26	14	12
Nagaland	13	13	0	40	22	22	0	74	0	0	0	36	36	0
Odisha	80	80	0	371	270	121	148	609	304	304	0	655	506	148
Punjab	26	0	26	424	305	0	305	958	0	0	0	331	0	331
Rajasthan	143	143	0	1213	876	90	786	2217	1108	0	1108	2128	233	1894
Sikkim	12	12	0	20	12	8	4	25	0	0	0	24	20	4
Tamil Nadu	132	132	0	652	470	470	0	1290	645	0	645	1246	601	645
Telangana	93	93	0	544	353	353	0	937	468	0	468	915	446	468
Tripura	18	18	0	40	23	23	0	79	0	0	0	41	41	0
Uttar Pr.	354	354	0	2387	1716	282	1434	4884	2442	0	2442	4513	637	3876
Uttarakhand	23	23	0	119	65	44	22	236	0	0	0	89	67	22
West Bengal	117	117	0	667	480	480	0	1460	0	0	0	597	597	0
Total	2390	1926	464	13642	9644	5636	4008	26346	10249	2497	7751	22282	10059	12223

Soil Health Card

 Department of Agriculture & Cooperation
 Ministry of Agriculture & Farmers Welfare
 Government of India

 Directorate of Agriculture
 Government of Goa

 Soil Health Card No. : _____
 Name of Farmer : _____
 Validity : From _____ To _____

SOIL HEALTH CARD				Name of Laboratory	
Farmer's Details					
Name					
Address					
Village					
Sub-District					
District					
PIN					
Aadhaar Number					
Mobile Number					
Soil Sample Details					
Soil Sample Number					
Sample Collected on					
Survey No.					
Khasra No. / Dag No.					
Farm Size					
Geo Position (GPS)	Latitude:	Longitude:			
Irrigated / Rainfed					

SOIL TEST RESULTS					
S. No.	Parameter	Test Value	Unit	Rating	
1	pH				
2	EC				
3	Organic Carbon (OC)				
4	Available Nitrogen (N)				
5	Available Phosphorus (P)				
6	Available Potassium (K)				
7	Available Sulphur (S)				
8	Available Zinc (Zn)				
9	Available Boron (B)				
10	Available Iron (Fe)				
11	Available Manganese (Mn)				
12	Available Copper (Cu)				

Secondary & Micro Nutrients Recommendations		
Sl. No.	Parameter	Recommendations for Soil Applications
1	Sulphur (S)	
2	Zinc (Zn)	
3	Boron (B)	
4	Iron (Fe)	
5	Manganese (Mn)	
6	Copper (Cu)	
General Recommendations		
1	Organic Manure	
2	Biofertiliser	
3	Lime / Gypsum	
<div> <div>International Year of Soils</div> <div>2015</div> <div> </div> <div>Healthy Soils for a Healthy Life</div> </div>		

Fertilizer Recommendations for Reference Yield (with Organic Manure)					
Sl. No.	Crop & Variety	Reference Yield	Fertilizer Combination-1 for N P K		Fertilizer Combination-2 for N P K
1	Paddy (Dhaan)				
2					
3					
4					
5					
6					