Grazers and fires. Their role in shaping the structure and functioning of the Río de la Plata Grasslands

JOSÉ MARÍA PARUELO ^{1,2,3}, MARTÍN OESTERHELD ², ALICE ALTESOR ³, GERVASIO PIÑEIRO ^{2,4},
CLAUDIA RODRÍGUEZ ³, PABLO BALDASSINI ², GONZALO IRISARRI ^{2,6}, LUIS LÓPEZ-MÁRSICO³,
VALÉRIO D. PILLAR⁵.

¹INIA La Estanzuela. Ruta 50 km 11. Colonia, Uruguay. ²IFEVA, Universidad de Buenos Aires, CONICET, Facultad de Agronomía.
Buenos Aires, Argentina. ³IECA, Facultad de Ciencias, Universidad de la República. Montevideo, Uruguay. ⁴Facultad de Agronomía,
Universidad de la República. Montevideo, Uruguay. ⁵Departamento de Ecología, Universidade Federal do Rio Grande do Sul. Porto Alegre,
Brasil. ⁶Sustainable Agriculture Sciences, Rothamsted Research. North Wyke, UK

José María Paruelo: jparuelo@inia.org.uy; Martín Oesterheld: oesterhe@agro.uba.ar; Alice Altesor:
aaltesor@fcien.edu.uy; Gervasio Piñeiro: pineiro@agro.uba.ar; Claudia Rodríguez:
claudia@fcien.edu.uy; Pablo Baldassini: pbaldass@agro.uba.ar; Gonzalo Irisarri: irisarri@agro.uba.ar
gonzalo.irisarri@rothamsted.ac.uk; Luis López-Mársico: luislo81@gmail.com; Valerio Pillar:
vpillar@ufrgs.br

ABSTRACT. The Río de la Plata Grasslands (RPG) are one of the largest areas of open ecosystems (grasslands, shrublands and savannas) in the world. Historically these systems have experienced, and continue to experience, an enormous loss of natural habitats. Moreover, their importance has been largely invisible in comparison to forested systems. The remaining area of open ecosystems in the RPG region varies according to the source from 38% to 58% of the original area. Open Ecosystems (OE) are a special case of agroecosystems because they can combine the provision of both provisioning and regulating and supporting ecosystem services (ES). Preserving the provision of ES in these natural habitats depends, in part, on understanding the role of the two main disturbances operating in them: grazing and fire. Although these two disturbances are natural components of OE, both are manipulated by humans. In this paper we reviewed the role played by fire and grazing in the structure and functioning of the RPG starting from the late Pleistocene and summarizing current evidence on the effects of fire and grazing on vegetation, fauna and biogeochemical processes. The evidence indicates that among agricultural activities, direct grazing systems in OE have the lowest environmental footprint. And, at the same time are the key for habitat preservation and ES supply. Overall, the OE of the RPG still represent a high proportion of the area, are capable of covering 2.5% of the world's population needs of high quality protein, and at the same time guarding the regulation of key processes.

[KEYWORDS: Disturbances, grasslands, ecosystem services, environmental footprint]

RESUMEN. Herbívoros y fuegos. Su rol en la configuración de la estructura y el funcionamiento de los pastizales del Río de la Plata. Los Pastizales del Río de la Plata (PRP) son una de las áreas más extensas de ecosistemas abiertos (pastizales, arbustales y sabanas) del mundo. Históricamente estos sistemas han experimentado, y lo siguen haciendo, una enorme pérdida de hábitats naturales. Más aún en buena medida su importancia ha sido invisibilizada frente a sistemas boscosos. El área remanente de ecosistemas abiertos (EA) varía según las fuentes entre un 38% a un 58% del área original. Los EA son agroecosistemas particulares porque pueden combinar el suministro tanto de servicios ecosistémicos (SE) de provisión como de regulación y soporte. La maximización del suministro de SE en estos hábitats naturales depende, en parte, de la comprensión del papel de las dos principales perturbaciones que operan en ellos: el pastoreo y el fuego. Aunque estas dos perturbaciones son componentes naturales de los EA, ambas son manipuladas por prácticas humanas. En este artículo revisamos el papel del fuego y el pastoreo en la estructura y el funcionamiento del PRP partiendo de su papel durante el final del Pleistoceno y el Holoceno y resumiendo evidencias de los efectos actuales del fuego y el pastoreo sobre la vegetación, la fauna y los procesos biogeoquímicos. Las evidencias muestran que los sistemas ganaderos en EA tienen, en dimensiones claves de la huella ambiental tales como la preservación de hábitats y la oferta de SE, el menor impacto entre las actividades agropecuarias. Los valores estimados de producción de carne equivalente en los EA representan una alta proporción de la producción total de la región, un volumen de producción capaz de cubrir el consumo de un 2,5% de la población mundial.

[Palabras clave: perturbaciones, pastizales, servicios ecosistémicos, huella ambiental]

Short title: Grazers and fire's role in RP grasslands

INTRODUCTION

Grasslands, savannas and shrublands are grassy or Open Ecosystems (OE) (Bond, 2019) occurring in a wide range of mean annual precipitation, from 200 to 1500 mm, and temperature, from 0 to 30 °C (Lauenroth 1979). Their main physiognomic characteristics include, among other features, a relatively continuous herbaceous layer with or without scattered woody components. Temperate and subtropical subhumid grasslands and savannas, in particular, are one of the most threatened biomes of the world (Hoekstra et al. 2005). The vast plains that once dominated central parts of Eurasia, and North America, eastern parts of Africa and the southeastern end of South America have been profoundly altered by agricultural expansion. Surprisingly, such changes have acquired less popular visibility than those occurring in, for example, wetlands or forest (Overbeck et al. 2015). Denial becomes clear when the area devoted to conservation is considered. Hoekstra et al. (2005) calculated a Conservation Risk Index (CRI) and temperate grasslands, savannas and shrublands ranked first. Moreover, these ecosystems are also threatened by worldwide initiatives of massive afforestation (Bastin et al. 2019) to deal with global changes issues, neglecting the importance of OE in biodiversity conservation, C sequestration and ecosystem services in general.

Grazing, droughts and fire are key factors in shaping these OE (Oesterheld et al. 1999; Gibson 2009). Year-to-year changes, and even larger-scale trends in climatic variables, affect water availability, a major driving force of the structure and function of OE (Sala et al. 1988; Lauenroth and Sala 1992; Paruelo et al. 1999; Paruelo and Lauenroth 1996, 1998; Paruelo et al. 1998; Oesterheld et al. 1992). Grazing and fires are two paradigmatic disturbances in OE and a major pathway to release fixed C back to the atmosphere (Gibson 2009; Chapin 2011). Grazing is a complex syndrome that involves biomass removal, selectivity, trampling, defecation, and urination (Mikola et al. 2009). The combined effects of these components of grazing determine changes in physiognomy (i.e. Sala et al. 1996; Altersor et al. 2019; Ferreira et al. 2020), soil properties (i.e. Milchunas and Lauenroth 1993; Taboada and Lavado 1988) and ecosystem functioning (Rusch and Oesterheld 1997; Altersor et al. 2005). The frequency and intensity of fires determine a pattern of biomass removal, nutrient volatilization, and ash deposition which affect the whole ecosystem (Daubenmire 1968; Vogl 1974; Hulbert 1988; Hobbs et al. 1991; Bond and Keeley 2005; Veldman et al. 2015; Buisson et al. 2019).

One of the largest plains in the world are the Río de la Plata Grasslands (RPG) region which extend in the eastern part of South America from 28 to 38° Latitude S and from 50 to 66.5° Longitude W. They cover about 853000 km² and represent one of the most diverse, largest and less transformed grassland areas in the world, despite the important land cover transformation that they experienced (Oyarzabal et al. 2020). Soriano (1992) defined nine subregions within the whole region (Figure 1): the Northern and Southern Campos and seven units within the Pampa: Mesopotamic, Rolling, Flat Inland, West Inland, Austral, Fluvial and Flooding Pampa. They occupy the vast and continuous plains of central-eastern Argentina, Uruguay and southern Brazil that surround the Río de la Plata estuary and its major tributaries. The RPG covers a range of average annual temperature from 20 °C in the north to 13 °C in the south and an annual rainfall from 1500 mm in the northeast to 400 mm in the southwest (Paruelo et al. 2007). C3 and C4 grasses are the co-dominant life forms, with a low cover of trees and shrubs. Other physiognomic types, such as meadows, savannas, shrublands and forests, occupy a relatively small area and become abundant in particular environments (Boldrini 1997; Overbeck et al. 2007; Perelman et al. 2001, 2017; Oyarzabal et al. 2018; Lezama et al. 2019; Batista et al. 2014). Forests are mainly restricted to riparian corridors along the large Paraná and Uruguay rivers and their tributary streams, to rocky hills and ravines, and to azonal soils formed on top of shell debris deposits (Lewis and Collantes 1973; Cabrera et al. 1976; León et al. 1979; Morello et al. 2012; Brazeiro et al. 2020). An abundant regional literature summarizes the biotic and abiotic characteristics of this biome (Vervoot 1967; Soriano 1992; Paruelo et al. 2005; Pillar et al. 2009; Perelman et al. 2001, 2017; Oyarzabal et al. 2020).

Several studies described the changes in land cover across the whole region and the spatial heterogeneity of the transformation (i.e. Hall et al. 1992; Guerschman et al. 2003; Baldi et al. 2006; Baldi and Paruelo 2008; Baeza et al. 2014; Baeza and Paruelo 2020, and recently MapBiomas initiatives).

Changes are mainly associated with the substitution of permanent grazing lands with cash crops or tree plantation expansion (Vega et al. 2009). At least three cartographic products show the present distribution of grazing lands in the whole region (Figure 1) (Baeza and Paruelo 2020; Copernicus Global Land Cover (Buchhorn et al. 2020), MapBiomias Pampa and Atlantic Forest Trinational initiatives (Souza et al. 2020; Vallejos et al. 2021; Milkovic et al. 2021)). From these three independent sources, it is clear that the current area of remnant OE is controversial. The sources of this controversy are associated with the classification protocols used to construct each one of the three maps (data sources, classification techniques, etc.), the conceptual resolution of the estimates, and the quality of “ground-truth” data may also jeopardize the estimates. Moreover, the actual definition of “natural” grasslands is controversial (Allen et al. 2011). Old sowed pastures, intersowed grasslands or old fields may be included or not as “natural grasslands” depending on the criteria used. The estimates of remnant OE ranged from 328739 ha (MapBiomias project) to 498987 ha (Baeza and Paruelo 2020) (a 38% and 58% of the estimated original area respectively). In this article we used an ensemble of the three sources presented in Figure 1. We considered an area as grassy Open ecosystems OE if it was classified as such by at least two sources (Figure 1).

The main economic activity on the remnant grasslands, savannas and shrublands (OE) of the RPG is cattle and sheep ranching (Oyarzabal et al. 2020). Livestock production has been strongly questioned from an environmental perspective. Probably the FAO’s document “Livestock’s long shadow” from 2006 started the controversy on the negative effects of livestock farming on the environment but a series of articles and books have further elaborated on this issue (i.e. Oppenlander 2013 for the general public and Garibaldi et al. 2018 in an academic context). The concerns are linked, in general, to three aspects: trophic efficiency of grain-feed production in confined systems, the destruction of natural habitats (mainly forests) and biodiversity losses and the impact on ecosystem services supply, particularly on biodiversity and greenhouse gases (GHG) emissions.

Open ecosystems (OE) are a special kind of agroecosystems because they may combine the supply of both provisioning (i.e. meat and wool) and regulating and supporting ecosystem services (ES) (i.e. C sequestration, water supply, biodiversity conservation). Maximizing ES supply on these natural habitats depends, in part, on understanding the role of the two main disturbances that operate on them: grazing and fire. Though these two disturbances are natural components of OE, both are manipulated by humans. In this article we reviewed the role of fire and grazing on the structure and functioning of the RPG. We first present a synthetic review of the importance of both disturbances over the end of Pleistocene and the Holocene. Then, the evidence of the current effects of fire and grazing on vegetation, fauna and biogeochemical processes is presented. Finally, we evaluate the potential for meat production in native grasslands, shrublands and savannas of the RPG and two estimates of the environmental footprint of ranching in OE.

The evolution of the grass-dominated vegetation and the influence of large herbivores and fire

At present, grass-dominated vegetation characterizes the RPG, which has been an important natural asset for the economy of the region since the introduction of livestock by Europeans. These ecosystems present a long evolutionary history of herbivory by large grazers that existed throughout South America (MacFadden 2005; Strömberg 2011). Data from plant silica (phytoliths) in sediment cores collected in Patagonia indicate that grass-dominated vegetation developed under the influence of large grazers about 18 million years before present (BP) (Strömberg et al. 2013), and possibly since about 8 million years BP in the RPG (Strömberg 2011). The now extinct megafauna of large herbivores included species such as *Equus neogeus*, *Toxodon platensis*, and *Notiomastodon platensis*, which presented C4 plants in their diets, as inferred by carbon isotopes, a clear indication that they were grazers (MacFadden 2005; Morosi and Ubilla 2019; Domingo et al. 2020; Omena et al. 2020).

Large herbivores became extinct in South America by the end of the Pleistocene and the beginning of the Holocene (Prado et al. 2015), and the extinction process was likely human-driven

(Prates and Perez 2021). The time window of ca. 10 thousand years without grazing by large herbivores until the introduction of domestic livestock was not enough, in evolutionary terms, for substantially changing the regional pool of grassland species, as shown by pollen and phytolith evidence (e.g. Behling and Pillar 2007; Iriarte 2006). Further, by the end of the Pleistocene, fire frequency increased in some regions of the RPG (Behling et al. 2005), likely due to the change from cold and dry to warm and humid climatic conditions, which increased primary productivity, and to the direct fire ignition by humans. In addition, the lack of large grazers may have been an important factor for this change of fire frequency. Thus, without grazing, fire may have played the role of herbivory (Bond and Keeley 2005) by maintaining in humid ecosystems an analogous selective pressure towards traits that are also adapted to grazing, such as fast regrowth after biomass loss (Díaz et al. 2007). Nevertheless, in addition to fast regrowth, it is expected that the evolution under grazing in humid conditions led also to plants presenting a prostrate habit.

This evolutionary long-term dynamics of plants under grazing and, possibly, under fire, was embedded into the dynamics of the climate during the Quaternary. The RPG, which at present is characterized by a strong NE-SW precipitation gradient (from ca. 1500 mm in southern Brazil to 400 mm in the West Pampa in Argentina) has undergone climatic shifts during the late Pleistocene and the Holocene, when species from the north have expanded their distribution towards the south and vice-versa (Tonni et al. 1999; Quattrocchio et al. 2008). This has been reflected in the vegetation (Behling and Pillar 2007; Iriarte 2006) and in the soils (Quattrocchio et al. 2008). These climatic changes explain the “aeolian deposits interbedded with paleosols” that are observed in the Pampa (Iriarte 1999). At present, species richness is related to this climatic gradient, with higher diversity towards the NE, where the climate is warmer, and the precipitation regime is less seasonal (Bergamin et al. 2021; Perelman et al. 2017).

The last 500 years of the RPG were characterized by increased human influence. First, livestock reinstated grazing as a disturbance throughout the area and reached an average density six times greater than the density of large natural grazers in comparable ecosystems of the world (Soriano 1992; León et al. 1984; Oesterheld et al. 1992). Thus, livestock grazing was a novel disturbance at a very short-time scale, which led some authors to assign these grasslands a short evolutionary history of grazing. However, as seen above, in a broader framework, these 10 thousand years may be seen as a brief interruption of a several million-year coevolution with large grazers.

During the 20th century, crops spread in the most favorable subregions and coexisted with livestock in the same farms within a rotation between sown pastures and crops (Hall et al. 1992). Native OE were restricted, in general, to habitats with limited conditions for cropping. In the late 20th century and over the course of the current century, technological and global market changes further facilitated the expansion of crops to marginal soils and the lengthening of the crop phase in the rotation. Grass-dominated ecosystems are now restricted to areas with strong limitations for cropping, and grazing by livestock in these ecosystems is so widespread that fire is also limited to particular vegetation units or conditions.

Effects of grazing and fire on the structure and functioning of the RPG

Effects on vegetation structure

Several studies in the RPG have evidenced profound consequences of grazing on vegetation structure, with the magnitude of its impact being greater as habitat productivity increases (Lezama et al. 2014) (Figure 2). Grazing induces changes in species composition, by replacing the dominant tall grasses with prostrate warm-season species (Facelli et al. 1989; Rodríguez et al. 2003; Altesor et al. 2005, 2006; Ferreira et al. 2020). This leads to a strong change in the vertical distribution of leaves, which concentrate their biomass in the first centimeters above the ground (Sala et al. 1986; Altesor et al. 2006; Ferreira et al. 2020). Grazing also exerts a positive effect on species richness (Rusch and Oesterheld

1997; Altesor et al. 2006; Lezama et al. 2014; Ferreira et al. 2020), particularly of graminoids and forbs (Lezama et al. 2014). In the Flooding Pampa, species richness increases mainly through the addition of exotic cool-season forbs (Rush and Oosterheld 1997; Facelli et al. 1989), differently from what occurs in the Uruguayan Campos, where no impact of grazing on the number of exotics has been observed, probably related to the relative importance of C4 grasses in this region (Altesor et al. 2006; Bresciano et al. 2014). Unlike richness, graminoid cover does not differ with grazing regime, while forb cover tends to be higher with grazing (Lezama et al. 2014). At larger scales, grazing reduces the spatial heterogeneity or β diversity (Chaneton and Faceli 1991; Chaneton et al. 2002; Lezama et al. 2014).

A particular pattern observed as a response to grazing in the RPG is the decrease in the richness and cover of shrubs (Lezama et al. 2014; Altesor et al. 2006; Ferreira et al. 2020). Despite the apparent homogeneity in grazing response patterns in the RPG, there are climatic, topographic and edaphic differences that interact with disturbances and lead to the emergence of different patterns. In areas with the presence of boundaries between forests and grasslands, as the Mesopotamic Pampa (Biganzoli et al. 2009; Chaneton et al. 2012), southern Brazil (Muller et al. 2012; Overbeck et al. 2016), and eastern Uruguay (Altesor et al. 2006; Gallego et al. 2020), it has been observed that grazing and/or fire suppression increases shrub and tree species cover, either native or exotic (Figure 2).

In the RPG, prescribed fires are common in areas dominated by tussock grasses. Fire spreads fueled by standing dead biomass of the tussocks and temporarily reduces its coverage and height (Latterra et al. 1998; Overbeck et al. 2005; Overbeck and Pfadenhauer 2007; Beal-Neves et al. 2020; López-Mársico et al. 2021). Soon after fire, the tussocks begin to produce fresh green leaves, which are attractive and consumed by livestock (Latterra et al. 1998; Latterra et al. 2003; López-Mársico et al. 2021). Also, fire generates bare soil spaces that are quickly occupied by a large number of species found in the regional species pool, but poorly represented before fire (Latterra et al. 1998; Overbeck et al. 2005; Fidelis et al. 2012; Beal-Neves et al. 2020; Cuello et al. 2020; López-Mársico et al. 2021), such as rosette forbs (Overbeck and Pfadenhauer 2007; Fidelis et al. 2012). Specifically in the Flooding Pampa, fire promotes the invasion of exotic species (Latterra 1997; Juan et al. 2000; Latterra et al. 2003). Accordingly, fire increases spatial heterogeneity that results in higher plant species richness (Overbeck et al. 2005; Beal-Neves et al. 2020; López-Mársico et al. 2021) or, occasionally, in equal numbers (Fidelis et al. 2012). As time since the fire increases, or when it is suppressed, standing dead biomass of the dominant tussock grasses accumulates and prevents the establishment of many less competitive herbaceous species.

Effects on fauna

It has been proposed that disturbances in grasslands, such as grazing and fire, create a shifting mosaic of patch types that differ in plant community composition and structure, translating into greater diversity at higher trophic levels (Fuhlendorf et al. 2006). Birds have been one of the most studied taxa in the RPG. Most bird species use grasslands for nesting and foraging. In addition, several species are long-distance migrants that use grasslands for wintering or breeding (Aspiroz et al. 2012). Spatial heterogeneity of vegetation enhances diversity of birds, as some species are either restricted to short (Isacch and Martínez 2003; Aldabe et al. 2019), or tall grasslands (Isacch and Cardoni 2011) while many others make broader use of both grassland types (Aspiroz et al. 2012). Accordingly, it has been proposed that heterogeneous grazing and fire management optimizes conservation of grassland birds (Develey et al. 2008; Isacch and Cardoni 2011; Jakobosky et al. 2017; Beal-Neves et al. 2020).

Conversely, it seems that disturbances negatively affect the presence of non-flying small mammals (such as rodents and marsupials), which have more restricted mobility than birds. Pedó et al. (2010) found that species richness and abundance of small mammals more than tripled inside protected areas, compared to grazed or periodically burned grasslands of southern Brazil. Similarly, Luza et al. (2018) found that three rodent species, that are among the most abundant species in the

South Brazilian grasslands, preferentially occupy paddocks subjected to moderate or low grazing intensities.

Information about the effect of disturbances on invertebrate diversity is scarce and seems to be group-specific. An increase in resource quantity (i.e. plant biomass) or quality (i.e. plant diversity) as well as a greater complexity in vegetation architecture may lead to an increase in the richness and abundance of arthropods (Evans et al. 2015). A study in southern Brazil showed that high-taxa richness and abundance of vegetation-dwelling arthropods increased under deferred grazing or grazing exclusion, while epigeic arthropod communities did not respond to grazing management (Ferreira et al. 2020). Also, it has been observed that certain communities of below-ground biota, such as mites (Altesor et al. 2006) and earthworms (Zerbino et al. 2006) show striking differences in species composition when grazed/ungrazed plots are compared. On the other hand, given the low intensity and rapid spreading of grass-fueled fires, arthropod communities would not be greatly affected by this disturbance. Although individual mortality by heating may instantly reduce populations, spider (Podgaiski et al. 2013) and grasshopper (Ferrando et al. 2016) communities recover promptly from small-scale fires through fast recolonization from unburned patches, tracking changes in vegetation. After one year since the fire, communities from burned sites were indistinguishable from unburned ones. Also, fire favors the growth and reproduction of several low competitive plant species, which in turn may also benefit species from other trophic levels, like pollinators (da Silva et al. 2021), extrafloral nectaries-visiting ants (da Silva et al. 2020) or thrips (Podgaiski et al. 2018).

Effects on primary production

More than 75% of the variation in aboveground net primary production of the herbaceous layer of OE can be accounted for by their mean annual precipitation (McNaughton 1985; Sala et al. 1988a; McNaughton et al. 1993; Paruelo et al. 1999). This pattern was also observed in the RPG region, despite the limited range of mean annual precipitation studied (Paruelo et al. 2010; Durante et al. 2017). However, at the local scale, landscape and soil characteristics modulate the vegetation response to rainfall (Sala et al. 1988; Lane et al. 1998).

On top of these abiotic factors primary production is affected by the disturbance regime. Grazing increases or decreases ANPP when compared to non-grazed nearby areas, but the relative effect was in general, no larger than 50% and showed no relationship with mean annual precipitation (Oesterheld et al. 1999). In the RPG only two studies specifically addressed the effect of grazing on ANPP, with contrasting results. In the Flooding Pampa grazing reduced ANPP (Rusch and Oesterheld 1997), while in the Campos region increased it (Altesor et al. 2005), compared to an adjacent enclosure. The intra-regional differences indicate that the effect of grazing on primary production is difficult to generalize, as it can be mediated by different components of the vegetation, like species composition/diversity or canopy structure, or the grazing regime, like stock density or defoliation frequency and intensity. Four studies addressed the effect of grazing on belowground net primary production in the RPG (Doll and Deregibus 1986; Soriano 1992; López-Mársico et al. 2015, 2016). Generally, grazing promoted an increase in root productivity, reaching a maximum in spring or summer.

The synthesis from Oesterheld et al. (1999) showed that for OE worldwide the effects of fire on productivity were relatively larger than the effects of grazing and showed a significant pattern along the precipitation gradient. Both positive and negative effects of fire on productivity appear to be more intense than the effects of grazing. The effect of fire on productivity was positively associated with mean annual precipitation, showing that the increases in productivity in burned areas are larger in the wet portion of gradient, which include a substantial portion of the Campos of the RPG. The local evidence on the effect of fire on primary production is scarce. Lateral et al. (1998) indicated that fire effects on tall grasslands of the Flooding Pampa showed no clear patterns. The effects ranged from nil to a 50% increase (Hidalgo and Lateral 1993, 1995). Prescribed fires are used as a management tool in some areas of the RPG to control shrubs or tall grasses species (Lateral et al. 1998; Overbeck et al.

2007; López-Mársico et al. 2021). A study carried out in eastern Uruguayan grasslands using NDVI (Normalized Difference Vegetation Index), a spectral index closely related to primary production, showed that, after a reduction of biomass due to fire, NDVI experience a rapid recovery (72 ± 37 days) and values of burned sites can even exceed those of adjacent non-burned sites (Bruzzone et al. 2018; Bruzzone 2019).

Effects on biogeochemical cycles

Grazing and fire both have the fundamental effect of opening biogeochemical cycles in ecosystems. Fire consumes biomass and releases large amounts of inorganic or simple organic compounds, increasing the exchange of nutrients with the atmosphere and the surrounding ecosystems (Bond and Keeley 2005). Therefore, large amounts of nutrients are usually lost from burned ecosystems, although nutrient availability also increases, potentially increasing some processes such as C gains (Harris et al. 2007; Oesterheld et al. 1999). Grazing, on the other hand, though not as fast as fire, also consumes biomass and releases inorganic nutrients and easily decomposable feces, fastening nutrient cycles (Bond and Keeley 2005). The concentration of nutrients in urine and dung patches also increases nutrient losses from ecosystems (Piñeiro et al. 2006). Large herbivores concentrate nutrients over the landscape due to their behavior patterns (affected by water availability and sleeping sites), favoring volatilization and leaching of inorganic compounds (Pastor et al. 2006; Chapin et al. 2002).

Soil organic matter (SOM) is the main reservoir of nutrients and C in most grasslands (Parton et al. 1993). SOM may store energy that equals more than 10 years of grassland's primary productivity and large stocks of nutrients and carbon. Grazing can change soil structure, function and its capacity to store SOM and, then, C stocks. Energy stored in SOM fuels soil microorganisms that make nutrients available for plants and maintains biogeochemical cycles (Chapin et al. 2002). Changes in SOM would influence soil physical properties and intermediate ES supply (nutrient retention and mineralization, water holding capacity, soil biodiversity, etc.). SOM is a key structural aspect in determining, regulating and supporting ES (Paruelo and Vallejos 2013). Grazing and fire affect SOM by affecting either carbon inputs or outputs to the soil. Such inputs and outputs are in turn affected, directly or indirectly, by changes in the nitrogen cycle (Piñeiro et al. 2009; Piñeiro et al. 2010). Nitrogen plays a central role in controlling carbon inputs and outputs from SOM because it controls simultaneously primary productivity, plant carbon allocation and SOM stabilization in the mineral associated organic matter (MAOM), the main SOM fraction in grasslands (Lavallo et al. 2020; Piñeiro et al. 2010) (Figure 3). Increasing N inputs (i.e. by overseeding legumes to increase biological N fixation) has been proposed as an alternative to increase C sequestration in grasslands soils (Figure 3). However, empirical data shows that SOM may increase or decrease with more legumes depending on grazing management (Bondaruk et al. 2020).

Both grazing and fire reduce the amount of carbon inputs to the soil, due to biomass consumption, potentially decreasing SOM stocks. Heavily and frequently burned areas and overgrazing undoubtedly decrease SOM stocks, but moderate grazing and infrequent burning have shown variable results (Abdalla et al. 2018). Several works report both increases, decreases or unchanged SOM stocks under contrasting grazing conditions and fire regimes across temperature and precipitation gradients, suggesting that grazing and fire influence the factors that control SOC accumulation in complex ways (Piñeiro et al. 2010). As shown in previous sections, grazing and fire impacts on both below and aboveground primary production by modifying C inputs (Figure 3). Grazing may increase root allocation in some environments and therefore increase SOM stocks (because of higher SOM formation efficiency of roots), particularly in the particulate organic matter fraction (POM) (Piñeiro et al. 2010). Such changes may derive from the effect of disturbance on species and plant functional type composition (Figure 3). Disturbance would also impact on soil respiration affecting directly SOC accumulation (Figure 3).

Environmental Footprint of livestock ranching in the RPG

The environmental footprint of agricultural activities is directly linked to its impact on biodiversity and ES supply. Previous sections on the effect of grazing and fire management on vegetation structure and fauna stressed the importance of disturbances on maintaining or even increasing plant and animal biodiversity. Both disturbances play, then, a critical role in determining the supply of ES.

Sala and Paruelo (1997) summarized the potential contribution of grassland areas to the supply of ES. The referred article focused on services whose value is not reflected in the market (atmospheric gases regulation, genetic library, soil conservation and climate regulation). As in a foundational article on the subject (Costanza et al. 1997), the focus was on monetizing the contribution of rangelands to the supply of these services. The economic valuation of ES has been controversial (i.e. Silvertown (2019)) and it was perceived as “commoditization” of Nature. The development of integrative conceptual frameworks that connect ecosystem structure and functioning with ecosystem services and benefits to the society (the “cascade model”, Haines-Young and Potschin 2010) helps developing alternatives to quantify the supply of ES and to integrate them into decision making processes (Paruelo and Laterra 2019; Staiano et al. 2020). The contribution of OE is particularly important because it includes both provisioning (meat, wool, water supply) and regulating services (pollination, C sequestration, hydrologic regulation) (Yahdjian et al. 2015). Moreover, Yahdjian et al. (2015) strongly emphasized the importance of considering not only the supply of ES but also the demand. Clearly, the society qualitatively modified the demand of ES in OE. Supporting and regulating ES and the conservation of natural habitats became critical in defining the environmental footprint of systems that focus on provisioning ES such as meat or wool. In fact, documenting the impact of production systems, its environmental footprint, on such ES is crucial to access to some markets (Kerhoe et al. 2020).

The previous sections documented the effect of grazing and fire, often associated to ranching activities, on several structural and functional aspects of OE. Such effects, in many cases, determine a higher supply of regulating ES compared to other land uses and even to the absence of grazing (Gallego et al. 2020). Empirical studies all over the RPG documented that grasslands, regardless of the grazing and fire regime, accumulate more soil organic C, have more biodiversity and emit less N₂O than most of the other land covers in the region (Della Chiesa et al. 2019, Mazzilli 2015). Two metrics provided synoptic estimates of the environmental footprint of agricultural activities over the RPG: the Human Appropriation of Net Primary Productivity (HANPP) (Baeza and Paruelo 2018) and the Ecosystem Service Supply Index (ESSI) (Paruelo et al. 2016).

Human Appropriation of Net Primary Productivity (HANPP)

HANPP results from the difference between the NPP in the absence of human influence (NPP of potential vegetation: NPP₀) and the NPP of the actual vegetation remaining after harvest (NPP_{REM}). NPP_{REM} was calculated as the NPP of the current vegetation (NPP_{ACT}) minus the harvested NPP (NPP_H), directly appropriated by humans as agricultural products (grain, wood, meat, etc.) or destroyed during harvest

$$\text{HANPP} = \text{NPP}_0 - \text{NPP}_{\text{REM}} = \text{NPP}_0 - (\text{NPP}_{\text{ACT}} - \text{NPP}_{\text{H}}) \quad (1)$$

The complement of HANPP quantifies the energy available for ecosystem processes which, in turn, are responsible of regulating ES supply. HANPP in the RPG ecosystems, considering all land uses, increased from 41.8% to 46.4% from 2001/2002 to 2012/2013 (Baeza and Paruelo 2018). Appropriation surpassed the 70–80% of NPP₀ in agricultural and afforested areas. Using the same approach than Baeza and Paruelo (2018) (see Supplementary Material) we calculated the HANPP for the remnant OE areas of the RPG (Figure 4). On average the HANPP on the OE of the RPG is less than 11% and it varied from 9 (West Inland Pampa) to 15% (Northern Campos of Uruguay) (Figure 4), values substantially lower than the regional average (42%).

393

394 *Ecosystem Service Supply Index (ESSI)*

395 Paruelo et al. (2016) presented a synoptic indicator of “bundles” supporting and regulating ES related
396 to carbon and water dynamics, the “Ecosystem Services Supply Index” (ESSI). It is based on
397 vegetation indices derived from remote sensing data, which constitute robust estimators of Net
398 Primary Productivity (NPP) (Monteith 1972; Piñeiro et al. 2006), an integrating variable of ecosystem
399 functioning (McNaughton et al. 1989). The ESSI merges two attributes of the NDVI annual dynamics:
400 the annual average (NDVI_{MEAN}, a proxy of total C gains) and the intra-annual coefficient of variation
401 (NDVI_{CV}, an indicator of seasonality): $ESSI = NDVI_{MEAN} * (1 - NDVI_{CV})$. Those sites where annual
402 productivity is higher and more seasonally stable would have a higher ES supply.

403 The foundation of the ESSI is based on both the conceptual framework of the ES cascade
404 model and the ES bundles concept (Raudsepp-Hearne et al. 2010). According to this scheme, the ESSI
405 represents an integrative index of ecosystem functioning which gives rise to the cascade. It can
406 describe the variation in different regulating and supporting ES (some of them intermediate and
407 others final ES) that vary together in the same direction (ES bundles). The support for using ESSI as a
408 proxy of ES supply was originally based on its positive relationship with four ES estimated from
409 empirical data or mechanistic models: groundwater recharge and avian richness in Dry Chaco forests
410 and soil organic carbon (SOC) in the RPG (Paruelo et al. 2016). Two additional studies provided
411 additional support to the use of ESSI (Weyland et al. 2019; Staiano et al. 2020). The index has been
412 used in a variety of systems to evaluate spatial and temporal patterns of the environmental footprint
413 of agricultural activities (Verón et al. 2017; Staiano et al. 2020; Gallego et al. 2020; Jullian et al. 2021;
414 Camba Sans et al. 2021).

415 The ESSI of the natural OE of the RPG was higher than the values of transformed land cover
416 (in average a 15% higher), except for tree plantations (Figure 5). Differences were more important in
417 those subregions where the level of agricultural intensification was higher (i.e. the Rolling Pampa in
418 Argentina, 22%) than in those experiencing lower levels (i.e. the Northern Campos in Uruguay, 5%).
419 The temporal trends of ecosystem supply showed even larger differences. Less than 4% of the
420 grasslands, savannas and shrublands of the RPG showed a significant reduction in ES supply for the
421 period 2001-2019 (Figure 6). For the rest of the region, almost 25% of the area experienced a loss in ES
422 supply over the last two decades. As any indicator ESSI must be used in a multidimensional
423 diagnosis. For example, higher ESSI values for tree plantations have to be considered with the level of
424 HANPP (Figure 4) or the proportion of habitat conservation (Figure 1).

425

426 *Meat equivalent production in Open Ecosystems (OE) of the RPG*

427 Oesterheld et al. (1992) showed that in livestock systems of South America, herbivore biomass
428 increased exponentially with ANPP and that herbivore biomass per unit of ANPP was 6 times higher
429 than in natural ecosystems. Over regional scales, ANPP is the major determinant of meat equivalent
430 (meat + milk + wool) production (ME), a substantial portion of the Net Secondary Production.
431 Compared to natural OE dominated by homeotherms, Trophic Efficiency (NSP/ANPP) in managed
432 systems is higher due to increases in Production Efficiency and, mainly, on Consumption Efficiency
433 (Irisarri et al. 2014). Such increases in Efficiency results from several management actions such as
434 animal health care, water supply, breeding, etc. In general livestock managed systems reach higher
435 NSP by (1) diverting a major proportion of ANPP from the detritus to the grazing chain, (2)
436 converting more efficiently consumption into NSP, and (3) stabilizing herbivore biomass across years.

437 Combining the data from Irisarri et al. (2014) and Gutierrez et al. (2020) we generated a model
438 on the relationship between ANPP and PSN for the entire region (Figure A1, Supplementary
439 Material). A particular feature of the fitted model indicates that a unit change in ANPP generates a
440 more than proportional change in ME production. Because of this pattern, the conversion of ANPP

dry matter into live weight decreased from more than 100 kg DM per kg of live weight, for sites below 2500 kg DM.ha⁻¹.yr⁻¹ of ANPP, to 80 to 60 kg of DM per kg of LW or even less than 60 kg of DM per kg of LW for sites located above 4500 kg DM.ha⁻¹.yr⁻¹ of ANPP (Figure A1, Supplementary Material).

Based on this model and the ensemble map of remnant OE in the RPG (see Figure 1), we estimated the production of ME on OE of the RPG. The average value per hectare was 78 kg ha⁻¹.yr⁻¹, with extreme values of 92.5 and 35.2 kg ha⁻¹.yr⁻¹ for the Southern Campos of Uruguay and the West Inland Pampa, respectively. Over the whole region the total volume of ME production on OE was 3488.6x10³ ton liveweight.yr⁻¹, for the ensemble map of distribution of remnant OE. These figures represent the ME production volume of the region with the lowest environmental footprint, at least based on Soil C dynamics and biodiversity conservation. The estimated values of ME production on the grassy ecosystems of the RPG represents a high proportion of the total production in the region. For the case of Uruguay the potential production of ME with minimum environmental footprint is equivalent to the total slaughtered bovines (1160 vs 1173x10³ ton liveweight.y⁻¹ for 2019 according to OPyPA (2020)). Assuming a per capita meat consumption of 6.27 kg.y⁻¹ (Consumption, 2020) and a yield of 35%, the RPG region may feed more than 195x10⁶ people, approximately a 2.5% of the world population. The people from the RPG may invite an “asado con cuero” to half the world population based on meat with the lowest possible environmental footprint.

The contribution of Open Ecosystems (OE) to local and global sustainability

Grasslands, shrublands and savannas are under threat worldwide and particularly in South America. The proportion of the remaining grassland, shrublands and savannas area under protection is the lowest among the different terrestrial ecosystems (Jones et al. 2018). In the Argentine Pampa the proportion is lower than 1% (Burkart 2005; Moreno et al. 2008). The grassland area under protection is even lower in Uruguay. Brazil has a 1% of the RPG under strict protection and a 2% more under sustainable use (Metzger et al. 2019), but Legal Reserves in private land could protect almost 300 thousand hectares of grassland remnants if the Brazilian legislation on native vegetation were enforced (Overbeck et al. 2015). Agricultural conversion during the first 14 years of XXI century expanded 23% (Baeza and Paruelo 2020), a value similar to the rate observed for subtropical forests (Volante et al. 2016). Moreover, two key components of these systems, grazing and fire, are often demonized, perhaps for its deleterious role in forested ecosystems. The evidence suggests that fire and grazing are critical to maintain the biodiversity at the species and landscape level and to increase the supply of ES. In areas where most of the land is privately owned, well managed ranching is the most affordable alternative to preserve natural habitats and to develop sustainable systems in environmental, social and economic terms. Grazing lands have the highest potential for soil C sequestration (Caride et al. 2012; Li et al. 2018; Baethgen et al. 2021), reduction of GHG emissions (Della Chiesa et al. 2019), water regulation (Nosetto et al. 2005) and biodiversity preservation (Altesor et al. 2006, Lezama et al. 2019) among the different land cover that occupied the RPG landscapes. Here we showed that those areas have a higher supply of regulating ES than those that were transformed. Moreover, the area of OE experiencing positive trends in regulating ES supply is three times higher than that showing negative trends. The higher supply of regulating ES would be directly linked to the higher availability of energy as indicated by the lower C exports from these ecosystems.

Livestock production, the main economic activity on OE, generates, though, a major concern: enteric methane emissions (Friedlingstein et al. 2019, Clark et al. 2020). This is a highly controversial issue regarding the environmental impacts of cattle and sheep ranching. The role of CH₄ as a GHG is in itself controversial (Cain et al. 2019). Its radiative forcing potential is higher than the CO₂ but its residence time is substantially lower. CO₂ accumulates in the atmosphere becoming then a “stock gas”, while 50% of the CH₄ emitted gets transformed into CO₂ in a decade. It is considered accordingly a “flux gas” (Cain et al. 2019). Moreover, enteric methane derives from anaerobic respiration of CO₂ previously removed from the atmosphere by photosynthesis. The baseline values of ruminant

emissions are also controversial. Smith et al. (2016) used a large spatially explicit database of late Quaternary mammals and body mass, and allometric regressions that relate population density, geographic range, and methane output to mass. Their calculations showed that CH₄ emissions from large mammals at the late Pleistocene would be equivalent to current domestic herbivores emissions. Human driven megafauna extinctions would be responsible for the lower CH₄ emissions during most of the Holocene. For Uruguayan livestock systems producing 136 ME kg.ha⁻¹.yr⁻¹ (in farms including natural grasslands and 50% of sowed pastures), CH₄ emissions, in CO₂ equivalent, was estimated in 421 kg of C ha⁻¹.year⁻¹ (1549 kg CO₂eq.ha⁻¹.year⁻¹) (Becoña et al. 2013). Tough it is not possible to claim for a generalized neutral C scenario (Villarino et al. 2020), field data showed that, in the RPG, grazing can increase or slightly reduce (+- 8%) the amount of C accumulated in the soil compared to not grazed situations for long periods of time (Piñeiro et al. 2009). Only under particular circumstances, the annual C gain in soils in grazed environments with respect to ungrazed ones can exceed 1 ton C/year, more than compensating CH₄ emissions. Simulation studies and long-term experiments on sowed pastures showed also a high potential of C sequestration (Baetghen et al. 2021). Terrer et al. (2021) show from a synthesis of CO₂ enrichment experiments from different ecosystems that Soil Organic Carbon (SOC) can increase by approximately 5% in response to a 65% step increase in CO₂ with a strong coupling between driven changes in aboveground plant biomass and SOC. However, the coupling between plant biomass and soils is an inverse relationship opposite to that simulated by many ecosystem models. That is, if it accumulates in plant biomass it does not accumulate in the soil. The results highlight the potential for grassland soils to store carbon as atmospheric CO₂ levels continue to rise. An estimate of this is that C accumulation should increase by about 187 kg.ha⁻¹.year⁻¹, a figure of the same order of magnitude as those from preliminary long-term studies on SOC stocks changes in Uruguay (Bazzoni, personal communication).

The main determinant of the negative environmental impact of livestock production systems is the conversion of natural habitats (Griscom et al. 2020; Cusack et al. 2021). Preserving grasslands, savannas and woodlands would warrant lower GHG emissions and biodiversity preservation. The actual opportunity to accumulate C depends on OE management (Cusack et al. 2021), being grazing management a key factor. Abdala et al. (2018) showed that moist/warm grasslands, such as those in the RPG, have the highest potential of C sequestration under grazing. For the main plant communities of the RPG the conservation status has a major effect on preserving functional, specific and structural diversity (Altersor et al. 2019). Specific management actions have been crafted to both maximize ME production and C accumulation and biodiversity preservation (Jaurena et al. 2020).

The Mercosur countries faced some serious challenges to add value (from both an ethical and economical perspective) to their agricultural products (see Kerhoe et al. 2020). Showing the connection between OE conservation and the environmental footprint of the production is critical. The importance of preserving forest ecosystems is well established and it concentrates important resources. Grassland, shrublands and savannas conservation has not been incorporated into the conservation political agenda. For example, Argentina has since 2007 a National Law to regulate deforestation (Aguilar et al. 2018) and a similar one aimed to preserve wetlands is under discussion, while in Brazil the federal Law on Native Vegetation Protection has not been enforced to protect native grassland vegetation (http://ecoqua.ecologia.ufrgs.br/arquivos/Agonia_do_Pampa.pdf). Open Ecosystems (OE) conservation requires a myriad of public policies, ranging from restrictions to incentives. The regional scientific system is able to provide policy makers not only the conceptual basis, but a number of building blocks to develop such policies, including:

- Management schemes of extensive livestock systems (Jaurena et al. 2020)
- Monitoring systems of remnant OE areas (MapBiomass Pampa)
- Grassland inventories (Lezama et al. 2019; Perelman et al. 2001; Batista et al. 2014)
- Estimates of Forage Production and Net Secondary Production of livestock systems (Grigera et al. 2007; Irisarri et al. 2014; Gutierrez et al. 2020)

- Indices and protocols for monitoring ES supply, environmental footprint and/or conservation status at the field level (Altesor et al. 2019; Blumetto et al. 2019) and regional level (Paruelo et al. 2016; Baeza and Paruelo 2018; Teixeira et al. 2019).
- Schemes of co-innovation to connect the scientific system and the stakeholders (Albicette et al. 2017; Rugia et al. 2021)

ACKNOWLEDGEMENTS

This work was funded by CONICET, FONCYT, UBA (Argentina), INIA, ANII and CSIC (Uruguay).and CNPq (grants 307689/2014-0 and 441280/2017-0).

REFERENCES

- Abdalla, M., Hastings, A., Chadwick, D. R., Jones, D. L., Evans, C. D., Jones, M. B., ... & Smith, P. 2018. Critical review of the impacts of grazing intensity on soil organic carbon storage and other soil quality indicators in extensively managed grasslands. *Agriculture, Ecosystems & Environment*, 253, 62-81.
- Aldabe, J., Lanctot, R. B., Blanco, D., Rocca, P., & Inchausti, P. 2019. Managing grasslands to maximize migratory shorebird use and livestock production. *Rangeland Ecology & Management*, 72(1), 150-159.
- Aguiar, S., Mastrangelo, M. E., Garcia Collazo, M. A., Camba Sans, G. H., Mosso, C. E., Ciuffoli, L., ... & Verón, S. R. 2018. ¿Cuál es la situación de la Ley de Bosques en la Región Chaqueña a diez años de su sanción? Revisar su pasado para discutir su futuro.
- Albicette, M. M., Leoni, C., Ruggia, A., Scarlato, S., Blumetto, O., Albín, A., & Aguerre, V. 2017. Co-innovation in family-farming livestock systems in Rocha, Uruguay: A 3-year learning process. *Outlook on agriculture*, 46(2), 92-98.
- Allen, R., Irmak, A., Trezza, R., Hendrickx, J. M., Bastiaanssen, W., & Kjaersgaard, J. 2011. Satellite-based ET estimation in agriculture using SEBAL and METRIC. *Hydrological Processes*, 25(26), 4011-4027.
- Altesor, A., Gallego, F., Ferrón, M., Pezzani, F., López-Mársico, L., Lezama, F., ... & Paruelo, J. M. 2019. Inductive approach to build state-and-transition models for Uruguayan Grasslands. *Rangeland Ecology & Management*, 72(6), 1005-1016.
- Altesor, A., Piñeiro, G., Lezama, F., Jackson, R. B., Sarasola, M., & Paruelo, J. M. 2006. Ecosystem changes associated with grazing in subhumid South American grasslands. *Journal of Vegetation science*, 17(3), 323-332.
- Altesor, A., Oesterheld, M., Leoni, E., Lezama, F., & Rodríguez, C. 2005. Effect of grazing on community structure and productivity of a Uruguayan grassland. *Plant Ecology*, 179(1), 83-91.
- Azpiroz A B, Isacch J P, Dias R A, Di Giacomo A S, Fontana C S and Palarea C M. 2012. Ecology and conservation of grassland birds in southeastern South America: a review. *J. Ornithol.* 83: 217–246.
- Baethgen, W. E., Parton, W. J., Rubio, V., Kelly, R. H., & Lutz, S. M. 2021. Ecosystem dynamics of crop-pasture rotations in a fifty-year field experiment in Southern South America: Century model and field results.
- Baeza, S., & Paruelo, J. M. 2020. Land use/land cover change (2000–2014) in the Rio de la Plata grasslands: an analysis based on MODIS NDVI time series. *Remote Sensing*, 12(3), 381.

581 Baeza, S., & Paruelo, J. M. 2018. Spatial and temporal variation of human appropriation of net primary
582 production in the Rio de la Plata grasslands. *ISPRS journal of photogrammetry and remote*
583 *sensing*, 145, 238-249.

584 Baeza, S., Baldassini, P., Bagnato, C., Pinto, P., & Paruelo, J. 2014. Caracterización del uso/cobertura
585 del suelo en Uruguay a partir de series temporales de imágenes MODIS. *Agrociencia*
586 (Uruguay), 18(2), 95-105.

587 Baldi, G., & Paruelo, J. M. 2008. Land-use and land cover dynamics in South American temperate
588 grasslands. *Ecology and Society*, 13(2).

589 Baldi, G., Guerschman, J. P., & Paruelo, J. M. 2006. Characterizing fragmentation in temperate South
590 America grasslands. *Agriculture, Ecosystems & Environment*, 116(3-4), 197-208.

591 Bastin, J. F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., ... & Crowther, T. W. 2019.
592 The global tree restoration potential. *Science*, 365(6448), 76-79.

593 Batista WB, Rolhauser AG, Biganzoli F, Burkart SE, Goveto L, Maranta A, Pignataro AG, Morandeira
594 NS, Rabadán M. 2014. Las comunidades vegetales de la sabana del Parque Nacional El Palmar
595 (Argentina). *Darwiniana* 2: 5–38

596 Beal-Neves, M., Chiarani, E., Ferreira, P. M. A., & Fontana, C. S. 2020. The role of fire disturbance on
597 habitat structure and bird communities in South Brazilian Highland Grasslands. *Scientific*
598 *reports*, 10(1), 1-14.

599 Becoña, G., Ledgard, S., Wedderburn E.. 2013. A Comparison of Greenhouse Gas Emissions from
600 Uruguayan and New Zealand Beef Systems. *Agrociencia*, 17:120-130 -

601 Behling, H., Pillar, V.D., 2007. Late Quaternary vegetation, biodiversity and fire dynamics on the
602 southern Brazilian highland and their implication for conservation and management of modern
603 Araucaria forest and grassland ecosystems. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 362, 243–51.

604 Behling, H., Pillar, V.D., Bauermann, S.G., 2005. Late Quaternary grassland (Campos), gallery forest,
605 fire and climate dynamics, studied by pollen, charcoal and multivariate analysis of the São
606 Francisco de Assis core in western Rio Grande do Sul (southern Brazil). *Rev. Palaeobot. Palynol.*
607 133, 235–248.

608 Bergamin et al. 2021. Environmental and anthropogenic drivers of native and alien grassland diversity
609 in the Rio de la Plata ecoregion. *Journal of Vegetation Science* (submitted).

610 Biganzoli, F., Wiegand, T., & Batista, W. B. 2009. Fire-mediated interactions between shrubs in a South
611 American temperate savannah. *Oikos*, 118(9), 1383-1395.

612 Blumetto, O., Castagna, A., Cardozo, G., García, F., Tiscornia, G., Ruggia, A., ... & Albin, A. 2019.
613 Ecosystem Integrity Index, an innovative environmental evaluation tool for agricultural
614 production systems. *Ecological indicators*, 101, 725-733.

615 Boldrini II. 1997. Campos do Rio Grande do Sul: caracterização fisionômica e problemática
616 ocupacional. *Boletim do Instituto de Biociências* 56: 1–39.

617 Bond, W. J., & Keeley, J. E. 2005. Fire as a global 'herbivore': the ecology and evolution of flammable
618 ecosystems. *Trends in ecology & evolution*, 20(7), 387-394.

619 Bond, W. J. 2019. Vertebrate herbivory and open ecosystems. In *Open Ecosystems* (pp. 121-140).
620 Oxford University Press.

621 Bondaruk, V., Lezama, F., Del Pino, A., & Piñeiro, G. 2020. Overseeding legumes in natural
622 grasslands: Impacts on root biomass and soil organic matter of commercial farms. *Science of The*
623 *Total Environment*, 743, 140771.

624 Brazeiro A., Betancourt A. y Haretche, F. 2020. Bosques nativos de Uruguay: distribución, diversidad
625 y propuesta de clasificación. *Plantae*, 3: 18-33.

626 .

627 Bresciano, D., Rodríguez, C., Lezama, F. & Altesor, A. 2014. Patrones de invasión de los pastizales de
628 Uruguay a escala regional. *Ecología Austral*, 24:83- 93

629 Bruzzone, L. Gallego, F. López-Mársico, L. 2018. Caracterización funcional de un pastizal ante eventos
630 de quema en Sierras del Este, Uruguay: una aproximación basada en sensores remotos. XXVIII
631 Reunión Argentina de Ecología, Mar del Plata, Argentina. Pp, 512.

632 Bruzzone, L. 2019. Evento de quemas en pastizales de Sierras del Este: una caracterización basada en
633 sensores. Tesis de grado, Facultad de Ciencias, Universidad de la República, Montevideo,
634 Uruguay. Pp 41.

635 Buchhorn, M. ; Lesiv, M. ; Tsendbazar, N. - E. ; Herold, M. ; Bertels, L. ; Smets, B. 2020. Copernicus
636 Global Land Cover Layers—Collection 2. Remote Sensing, 12Volume 108, 1044.

637 Buisson, E., Le Stradic, S., Silveira, F. A., Durigan, G., Overbeck, G. E., Fidelis, A., ... & Veldman, J. W.
638 2019. Resilience and restoration of tropical and subtropical grasslands, savannas, and grassy
639 woodlands. *Biological Reviews*, 94(2), 590-609.

640 Burkart, R. 2005. Las áreas protegidas de la Argentina. La situación ambiental argentina, 399-404.

641 Cabrera, A. L. 1976. Regiones fitogeográficas argentinas.

642 Cain, M., Hornsby, E., Lynch, J., & Allen, M. R. 2019. What do" net zero" and" climate neutrality" mean
643 for agriculture and for global warming?. In AGU Fall Meeting Abstracts (Vol. 2019, pp. PA32A-
644 03).

645 Sans, G. H. C., Verón, S. R., & Paruelo, J. M. 2021. Forest strips increase connectivity and modify
646 forests' functioning in a deforestation hotspot. *Journal of Environmental Management*, 290,
647 112606.

648 Caride, C., Pineiro, G., & Paruelo, J. M. 2012. How does agricultural management modify ecosystem
649 services in the argentine Pampas? The effects on soil C dynamics. *Agriculture, ecosystems &
650 environment*, 154, 23-33.

651 Chaneton, E. J., & Facelli, J. M. 1991. Disturbance effects on plant community diversity: spatial scales
652 and dominance hierarchies. *Vegetatio*, 93(2), 143-155.

653 Chaneton, E. J., Perelman, S. B., Omacini, M., & León, R. J. 2002. Grazing, environmental
654 heterogeneity, and alien plant invasions in temperate Pampa grasslands. *Biological
655 Invasions*, 4(1), 7-24.

656 Chaneton, E. J., Mazía, N., Batista, W. B., Rolhauser, A. G., & Ghera, C. M. 2012. Woody plant
657 invasions in Pampa grasslands: a biogeographical and community assembly perspective.
658 In *Ecotones between forest and grassland* (pp. 115-144). Springer, New York, NY.

659 Chapin, F. S., Matson, P. A., Mooney, H. A., & Vitousek, P. M. 2002. Principles of terrestrial ecosystem
660 ecology.

661 Chapin III, F. S., Power, M. E., Pickett, S. T., Freitag, A., Reynolds, J. A., Jackson, R. B., ... & Bartuska,
662 A. 2011. Earth Stewardship: science for action to sustain the human-earth system. *Ecosphere*, 2(8),
663 1-20.

664 Clark, M., Domingo, N., Colgan, K., Thakrar, S., Tilman, D., Lynch, J., & Hill, J. 2020. Global food
665 system emissions could preclude achieving the 1.5° and 2° C climate change targets. *Science*,
666 370(6517), 705–708.

667 Consumption, O. M. 2020. OECD: Paris.

668 Costanza R., D'Arge R., de Groot R., Farber S., Grasso M., Hannon B., Limburg K., Naeem S, O'Neill
669 R.W., Paruelo J.M., Raskin R.J., Sutton P. y Van den Belt M. 1997. The value of the world's
670 ecosystem services and natural capital. *Nature*, 357:253-260.

671 Cuello, N., López-Mársico, L., & Rodríguez, C. 2020. Field burn versus fire-related cues: germination
672 from the soil seed bank of a South American temperate grassland. *Seed Science Research*, 30(3),
673 206-214.

674 Cusack, D. F., Kazanski, C. E., Hedgpeth, A., Chow, K., Cordeiro, A. L., Karpman, J., & Ryals, R. 2021.
675 Reducing climate impacts of beef production: A synthesis of life cycle assessments across
676 management systems and global regions. *Global Change Biology*, 27(9), 1721-1736.

677 da Silva, C. V. C., da Silva Goldas, C., Dattilo, W., Dröse, W., de Souza Mendonça Jr, M., & Podgaiski,
678 L. R. 2020. Effects of time-since-fire on ant-plant interactions in southern Brazilian
679 grasslands. *Ecological Indicators*, 112, 106094.

680 da Silva Goldas, C., Podgaiski, L. R., da Silva, C. V. C. & de Souza Mendonça Jr, M. 2021. Burning for
681 grassland pollination: Recently burned patches promote plant flowering and insect pollinators.
682 *Austral Ecology*. doi:10.1111/aec.13108

683 Daubenmire, R. 1968. Ecology of fire in grasslands. *Advances in ecological research*, 5, 209-266.

684 Della Chiesa, T., Piñeiro, G., & Yahdjian, L. 2019. Gross, background, and net anthropogenic soil
685 nitrous oxide emissions from soybean, corn, and wheat croplands. *Journal of environmental*
686 *quality*, 48(1), 16-23.

687 Develey, P. F., Setubal, R. B., Dias, R. A., & Bencke, G. A. 2008. Conservação das aves e da
688 biodiversidade no bioma Pampa aliada a sistemas de produção animal. *Revista Brasileira de*
689 *Ornitologia*, 16(4), 308-315.

690 Díaz, S., Lavorel, S., McIntyre, S., Falczuk, V., Casanoves, F., Milchunas, D.G., Skarpe, C., Rusch, G.,
691 Sternberg, M., Noy-Meir, I., Landsberg, J., Zhang, W., Clark, H., Campbell, B.D. 2007. Plant trait
692 responses to grazing? a global synthesis. *Glob. Chang. Biol.* 13, 313–341.

693 Doll, U. M., & Deregibus, V. A. 1986. Efecto de la exclusión del pastoreo sobre el subsistema
694 subterráneo de un pastizal templado húmedo. *Turrialba*, 36(3), 337-344.

695 Domingo, L., Tomassini, R. L., Montalvo, C. I., Sanz-Pérez, D., & Alberdi, M. T. 2020. The Great
696 American Biotic Interchange revisited: a new perspective from the stable isotope record of
697 Argentine Pampas fossil mammals. *Scientific reports*, 10(1), 1-10.

698 Durante, M., Piñeiro, G., Irisarri, J. G. N., & Oesterheld, M. 2017. Primary production of lowland
699 natural grasslands and upland sown pastures across a narrow climatic
700 gradient. *Ecosystems*, 20(3), 543-552.

701 Evans DM, Villar N, Littlewood NA, Pakeman RJ, Evans SA, Dennis P, Skartveit J and Redpath SM .
702 2015. The cascading impacts of livestock grazing in upland ecosystems: a 10-year experiment.
703 *Ecosphere* 6: 1–15.

704 Facelli, J. M., Leon, R. J., & Deregibus, V. A. 1989. Community structure in grazed and ungrazed
705 grassland sites in the Flooding Pampa, Argentina. *American Midland Naturalist*, 125-133.

706 Ferrando, C. P. R., Podgaiski, L. R., Costa, M. K. M., & Mendonca, M. D. S. 2016. Taxonomic and
707 functional resilience of grasshoppers (Orthoptera, Caelifera) to fire in south Brazilian
708 grasslands. *Neotropical entomology*, 45(4), 374-381.

709 Ferreira, P. M., Andrade, B. O., Podgaiski, L. R., Dias, A. C., Pillar, V. D., Overbeck, G. E., ... &
710 Boldrini, I. I. 2020. Long-term ecological research in southern Brazil grasslands: Effects of grazing
711 exclusion and deferred grazing on plant and arthropod communities. *PloS one*, 15(1), e0227706.

712 Fidelis, A., Blanco, C. C., Müller, S. C., Pillar, V. D., & Pfadenhauer, J. 2012. Short-term changes caused
713 by fire and mowing in Brazilian Campos grasslands with different long-term fire
714 histories. *Journal of Vegetation Science*, 23(3), 552-562.

715 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W.,
716 Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B.,
717 Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., ... Zaehle, S. 2019. Global carbon
718 budget 2019. *Earth System Science Data*, 11(4), 1783-1838.

719 Fuhlendorf, S. D., Harrell, W. C., Engle, D. M., Hamilton, R. G., Davis, C. A., & Leslie Jr, D. M. 2006.
720 Should heterogeneity be the basis for conservation? Grassland bird response to fire and
721 grazing. *Ecological applications*, 16(5), 1706-1716.

722 Gallego, F., Paruelo, J. M., Baeza, S., & Altesor, A. 2020. Distinct ecosystem types respond
723 differentially to grazing exclosure. *Austral Ecology*, 45(5), 548-556.

724 Garibaldi, L. A., Andersson, G., Fernández Ferrari, C., & Pérez-Méndez, N. 2018. Seguridad
725 alimentaria, medio ambiente y nuestros hábitos de consumo. *Ecología austral*, 28(3), 572-580.

726 Gibson, D. J. 2009. Grasses and grassland ecology. Oxford University Press.

727 Grigera, G., Oesterheld, M., & Pacín, F. 2007. Monitoring forage production for farmers' decision
728 making. *Agricultural Systems*, 94(3), 637-648.

729 Griscom, B. W., Busch, J., Cook-Patton, S. C., Ellis, P. W., Funk, J., Leavitt, S. M., ... & Worthington, T.
730 2020. National mitigation potential from natural climate solutions in the tropics. *Philosophical*
731 *Transactions of the Royal Society B*, 375(1794), 20190126.

732 Guerschman, J. P., Paruelo, J. M., & Burke, I. C. 2003. Land use impacts on the normalized difference
733 vegetation index in temperate Argentina. *Ecological applications*, 13(3), 616-628.

734 Gutiérrez, F., Gallego, F., Paruelo, J. M., & Rodríguez, C. 2020. Damping and lag effects of
735 precipitation variability across trophic levels in Uruguayan rangelands. *Agricultural*
736 *Systems*, 185, 102956.

737 Haines-Young, R., & Potschin, M. 2010. The links between biodiversity, ecosystem services and
738 human well-being. *Ecosystem Ecology: a new synthesis*, 1, 110-139.

739 Hall, A. J., Rebella, C. M., Ghersa, C. M., & Culot, J. P. 1992. Field-crop systems of the
740 Pampas. *Ecosystems of the World*, 18, 413-450.

741 Harris, W. N., Moretto, A. S., Distel, R. A., Boutton, T. W., & Boo, R. M. 2007. Fire and grazing in
742 grasslands of the Argentine Caldenal: effects on plant and soil carbon and nitrogen. *acta*
743 *oecologica*, 32(2), 207-214.

744 Hidalgo, L. & Laterra, P. 1993. Productividad primaria de un pajonal de paja colorada (*Paspalum*
745 *quadrifarium*). Pp. 138. In: Kunst, C; Sipowicz, A; Maceira, N & Bravo de Mitre, S (Eds). *Ecología*
746 *y manejo de fuego en ecosistemas naturales y modificados*. EEA Santiago del Estero, INTA.

747 Hidalgo, L. & Laterra, P. 1995. Efecto del fuego sobre la productividad primaria de un pajonal de paja
748 colorada (*Paspalum quadrifarium*). XVII Reunión Argentina de Ecología, Mar del Plata.

749 Hobbs, N. T., Schimel, D. S., Owensby, C. E., & Ojima, D. S. 1991. Fire and grazing in the tallgrass
750 prairie: contingent effects on nitrogen budgets. *Ecology*, 72(4), 1374-1382.

751 Hoekstra, J. M., Boucher, T. M., Ricketts, T. H., & Roberts, C. 2005. Confronting a biome crisis: global
752 disparities of habitat loss and protection. *Ecology letters*, 8(1), 23-29.

753 Hulbert, L. C. 1988. Causes of fire effects in tallgrass prairie. *Ecology*, 69(1), 46-58.

754 Iriarte, J., 2006. Vegetation and climate change since 14,810 14C yr B.P. in southeastern Uruguay and
755 implications for the rise of early Formative societies. *Quat. Res.* 65, 20-32.

756 Iriondo, M., 1999. Climatic changes in the South American plains: Records of a continent-scale
757 oscillation. *Quat. Int.* 57-58, 93-112. [https://doi.org/10.1016/S1040-6182\(98\)00053-6](https://doi.org/10.1016/S1040-6182(98)00053-6)

758 Irisarri, J. G. N., Oesterheld, M., Golluscio, R. A., & Paruelo, J. M. 2014. Effects of animal husbandry on
759 secondary production and trophic efficiency at a regional scale. *Ecosystems*, 17(4), 738-749.

760 Isacch, J. P., & Martínez, M. M. 2003. Habitat use by non-breeding shorebirds in flooding pampas
761 grasslands of Argentina. *Waterbirds*, 26(4), 494-500.

762 Isacch, J. P., & Cardoni, D. A. 2011. Different grazing strategies are necessary to conserve endangered
763 grassland birds in short and tall salty grasslands of the flooding Pampas. *The Condor*, 113(4), 724-
764 734.

765 Jacobosky LI, Paulsen RK and Hartz SM. 2017. Bird-grassland associations in protected and non-
766 protected areas in southern Brazil. *Perspectives in Ecology and Conservation* 15: 109-114.

767 Jaurena, M., Durante, M., Devincenzi, T., Savian, J. V., Bendersky, D., Moojen, F. G., ... & Lattanzi, F.
768 A. 2021. Native Grasslands at the Core: A New Paradigm of Intensification for the Campos of
769 Southern South America to Increase Economic and Environmental Sustainability. *Frontiers in*
770 *Sustainable Food Systems*, 5, 11.

771 Jones, K. R., Venter, O., Fuller, R. A., Allan, J. R., Maxwell, S. L., Negret, P. J., & Watson, J. E. 2018.
772 One-third of global protected land is under intense human pressure. *Science*, 360(6390), 788-791.

773 Juan, V. F., Monterroso, L., Sacido, M. B., & Cauhépé, M. A. 2000. Postburning legume seeding in the
774 Flooding Pampas, Argentina.

775 Jullian, C., Nahuelhual, L., & Littera, P. 2021. The Ecosystem Service Provision Index as a generic
776 indicator of ecosystem service supply for monitoring conservation targets. *Ecological*
777 *Indicators*, 129, 107855.

778 Kehoe, L., dos Reis, T. N., Meyfroidt, P., Bager, S., Seppelt, R., Kuemmerle, T., ... & Kiesecker, J. 2020.
779 Inclusion, transparency, and enforcement: how the EU-mercosur trade agreement fails the
780 sustainability test. *One Earth*, 3(3), 268-272.

781 Lane, D. R., Coffin, D. P., & Lauenroth, W. K. 1998. Effects of soil texture and precipitation on
782 above-ground net primary productivity and vegetation structure across the Central Grassland
783 region of the United States. *Journal of Vegetation Science*, 9(2), 239-250.

784 Littera, P. 1997. Post-burn recovery in the Flooding Pampa: impact of an invasive legume. *Rangeland*
785 *Ecology & Management/Journal of Range Management Archives*, 50(3), 274-277.

786 Littera, P., Vignolio, O. R., Hidalgo, L. G., Fernández, O. N., Cauhépé, M. A., & Maceira, N. O. 1998.
787 Dinámica de pajonales de paja colorada (*Paspalum* Spp) manejados con fuego y pastoreo en la
788 Pampa deprimida Argentina.

789 Littera, P., Vignolio, O. R., Linares, M. P., Giaquinta, A., & Maceira, N. 2003. Cumulative effects of fire
790 on a tussock pampa grassland. *Journal of vegetation Science*, 14(1), 43-54.

791 Lauenroth, W. K., & Sala, O. E. 1992. Long-term forage production of North American shortgrass
792 steppe. *Ecological applications*, 2(4), 397-403.

793 Lauenroth, W. K. 1979. Grassland primary production: North American grasslands in perspective.
794 In *Perspectives in grassland ecology* (pp. 3-24). Springer, New York, NY.

795 Lavalley, J. M., Soong, J. L., & Cotrufo, M. F. 2020. Conceptualizing soil organic matter into particulate
796 and mineral-associated forms to address global change in the 21st century. *Global Change*
797 *Biology*, 26(1), 261-273.

798 León, R.J.C, Burkart, S., & Movia, C.P. 1979. La Vegetación de la República Argentina. Relevamiento
799 fitosociológico del pastizal del Norte de la Depresión del Salado. Instituto Nacional de Tecnología
800 Agropecuaria. Serie Fitogeográfica 17:11-93.

801 León RJC, Rush GM, Oesterheld M. 1984. Pastizales pampeanos, impacto agropecuario.
802 *Phytocoenologia* 12:201–218

803 Lewis, J. P., & Collantes, M. B. 1973. El espinal periestépico. *Ciencia e investigación*, 29(11-12), 360-377.

804 Lezama, F., Baeza, S., Altesor, A., Cesa, A., Chaneton, E. J., & Paruelo, J. M. 2014. Variation of
805 grazing-induced vegetation changes across a large-scale productivity gradient. *Journal of*
806 *Vegetation Science*, 25(1), 8-21.

807 Lezama, F, Pereira, M, Altesor, A and Paruelo, JM. 2019. Grasslands of Uruguay: a floristic based
808 description of their heterogeneity. *Phytocoenologia* 49, 211–229

809 Li, G., Zhang, Z., Shi, L., Zhou, Y., Yang, M., Cao, J., ... & Lei, G. 2018. Effects of different grazing
810 intensities on soil C, N, and P in an alpine meadow on the Qinghai—Tibetan Plateau,
811 China. *International journal of environmental research and public health*, 15(11), 2584.

812 López-Mársico, L., Lezama, F., & Altesor, A. 2021. Heterogeneity decreases as time since fire increases
813 in a South American grassland. *Applied Vegetation Science*, 24(1), e12521.

814 López-Mársico, L., Altesor, A., Oyarzabal, M., Baldassini, P., & Paruelo, J. M. 2015. Grazing increases
815 below-ground biomass and net primary production in a temperate grassland. *Plant and*
816 *Soil*, 392(1), 155-162.

817 López-Mársico, L., Oyarzabal, M., Altesor, A., & Paruelo, J. M. 2016. Effect of grazing on below-
818 ground net primary productivity of a temperate grassland. *Ecología Austral*, 26(03), 256-263.

819 Luza, A. L., Trindade, J. P. P., Maestri, R., da Silva Duarte, L., & Hartz, S. M. 2018. Rodent occupancy
820 in grassland paddocks subjected to different grazing intensities in South Brazil. *Perspectives in*
821 *Ecology and Conservation*, 16(3), 151-157.

822 MacFadden, B. J. 2005. Diet and habitat of toxodont megaherbivores (Mammalia, Notoungulata) from
823 the late Quaternary of South and Central America. *Quaternary Research*, 64(2), 113-124.

824 Mazzilli, S.. 2015. Dinámica del carbono en sistemas agrícolas bajo siembra directa. nuevas evidencias
825 obtenidas mediante el uso de ^{13}C sobre la importancia de las raíces, la calidad de los residuos y
826 el laboreo". PhD Thesis. FAUBA, EPG. 152 p.

827 McNaughton, S. J. 1985. Ecology of a grazing ecosystem: the Serengeti. *Ecological monographs*, 55(3),
828 259-294.

829 McNaughton, S. J., Oesterheld, M., Frank, D. A., & Williams, K. J. 1989. Ecosystem-level patterns of
830 primary productivity and herbivory in terrestrial habitats. *Nature*, 341(6238), 142-144.

831 McNaughton, S.J., Sala, O.E. & Oesterheld, M. 1993. Compa-rative ecology of African and South
832 American arid tosubhumid ecosystems. In: Goldblatt, P. (ed.) *Biologicalrelationships between*
833 *Africa and South America*, pp. 548-567. Yale University Press, New Haven, NY.

834 Metzger, J.P., Bustamante, M.M.C., Ferreira, J., Fernandes, G.W., Librán-Embid, F., Pillar, V.D., Prist,
835 P.R., Rodrigues, R.R., Vieira, I.C.G., Overbeck, G.E., 2019. Why Brazil needs its Legal Reserves.
836 *Perspect. Ecol. Conserv.* 17, 91–103.

- 837 Mikola, J., Setälä, H., Virkajärvi, P., Saarijärvi, K., Ilmarinen, K., Voigt, W., & Vestberg, M. 2009.
838 Defoliation and patchy nutrient return drive grazing effects on plant and soil properties in a
839 dairy cow pasture. *Ecological Monographs*, 79(2), 221-244.
- 840 Milchunas, D. G., & Lauenroth, W. K. 1993. Quantitative effects of grazing on vegetation and soils
841 over a global range of environments: Ecological Archives M063-001. *Ecological*
842 *monographs*, 63(4), 327-366.
- 843 Milkovic, M.; Gonzalez Lanzillotta, M.S.; Eljall, A.; Baldassini, P.; Perucca, R.; Insaurralde, A.; Zurano,
844 J. P.; Minoli, I.; Leszczuk, A.; Orona, M.; Loran, D.; Gill, A.; Garay, A.; Serafini, J. y Insfrán, P.
845 2021. MapBiomás Bosque Atlántico: dinámica del uso y cobertura en Argentina y Paraguay. XXIX
846 Reunión Argentina de Ecología, Tucumán, Agosto 2021.
- 847 Monteith, J. L. 1972. Solar radiation and productivity in tropical ecosystems. *Journal of applied*
848 *ecology*, 9(3), 747-766.
- 849 Morello, J., Matteucci, S. D., Rodriguez, A. F., Silva, M. E., Mesopotámica, P., & Llana, P. 2012.
850 Ecorregiones y complejos Ecosistémicos de Argentina. Orientación Gráfica Editora, Buenos Aires.
- 851 Moreno, D., Carminati, A., Machain, N., & Roldán, M. (2008). Reseña sobre las reservas privadas en la
852 Argentina. *voluntad*, 7.
- 853 Morosi, E., & Ubilla, M. 2019. Feeding and environmental studies on late Pleistocene horses in mid-
854 latitudes of South America (northern Uruguay). *Quaternary Science Reviews*, 225, 106025.
- 855 Müller, S. C., Overbeck, G. E., Blanco, C. C., de Oliveira, J. M., & Pillar, V. D. 2012. South Brazilian
856 forest-grassland ecotones: dynamics affected by climate, disturbance, and woody species traits.
857 In *Ecotones between forest and grassland* (pp. 167-187). Springer, New York, NY.
- 858 Nosetto, M. D., Jobbágy, E. G., & Paruelo, J. M. 2005. Land-use change and water losses: the case of
859 grassland afforestation across a soil textural gradient in central Argentina. *Global Change*
860 *Biology*, 11(7), 1101-1117.
- 861 Oosterheld, M., Sala, O. E., & McNaughton, S. J. 1992. Effect of animal husbandry on herbivore-
862 carrying capacity at a regional scale. *Nature*, 356(6366), 234-236.
- 863 Oosterheld, M., Loreti, J., Semmartin, M., Paruelo, J.M., 1999. Grazing, fire, and climate effects on
864 primary productivity of grasslands and savannas. *Ecosyst. world* 287-306.
- 865 Omena, É. C., Silva, J. L. L. D., Sial, A. N., Cherkinsky, A., & Dantas, M. A. T. 2020. Late Pleistocene
866 meso-megaherbivores from Brazilian Intertropical Region: isotopic diet ($\delta^{13}\text{C}$), niche
867 differentiation, guilds and paleoenvironmental reconstruction ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$). *Historical Biology*,
868 1-6.
- 869 Oppenlander, R. 2013. Food choice and sustainability: Why buying local, eating less meat, and taking
870 baby steps won't work. Hillcrest Publishing Group.
- 871 OPYPA. 2020. Análisis sectorial y cadenas productivas Temas de política Estudios. MINISTERIO DE
872 GANADERÍA, AGRICULTURA Y PESCA. Uruguay. 668 pp
- 873 Overbeck, G. E., & Pfadenhauer, J. 2007. Adaptive strategies in burned subtropical grassland in
874 southern Brazil. *Flora-Morphology, Distribution, Functional Ecology of Plants*, 202(1), 27-49.
- 875 Overbeck, G. E., Müller, S. C., Pillar, V. D., & Pfadenhauer, J. 2005. Fine-scale post-fire dynamics in
876 southern Brazilian subtropical grassland. *Journal of Vegetation science*, 16(6), 655-664.
- 877 Overbeck, G. E., Müller, S. C., Fidelis, A., Pfadenhauer, J., Pillar, V. D., Blanco, C. C., ... & Forneck, E.
878 D. 2007. Brazil's neglected biome: the South Brazilian Campos. *Perspectives in Plant Ecology,*
879 *Evolution and Systematics*, 9(2), 101-116.

880 Overbeck, G. E., Vélez-Martin, E., Scarano, F. R., Lewinsohn, T. M., Fonseca, C. R., Meyer, S. T., ... &
881 Pillar, V. D. 2015. Conservation in Brazil needs to include non-forest ecosystems. *Diversity and*
882 *distributions*, 21(12), 1455-1460.

883 Overbeck, G. E., de Abreu Ferreira, P. M., & Pillar, V. D. 2016. Conservation of mosaics calls for a
884 perspective that considers all types of mosaic-patches. Reply to Luza et al. *Natureza &*
885 *Conservação*.

886 Oyarzabal, M., Clavijo, J. R., Oakley, L. J., Biganzoli, F., Tognetti, P. M., Barberis, I. M., ... & Leon, R. J.
887 C. 2018. Unidades de vegetación de la Argentina.

888 Oyarzabal, M., Andrade, B., Pillar, V.D., Paruelo, J., 2020. Temperate Subhumid Grasslands of
889 Southern South America. In: Goldstein, M.I., DellaSala, D.A. (Eds.), *Encyclopedia of the World's*
890 *Biomes*, vol. 3. Elsevier, pp. 577–593.

891 Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., ... &
892 Kinyamario, J. I. 1993. Observations and modeling of biomass and soil organic matter dynamics
893 for the grassland biome worldwide. *Global biogeochemical cycles*, 7(4), 785-809.

894 Paruelo, J., & Laterra, P. 2019. El lugar de la naturaleza en la toma de decisiones. Editorial CICCUS.
895 Buenos Aires. Pp 567rial.

896 Paruelo, J. M., & Lauenroth, W. K. 1998. Interannual variability of NDVI and its relationship to climate
897 for North American shrublands and grasslands. *Journal of Biogeography*, 25(4), 721-733.

898 Paruelo, J. M., & Lauenroth, W. K. 1996. Relative abundance of plant functional types in grasslands
899 and shrublands of North America. *Ecological applications*, 6(4), 1212-1224.

900 Paruelo, J. M., & Vallejos, M. 2013. Ecosystem services related to carbon dynamics: its evaluation using
901 remote sensing techniques. *Earth observation of ecosystem services*. CRC Press Group, Boca
902 Raton, FL, USA, 17-32.

903 Paruelo, J. M., Jobbágy, E. G., Sala, O. E., Lauenroth, W. K., & Burke, I. C. 1998. Functional and
904 structural convergence of temperate grassland and shrubland ecosystems. *Ecological*
905 *Applications*, 8(1), 194-206.

906 Paruelo, J. M., Lauenroth, W. K., Burke, I. C., & Sala, O. E. 1999. Grassland precipitation-use efficiency
907 varies across a resource gradient. *Ecosystems*, 2(1), 64-68.

908 Paruelo, J. M., Guerschman, J. P., & Verón, S. R. 2005. Expansión agrícola y cambios en el uso del
909 suelo. *Ciencia hoy*, 15(87), 14-23.

910 Paruelo, J. M., Jobbágy, E. G., Oesterheld, M., Golluscio, R. A., & Aguiar, M. R. 2007. The grasslands
911 and steppes of Patagonia and the Rio de la Plata plains. *The physical geography of South*
912 *America*, 232-248.

913 Paruelo, J. M., Pineiro, G., Baldi, G., Baeza, S., Lezama, F., Altesor, A., & Oesterheld, M. 2010. Carbon
914 stocks and fluxes in rangelands of the Rio de la Plata basin. *Rangeland Ecology &*
915 *Management*, 63(1), 94-108.

916 Paruelo, J. M., Texeira, M., Staiano, L., Mastrángelo, M., Amdan, L., & Gallego, F. 2016. An integrative
917 index of Ecosystem Services provision based on remotely sensed data. *Ecological Indicators*, 71,
918 145-154.

919 Pastor, J., Cohen, Y. O. S. E. F., & Hobbs, N. T. 2006. The roles of large herbivores in ecosystem
920 nutrient cycles. *CONSERVATION BIOLOGY SERIES-CAMBRIDGE-*, 11, 289.

921 Pedó, E., de Freitas, T. R., & Hartz, S. M. 2010. The influence of fire and livestock grazing on the
922 assemblage of non-flying small mammals in grassland-Araucaria Forest ecotones, southern
923 Brazil. *Zoologia (Curitiba)*, 27, 533-540.

924 Perelman SB, León RJ, Oesterheld M. 2001. Cross-scale vegetation patterns of Flooding Pampa
925 grasslands. *Journal of Ecology* 89: 562–77.

926 Perelman, S.B., Silvia E. Burkart, Oyarzabal, M., Bagnato, C. & Batista, W. 2017. Climatic and land-use
927 drivers along a latitudinal gradient: species diversity in temperate grasslands on agricultural
928 soils. *Journal of Vegetation Science* 28, 1230–1239.

929 Piñeiro, G., Paruelo, J.M.J.J.M., Jobbágy, E.E.G., Jackson, R.B., Oesterheld, M. 2009. Grazing effects on
930 belowground C and N stocks along a network of cattle exclosures in temperate and subtropical
931 grasslands of South America.

932 Piñeiro, G., Paruelo, J. M., & Oesterheld, M. 2006. Potential long-term impacts of livestock
933 introduction on carbon and nitrogen cycling in grasslands of Southern South America. *Global*
934 *Change Biology*, 12(7), 1267-1284.

935 Piñeiro, G., Paruelo, J. M., Oesterheld, M., & Jobbágy, E. G. 2010. Pathways of grazing effects on soil
936 organic carbon and nitrogen. *Rangeland Ecology & Management*, 63(1), 109-119.

937 Piñeiro, G., Paruelo, J. M., Oesterheld, M., & Jobbágy, E. G. 2010. Pathways of grazing effects on soil
938 organic carbon and nitrogen. *Rangeland Ecology & Management*, 63(1), 109-119.

939 Podgaiski, L. R., Joner, F., Lavorel, S., Moretti, M., Ibanez, S., Mendonça Jr, M. D. S., & Pillar, V. D.
940 2013. Spider trait assembly patterns and resilience under fire-induced vegetation change in South
941 Brazilian grasslands. *PloS one*, 8(3), e60207.

942 Podgaiski, L. R., Cavalleri, A., Ferrando, C. P. R., Pillar, V. D., & Mendonça Jr, M. D. S. 2018.
943 Prescribed patch burnings increase thrips species richness and body size in grassland
944 communities. *Insect Conservation and Diversity*, 11(2), 204-212.

945 Prado J, Martínez-Maza C, Alberdi MT. 2015. Megafauna extinction in South America: a new
946 chronology from the Argentine pampas. *Palaeogeogr Palaeoclimatol Palaeoecol* 425:41–49

947 Prates, L., & Perez, S. I. 2021. Late Pleistocene South American megafaunal extinctions associated with
948 rise of Fishtail points and human population. *Nature communications*, 12(1), 1-11.

949 Quattrocchio, M.E., Borromei, A.M., Deschamps, C.M., Grill, S.C., Zavala, C.A. 2008. Landscape
950 evolution and climate changes in the Late Pleistocene–Holocene, southern Pampa (Argentina):
951 Evidence from palynology, mammals and sedimentology. *Quat. Int.* 181, 123–138.

952 Raudsepp-Hearne, C., Peterson, G. D., & Bennett, E. M. 2010. Ecosystem service bundles for analyzing
953 tradeoffs in diverse landscapes. *Proceedings of the National Academy of Sciences*, 107(11), 5242-
954 5247.

955 Rodríguez, C., Leoni, E., Lezama, F., & Altesor, A. 2003. Temporal trends in species composition and
956 plant traits in natural grasslands of Uruguay. *Journal of Vegetation Science*, 14(3), 433-440.

957 Ruggia, A., Dogliotti, S., Aguerre, V., Albicette, M. M., Albin, A., Blumetto, O., ... & Rossing, W. A. H.
958 2021. The application of ecologically intensive principles to the systemic redesign of livestock
959 farms on native grasslands: A case of co-innovation in Rocha, Uruguay. *Agricultural*
960 *Systems*, 191, 103148.

961 Rusch, G. M., & Oesterheld, M. 1997. Relationship between productivity, and species and functional
962 group diversity in grazed and non-grazed Pampas grassland. *Oikos*, 519-526.

963 Sala, O. E., & Paruelo, J. M. 1997. Ecosystem services in grasslands. *Nature's services: Societal*
964 *dependence on natural ecosystems*, 237-251.

965 Sala, O. E., Oesterheld, M., León, R. J. C., & Soriano, A. N. D. A. 1986. Grazing effects upon plant
966 community structure in subhumid grasslands of Argentina. *Vegetatio*, 67(1), 27-32.

967 Sala, O. E., Parton, W. J., Joyce, L. A., & Lauenroth, W. K. 1988. Primary production of the central
968 grassland region of the United States. *Ecology*, 69(1), 40-45.

969 Sala, O.E., Lauenroth, W.K., McNaughton, S.J., Rusch, G. & Xinshi Zhang, A. 1996. Biodiversity and
970 ecosystem functioning in grasslands. In: Mooney, H.A., Cushman, J.H., Medina, E., Sala, O.E. &
971 Schulze, E.D. (eds.) *Functional roles of biodiversity: A global perspective*, pp. 129-149. Wiley,
972 Chichester, UK.

973 Silvertown, J. 2015. Have ecosystem services been oversold? *Trends in ecology & evolution*, 30(11),
974 641-648.

975 Smith, F. A., Tomé, C. P., Elliott Smith, E. A., Lyons, S. K., Newsome, S. D., & Stafford, T. W. 2016.
976 Unraveling the consequences of the terminal Pleistocene megafauna extinction on mammal
977 community assembly. *Ecography*, 39(2), 223-239.

978 Soriano, A. 1992. Rio de la Plata grasslands, p. 367-407. In: R.T.Coupland (ed), *Natural grasslands:*
979 *introduction and western hemisphere. Ecosystems of the World 8A*. Elsevier, Amsterdam.

980 Souza, C. M., Z Shimbo, J., Rosa, M. R., Parente, L. L., A Alencar, A., Rudorff, B. F., ... & Azevedo, T.
981 (2020). Reconstructing three decades of land use and land cover changes in brazilian biomes with
982 landsat archive and earth engine. *Remote Sensing*, 12(17), 2735.

983 Staiano, L., Sans, G. H. C., Baldassini, P., Gallego, F., Texeira, M. A., & Paruelo, J. M. 2021. Putting the
984 Ecosystem Services idea at work: Applications on impact assessment and territorial
985 planning. *Environmental Development*, 38, 100570.

986 Strömberg, C.A.E. 2011. Evolution of Grasses and Grassland Ecosystems. *Annu. Rev. Earth Planet. Sci.*
987 39, 517-544.

988 Strömberg, C.A.E., Dunn, R.E., Madden, R.H., Kohn, M.J., Carlini, A.A. 2013. Decoupling the spread of
989 grasslands from the evolution of grazer-type herbivores in South America. *Nat. Commun.* 4,

990 Taboada, M. A., & Lavado, R. S. 1988. Grazing effects of the bulk density in a Natraquoll of the
991 flooding pampa of Argentina. *Rangeland Ecology & Management/Journal of Range Management*
992 *Archives*, 41(6), 500-503.

993 Terrer, C., Phillips, R. P., Hungate, B. A., Rosende, J., Pett-Ridge, J., Craig, M. E., ... & Jackson, R. B.
994 2021. A trade-off between plant and soil carbon storage under elevated CO₂. *Nature*, 591(7851),
995 599-603.

996 Texeira, M., Veron, S., Irisarri, G., Oyarzabal, M., Staiano, L., Baeza, S., & Paruelo, J. 2019. Functional
997 syndromes as indicators of ecosystem change in temperate grasslands. *Ecological Indicators*, 96,
998 600-610.

999 Tonni, E.P., A.L. Cione, A.J. Figini. 1999. Predominance of arid climates indicated by mammals in the
1000 pampas of Argentina during the Late Pleistocene and Holocene. *Palaeogeography,*
1001 *Palaeoclimatology, Palaeoecology* 147, 257-281

1002 Vallejos, M., Gallego, F., Barbieri, A., Bruzzone, L., Ramos, S., Baeza, S. 2021. MAPBIOMAS PAMPA
1003 SUDAMERICANO: una iniciativa trinacional para conocer los cambios en el uso y cobertura del
1004 suelo. *Revista INIA* 65, 85-89.

1005 Vega, E., Baldi, G., Jobbágy, E. G., & Paruelo, J. 2009. Land use change patterns in the Río de la Plata
1006 grasslands: the influence of phytogeographic and political boundaries. *Agriculture, ecosystems &*
1007 *environment*, 134(3-4), 287-292.

1008 Veldman, J. W., Buisson, E., Durigan, G., Fernandes, G. W., Le Stradic, S., Mahy, G., ... & Bond, W. J.
1009 2015. Toward an old-growth concept for grasslands, savannas, and woodlands. *Frontiers in*
1010 *Ecology and the Environment*, 13(3), 154-162.

1011 Verón, S. R., Blanco, L. J., Texeira, M. A., Irisarri, J. G. N., & Paruelo, J. M. 2018. Desertification and
 1012 ecosystem services supply: The case of the Arid Chaco of South America. *Journal of Arid*
 1013 *Environments*, 159, 66-74.

1014 Vervoorst, F.B. 1967. Las comunidades vegetales de la Depresión del Salado. La Vegetación de la
 1015 República Argentina, Serie Fitogeográfica N° 7. Buenos Aires INTA. 259 p.

1016 Villarino, S. et al. 2020. The role of South American grazing lands in mitigating greenhouse gas
 1017 emissions. A reply to: "Reassessing the role of grazing lands in carbon-balance estimations: Meta-
 1018 analysis and review", by Viglizzo et al. (2019). *The Science of Total Environment* 749.

1019 Vogl, R. J. 1974. Effects of fire on grasslands. *Fire and Ecosystems*. TT Kozlowski & CE Ahlgren, eds.

1020 Volante, J. N., Mosciaro, M. J., Gavier-Pizarro, G. I., & Paruelo, J. M. 2016. Agricultural expansion in
 1021 the Semiarid Chaco: Poorly selective contagious advance. *Land use policy*, 55, 154-165.

1022 Weyland, F., Baudry, J., & Ghera, C. 2019. Short-term effects of a severe drought on avian diversity
 1023 and abundance in a Pampas Agroecosystem. *Austral Ecology*, 44(8), 1340-1350.

1024 Yahdjian, L., Sala, O. E., & Havstad, K. M. 2015. Rangeland ecosystem services: shifting focus from
 1025 supply to reconciling supply and demand. *Frontiers in Ecology and the Environment*, 13(1), 44-
 1026 51.

1027 Zerbino, M. S., Rodríguez, C., & Altier, N. 2006. Earthworms in Agro-ecosystems of
 1028 Uruguay. *Caribbean Journal of Science*, 42(3), 315.

1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048

Figures

Figure 1. Maps of the distribution of Open Ecosystems (OE) (grasslands, savannas, shrublands and wetlands) in the Río de la Plata Grasslands (RPG) region according to three sources highlighted in green, red and blue respectively: a) Copernicus Global Land Cover (ESA) (Buchhorn et al. 2020), b) Baeza and Paruelo (2020) and c) MapBiomias Project (<https://pampa.mapbiomas.org/initiatives>). The map on the right shows a combination of the three sources, highlighting in magenta, yellow, cyan and white the coincidence of at least two sources, while in grey land cover types other than open ecosystems are indicated. On top of the maps are outlined the subregions according to Soriano (1992). A. Rolling Pampa, B-1. Flat Inland Pampa, B-2. West Inland Pampa, C. Southern Pampa, D. Flooding Pampa, E. Mesopotamic Pampa, F. Fluvial Pampa, G. Southern Campos, H. Northern Campos.

Figura 1. Mapas de la distribución de Ecosistemas Abiertos (EA) (pastizales, sabanas, matorrales y humedales) en la región de los Pastizales del Río de la Plata (PRP) según tres fuentes resaltadas en verde, rojo y azul, respectivamente: a) Copernicus Global Land Cover (ESA) (Buchhorn et al.2020), b) Baeza y Paruelo (2020), y c) Proyecto MapBiomias (<https://pampa.mapbiomas.org/initiatives>). El mapa de la derecha muestra una combinación de las tres fuentes, destacando en magenta, amarillo, cian y blanco la coincidencia de al menos dos fuentes, mientras que en gris se indican tipos de cobertura terrestre distintos de los ecosistemas abiertos. Sobre los mapas se indican las subregiones según Soriano (1992). A. Pampa Ondulada, B-1. Pampa Interior Plana, B-2. Pampa Interior Oeste, C. Pampa Austral, D. Pampa Deprimida, E. Pampa Mesopotámica, F. Pampa Fluvial, G. Campos del Sur, H. Campos del Norte.

Figure 2. Schemes of the physiognomy of different Open Ecosystems (OE) across a gradient of grazing (Y axis left), fire (Y axis right) and geographic (X axis) in the RPG.

Figura 2. Esquemas de la fisonomía de diferentes Ecosistemas Abiertos (EA) a lo largo de gradientes de pastoreo (eje Y izquierdo), fuego (eje Y derecho) y geográfico (eje X) en los PRP.

Figure 3. Soil organic carbon (SOC) controls at different temporal scales. Dashed lines show which controls are affected by grazing and fire. ANPP is aboveground net primary production, and BNPP is belowground net primary production. Modified from Piñeiro et al. (2010).

Figura 3. Controles del carbono orgánico del suelo (COS) a diferentes escalas temporales. Las líneas discontinuas muestran qué controles se ven afectados por el pastoreo y el fuego. ANPP (PPNA, por sus siglas en español) es la producción primaria neta aérea y BNPP (PPNB, por sus siglas en español) es la producción primaria neta subterránea. Modificado de Piñeiro et al. (2010).

Figure 4. Mean Human Appropriation of Net Primary Productivity (HANPP) on the ensemble remnant Open Ecosystem (OE) of the Río de la Plata Grasslands (RPG) for the period 2001-2019, calculated according to Baeza and Paruelo (2018). The inserted map in the upper left corner shows the original map of Baeza and Paruelo (2018) including all land covers. The graph in the bottom right corner shows the mean HANPP (%) of OE areas per subregion and country. The lines on the bars indicate the standard deviation. F. Flooding Pampa, S. Southern Pampa, Fi. Flat Inland Pampa. Fl. Fluvial Pampa, M. Mesopotamic Pampa, R. Rolling Pampa, Wi. West Inland Pampa, Nc. Northern Campos, Sc. Southern Campos.

Figura 4. Apropiación Humana de la Productividad Primaria Neta (AHPPN) promedio de los Ecosistemas Abiertos (EA) remanentes del conjunto de los Pastizales del Río de la Plata (PRP) para el período 2001-2019, calculado según Baeza y Paruelo (2018). El mapa insertado en la esquina superior izquierda muestra el mapa original de Baeza y Paruelo (2018) incluyendo todas las coberturas terrestres. El gráfico de la esquina inferior derecha muestra la AHPPN promedio (%) de EA por subregión y país. Las líneas de las barras indican el desvío estándar. F. Pampa Deprimida, S. Pampa Austral, Fi. Pampa Interior Plana. Fl. Pampa Fluvial, M. Pampa Mesopotámica, R. Pampa Ondulada, Wi. Pampa Interior Oeste, Nc. Campos del Norte, Sc. Campos del Sur.

Figure 5. Mean Ecosystem Services (regulation/support) Supply Index (ESSI) on the ensemble remnant Open Ecosystems (OE) of the Río de la Plata Grasslands (RPG) for the period 2001-2019 calculated according to Paruelo et al. (2016). The graph in the upper left corner shows the mean ESSI for Open Ecosystem areas (oe) and other land covers (lc) per subregión and country. The lines on the bars indicate the standard deviation. F. Flooding Pampa, S. Southern Pampa, Fi. Flat Inland Pampa. Fl. Fluvial Pampa, M. Mesopotamic Pampa, R. Rolling Pampa, Wi. West Inland Pampa, Nc. Northern Campos, Sc. Southern Campos.

Figura 5. Índice de Oferta de Servicios Ecosistémicos (IOSE) de regulación y soporte promedio de los Ecosistemas Abiertos (EA) remanentes del conjunto de los Pastizales del Río de la Plata (PRP) para el período 2001-2019, calculado según Paruelo et al. (2016). El gráfico en la esquina superior izquierda muestra el IOSE promedio para áreas de Ecosistemas Abiertos (oe) y otras coberturas terrestres (lc) por subregión y país. Las líneas de las barras indican el desvío estándar. F. Pampa Deprimida, S. Pampa Austral, Fi. Pampa Interior Plana. Fl. Pampa Fluvial, M. Pampa Mesopotámica, R. Pampa Ondulada, Wi. Pampa Interior Oeste, Nc. Campos del Norte, Sc. Campos del Sur.

Figure 6. Trends of the Ecosystem Services Supply Index (ESSI) for the period 2001-2019, being positive (blue), negative (red) and no temporal trends (white). The graph in the upper left corner shows the proportion of the area of each subregión with significant both trends positive and negative and without trend, differentiating between Open Ecosystem areas (oe) and other land covers (lc). F. Flooding Pampa, S. Southern Pampa, Fi. Flat Inland Pampa. Fl. Fluvial Pampa, M. Mesopotamic Pampa, R. Rolling Pampa, Wi. West Inland Pampa, Nc. Northern Campos, Sc. Southern Campos.

Figura 6. Tendencias del Índice de Oferta de Servicios de los Ecosistemas (IOSE) para el período 2001-2019, siendo positivas (azul), negativas (rojo) y sin tendencias temporales (blanco). El gráfico de la esquina superior izquierda muestra la proporción del área de cada subregión con tendencias significativas, tanto positivas como negativas, y sin tendencia, diferenciando entre áreas de Ecosistemas Abiertos (oe) y otras coberturas de suelo (lc). F. Pampa Deprimida, S. Pampa Austral, Fi. Pampa Interior Plana. Fl. Pampa Fluvial, M. Pampa Mesopotámica, R. Pampa Ondulada, Wi. Pampa Interior Oeste, Nc. Campos del Norte, Sc. Campos del Sur.

Supplemental material: Estimation of ANPP and NSP across the Rio de la Plata Grasslands (RPG)

For the estimation of beef production across the Rio de la Plata Grasslands (RPG) a two steps analysis was performed. First, based on the radiative model proposed by Monteith (1972), we estimated the aboveground net primary productivity (ANPP) as the product between the solar incoming photosynthetically active radiation (PARi), the fraction of the photosynthetically active radiation intercepted by green vegetation (fPAR) and the radiation use efficiency to convert the absorbed radiation into biomass (RUE). To do so three land use and land cover classifications of the RPG were used to identify areas in which grassland cover was located. The three land use/land cover used were: 1) Copernicus Global Land Cover Classification developed by European Space Agency (ESA) (Buchhorn et al. 2020) for the year 2019 and which present a spatial resolution of 100 m, 2) Land use land cover classification of the Rio de la Plata Grasslands for the year 2012-2013 which identify perennial forage resources at pixel MODIS scale (~250 m) (Baeza and Paruelo 2020), and 3) MapBiomass Pampa and Atlantic Forest Tinational initiatives (Souza et al. 2020; Vallejos et al. 2021; Milkovic et al. 2021) for the year 2019, which provide a land cover and land use map on a 30 m scale (<https://mapbiomas.org/>). Within each land-use map, and only for areas labelled as grasslands, savannas, wetlands and shrublands, information of solar incoming photosynthetically active radiation and the spectral index EVI (Enhanced Vegetation Index) were obtained monthly from 2001 to 2020 through the web platform Google Earth Engine (Gorelick et al. 2017). PARi data was obtained from ERA5-Land dataset (Muñoz Sabater 2019), which provide monthly-mean averages of surface net solar radiation at global scale from 1981 to present. A scale factor to convert units from $\text{J.m}^{-2}.\text{day}^{-1}$ to $\text{MJ.m}^{-2}.\text{month}^{-1}$ was applied. Also, we considered that the photosynthetic active radiation was 48% of the total incoming solar radiation (McCree 1972). EVI spectral index was obtained from the MODIS satellite mission. Specifically, we used the MOD13Q1 synthetic product, which provides data at a 5.3 ha spatial resolution on a 16-day frequency. EVI data was transformed to fPAR using a locally-calibrated equation: $\text{fAPAR} = 1.4914 * \text{EVI} - 0.1382$. Finally, the product between PARi ($\text{MJ.m}^{-2}.\text{month}^{-1}$) and fPAR, the amount of energy absorbed by active vegetation (APAR), was converted into ANPP using the following equation: $\text{ANPP (g DM.m}^{-2}.\text{month}^{-1})} = -0.1978 + \text{APAR} * 0.4227 * 10$. Monthly ANPP data was integrated to the annual scale and expressed in $\text{kg DM.ha}^{-1}.\text{yr}^{-1}$ through the multiplication by 10. The fitted model to convert from EVI to ANPP was developed by the Laboratorio de Análisis Regional y Teledetección (LART)-IFEVA-CONICET belonging to the Faculty of Agronomy - University of Buenos Aires.

The second step of the analysis consisted in using the ANPP estimations to quantify the beef production across the RPG. To do so, two sources of data (Irisarri et al. 2014 and Gutierrez et al. 2020) were combined to cover the entire ANPP gradient of the RPG. Irisarri et al. (2014) included data from 113 farms arranged into 19 groups according to their mean annual precipitation and geographic location. For this study, we selected 6 groups that corresponded to the ones in the lowest extreme end of the ANPP gradient. Gutierrez et al. (2020) included data from 19 administrative sections from the Basaltic geomorphological unit, within the Campos physiognomic unit of the RPG, located in the highest end of the ANPP gradient. Both sources of data included ranches devoted to either sheep, cow-calf operations and in a minor proportion dairy system. At the same time both, net secondary production (NSP) and ANPP were originally expressed in terms of energy units, $\text{kJ.m}^{-2}.\text{yr}^{-1}$. NSP was transformed to beef production, expressed in $\text{kg LW.ha}^{-1}.\text{yr}^{-1}$, through the following constant: 9900 kJ.kg LW^{-1} . For ANPP we used the conversion factor of 16760 kJ.kg DM^{-1} .

Beef production, expressed in $\text{kg LW.ha}^{-1}.\text{yr}^{-1}$, was positively associated with ANPP, expressed in $\text{kg DM.ha}^{-1}.\text{yr}^{-1}$, across the ANPP gradient observed for the RPG (Figure A1). A particular feature of the fitted model indicates that a unit change in ANPP generates a more than proportionate change in beef production. Because of this pattern the conversion of ANPP dry matter into live weight decreased from more than 100 kg DM per kg of live weight, for sites below 2500 $\text{kg DM.ha}^{-1}.\text{yr}^{-1}$ of ANPP, to 80

to 60 kg of DM per kg of LW or even less than 60 kg of DM per kg of LW for sites located above 4500 kg DM.ha⁻¹.yr⁻¹ of ANPP (Figure A1).

Figure A1: Relation between beef production (BP) and ANPP (in kg.ha⁻¹.yr⁻¹) according to the following fitted model $BP = 0.00007489 * ANPP^{1.6164174}$ (n: 25; Adj. R²: 0.92; p-value < 0.0001). Beef production and ANPP were obtained from two sources, Irisarri et al. (2014) and Gutierrez et al. (2020). Through these two sources we included data from arid to humid rangeland within RPG. Each dot represents the average beef production for an area not less than 10000 ha.

REFERENCES

Baeza, S., & Paruelo, J. M. 2020. Land use/land cover change (2000–2014) in the Rio de la Plata grasslands: an analysis based on MODIS NDVI time series. *Remote Sensing*, 12(3), 381.

Buchhorn, M.; Lesiv, M. ; Tsendbazar, N. - E. ; Herold, M. ; Bertels, L. ; Smets, B. 2020. Copernicus Global Land Cover Layers—Collection 2. *Remote Sensing* (12), Volume 108, 1044.

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. 2017. Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote sensing of Environment*, 202, 18-27.

Gutiérrez, F., Gallego, F., Paruelo, J. M., & Rodríguez, C. 2020. Damping and lag effects of precipitation variability across trophic levels in Uruguayan rangelands. *Agricultural Systems*, 185, 102956.

Irisarri, J. G. N., Oesterheld, M., Golluscio, R. A., & Paruelo, J. M. 2014. Effects of animal husbandry on secondary production and trophic efficiency at a regional scale. *Ecosystems*, 17(4), 738-749.

1268 McCree, K.J., 1972. Test of current definitions of photosynthetically active radiation against leaf
1269 photosynthesis data. *Agric. Meteorol.* 10, 442–453.

1270 Milkovic, M.; Gonzalez Lanzillotta, M.S.; Eljall, A.; Baldassini, P.; Perucca, R.; Insaurrealde, A.; Zurano,
1271 J. P.; Minoli, I.; Leszczuk, A.; Orona, M.; Loran, D.; Gill, A.; Garay, A.; Serafini, J. y Insfrán, P.
1272 2021. MapBiomias Bosque Atlántico: dinámica del uso y cobertura en Argentina y Paraguay. XXIX
1273 Reunión Argentina de Ecología, Tucumán, Agosto 2021.

1274 Monteith, J.L. 1972. Solar radiation and productivity in tropical ecosystems. *J. Appl. Ecol.* 9, 747–766.

1275 Muñoz Sabater, J. 2019. ERA5-Land monthly averaged data from 1981 to present. Copernicus Climate
1276 Change Service (C3S) Climate Data Store (CDS).

1277 Souza, C. M., Z Shimbo, J., Rosa, M. R., Parente, L. L., A Alencar, A., Rudorff, B. F., ... & Azevedo, T.
1278 2020. Reconstructing three decades of land use and land cover changes in brazilian biomes with
1279 landsat archive and earth engine. *Remote Sensing*, 12(17), 2735.

1280 Vallejos, M., Gallego, F., Barbieri, A., Bruzzone, L., Ramos, S., Baeza, S. 2021. MAPBIOMAS PAMPA
1281 SUDAMERICANO: una iniciativa trinacional para conocer los cambios en el uso y cobertura del
1282 suelo. *Revista INIA* 65, 85-89.

1283