1 **Coronaviruses induce the expression of the dual specificity phosphatase DUSP1.**

2 Shahan Mamoor, MS¹

3 ¹Thomas Jefferson School of Law

4 San Diego, CA 92101

5 mammoorsk@tjssl.edu

6 Coronavirus infection is an emerging public health threat in the United States and worldwide
7 (1). We mined published microarray data to perform systems-level analysis of host cell
8 transcription following infection with multiple coronavirus types in order to identify therapeutic
9 targets and host cell vulnerabilities (2, 3). We identified the dual specificity phosphatase
10 DUSP1 as differentially expressed *in vitro* following infection with human coronavirus 229E,
11 HCoV-229E and with Middle East Respiratory Syndrome coronavirus, MERS-CoV. DUSP1
12 expression significantly increases after coronavirus infection *in vitro*. DUSP1 may be of
13 relevance to the pathology of coronavirus infection.

14
15
16
17
18
19
20
21
22
23
24 Keywords: DUSP1, coronavirus, MERS coronavirus, human coronavirus, COVID19, systems
25 biology of viral infection, virus host-cell interactions.

Coronaviruses are a family of enveloped single-stranded positive-sense RNA viruses (4), seven of which are capable of infecting humans (5-10). The COVID19 infection pandemic that is currently a public health problem in the United States and worldwide is caused by a novel coronavirus, the severe respiratory distress syndrome coronavirus 2, or SARS-CoV-2 (1, 10). The death toll from SARS-CoV-2 worldwide has surpassed 46,000 (11). There are no FDA-approved treatments for SARS-CoV-2 infection (12).

To facilitate discovery for the identification of therapeutic targets in coronavirus infection, we mined published transcriptome data from *in vitro* models of coronavirus infection (2, 3). We identified the dual specificity phosphatase DUSP1 as differentially expressed in human cells *in vitro* after infection with the coronaviruses HCoV 229E and MERS-CoV London. The fact that DUSP1 is induced after infection with both human coronavirus 229E and MERS-CoV London suggests that DUSP1 induction may be a property shared by the coronavirus family including SARS-CoV-2.

Methods

We used datasets GSE89167 (2) and GSE56677 (3) for this systems-level differential gene expression analysis of coronavirus infections in conjunction with GEO2R. GSE89167 was generated using Agilent-039494 SurePrint G3 Human GE v2 8x60K Microarray technology. GSE56677 was generated using Agilent-039494 SurePrint G3 Human GE v2 8x60K Microarray technology.

The Benjamin and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used.

A statistical test was performed to evaluate the significance of difference between mRNA expression levels of DUSP1 in the Huh-7 human liver cell line with and without coronavirus

infection, using a two-tailed, unpaired t-test with Welch's correction. A statistical test was performed to evaluate the significance of difference between mRNA expression levels of DUSP1 in the CALU3 2B4 lung adenocarcinoma cell line with coronavirus infection at 3, 7, 12, 18 and 24 hours post-infection as compared to baseline (0 hours) using an ordinary one-way ANOVA with Dunnett's multiple comparisons test. Only p-values less than 0.05 were considered statistically significant. We used PRISM for all statistical analyses (Version 8.4.0)(455).

Results

We mined published transcriptome data from *in vitro* coronavirus infection models in human cell lines to identify genes differentially expressed after coronavirus infection for therapeutic target discovery (2, 3).

DUSP1 is differentially expressed in human coronavirus 229E infected Huh-7 cells relative to uninfected cells.

We identified the dual specificity phosphatase DUSP1 as differentially expressed following infection of the human liver cell line Huh-7 with human coronavirus 229E when compared to uninfected Huh-7 cells. When sorting all of the transcripts produced in the cell based on change in expression between infected and uninfected cells, DUSP1 ranked 26 out of 62976 total transcripts (Table 1). Differential expression of DUSP1 following HCoV 229E infection was statistically significant (Table 1; $p=0.00004175$).

DUSP1 is differentially expressed in MERS-CoV London infected CALU3 2B4 cells when compared to cells infected at baseline (0 hours).

Analysis of a separate dataset revealed that DUSP1 was differentially expressed in the human lung cancer cell line CALU3 2B4 with Middle East Respiratory Syndrome (MERS)

1 coronavirus London (MERS-CoV London) when compared to cells at baseline (0 hours). When
2 sorting all of the transcripts produced in CALU3 2B4 cells based on change in expression
3 between MERS-CoV infection and cells infected at baseline, DUSP1 ranked 16 out of 28653
4 total transcripts (Table 2). Differential expression of DUSP1 following MERS-CoV infection was
5 statistically significant (Table 2; $p=3.79E-15$).
6

7
8 DUSP1 expression is transcriptionally induced in a human liver cell line following infection with
9 human coronavirus 229E.
10

11 Next, we obtained exact expression values for DUSP1 mRNA in cells infected with
12 HCoV 229E to compare DUSP1 expression with and without coronavirus infection. DUSP1 was
13 expressed at higher levels in Huh-7 cells following infection with HCoV 229E when compared to
14 uninfected Huh-7 cells, and this difference was statistically significant (Figure 1; $p=0.0334$). We
15 calculated a 1.678 fold change in DUSP1 expression between HCoV 229E infected and
16 uninfected HuH-7 cells (Table 1).
17

18
19 DUSP1 is initially transcriptionally repressed, and then transcriptionally induced after infection
20 of a human lung cancer cell line with Middle East Respiratory Syndrome (MERS) coronavirus
21 London (MERS-CoV London).

22 We also obtained exact expression values for DUSP1 mRNA in CALU3 2B4 cells
23 infected with MERS-CoV to compare DUSP1 expression with coronavirus infection at 3, 7, 12
24 18 and 24 hours post-infection. DUSP1 infection first dropped at 3 hours post-infection,
25 significantly (Figure 2; $p<0.0001$), then increased at 7 hours but to levels below that of baseline
26 (Figure 2; $p<0.0008$). At 12 hours post-infection, the level of DUSP1 rose to similar levels to
27 that at baseline (Figure 2; $p=0.5559$). At 18 and 24 hours, however, DUSP1 expression was
28 significantly higher than that of baseline. (Figure 2; $p<0.0001$ for both comparisons). We
calculated a 0.848 fold change in DUSP1 expression at 3 hours post-infection, a 0.932 fold

change in DUSP1 expression at 7 hours post-infection, a 1.017 fold change in DUSP1 expression at 12 hours post-infection, a 1.157 fold change in DUSP1 expression at 18 hours post-infection, and a 1.344 fold change in DUSP1 expression at 24 hours post-infection, all relative to baseline (0 hours) (Table 1).

Thus, we identified DUSP1 as among the genes whose expression changed most significantly following coronavirus infection of two separate human cell lines, and found that DUSP1 was expressed at significantly higher levels in both HuH-7 cells and in CALU3 2B4 cells following infection with HCoV 229 and MERS-CoV London, respectively. Importantly, DUSP1 induction following coronavirus infection was conserved between two different types of coronavirus infection: HCoV 229E and MERS-CoV.

Discussion

We mined two published microarray datasets to identify genes differentially expressed in human cells infected with coronaviruses in order to identify potential therapeutic targets. We found that in human cells *in vitro*, (2, 3), coronavirus infection lead to the transcriptional induction of the dual specificity phosphatase DUSP1. DUSP1 was among the genes whose expression changed most significantly when comparing the transcriptomes of a human liver cell line with and without human coronavirus 229E infection. DUSP1 was also among the genes whose expression changed most significantly when comparing the transcriptomes of a human lung cancer cell line infected with MERS coronavirus London infection at 3, 7, 12, 18 and 24 hours post-infection as compared to at baseline (0 hours). DUSP1 expression significantly increased in both cell lines after infection with either HCoV-229E or MERS-CoV London, except for the 3 hours post-infection of CALU3 2B4 cells with MERS-CoV London, where DUSP1 expression was significantly lower than at baseline, before rebounding to levels higher than baseline by 18 hours post-infection.

1 Dual specificity phosphatases function as regulators of MAPK signaling by antagonizing
2 kinases that post-translationally modify substrates in signal transduction pathways (13, 14). By
3 erasing the phosphorylation marks that these kinases generate to positively transduce signals
4 in a cascade, dual specificity phosphatases act as inhibitors of signal transduction. A key
5 substrate of DUSP1 is the p38 mitogen-activated protein kinase (p38 MAPK) (14). Another
6 name for DUSP1 is the MAPK phosphatase, or MKP1 (14).

7
8 Changes in expression of DUSP1 have been reported in association with multiple types
9 of viral infections. These viruses include vaccinia virus, hepatitis C virus, Respiratory Syncytial
10 Virus, Sendai Virus, and the Kaposi Sarcoma herpesvirus (KSHV) (15-18). In fact DUSP1 has
11 previously been implicated in coronavirus biology, in the case of a chicken coronavirus,
12 infectious bronchitis virus (IBV) (19). IL-6 and IL-8 are produced by cells following infection
13 with IBV, and p38 MAPK pathway is critical for this (19). DUSP1 induction by IBV was reported
14 to act as viral mechanism to dampen IL-6 and IL-8 induction by dephosphorylating p38 MAPK
15 (19). We previously reported that IL-6 is among the genes most differentially expressed after
16 human coronavirus infections (20).

17
18 DUSP1 is similarly endogenously transcriptionally activated *in vivo* in response to
19 lipopolysaccharide (LPS), functioning to control the expression of genes expressed following
20 endotoxin shock including CCL3, CCL4, CXCL2, IL-6 and IL-10 (21). DUSP1 is intertwined
21 with multiple cellular processes and pathways, including roles in the cell cycle as a p53
22 transcriptional target (22), involvement in adipocyte differentiation (23), AKT regulation in
23 macrophages during inflammatory responses (24), axon branching involving the brain derived
24 neurotrophic factor (BDNF) (25), and UV-mediated apoptosis (26).

25
26 In summary, we used published microarray datasets to determine that human
27 coronavirus 229E and MERS-CoV infections lead to the transcriptional induction of DUSP1.
28 DUSP1 induction may be a conserved feature of coronaviruses, shared by the novel
coronavirus SARS-CoV-2.

References

1. World Health Organization, 2020. Coronavirus disease 2019 (COVID-19): situation report, 49.
2. Poppe, M., Wittig, S., Jurida, L., Bartkuhn, M., Wilhelm, J., Müller, H., Beuerlein, K., Karl, N., Bhujju, S., Ziebuhr, J. and Schmitz, M.L., 2017. The NF- κ B-dependent and-independent transcriptome and chromatin landscapes of human coronavirus 229E-infected cells. *PLoS pathogens*, 13(3), p.e1006286.
3. Selinger, C., Tisoncik-Go, J., Menachery, V.D., Agnihothram, S., Law, G.L., Chang, J., Kelly, S.M., Sova, P., Baric, R.S. and Katze, M.G., 2014. Cytokine systems approach demonstrates differences in innate and pro-inflammatory host responses between genetically distinct MERS-CoV isolates. *BMC genomics*, 15(1), p.1161.
4. Lai, M.M. and Cavanagh, D., 1997. The molecular biology of coronaviruses. In *Advances in virus research* (Vol. 48, pp. 1-100). Academic Press.
5. Peiris, J.S.M., Lai, S.T., Poon, L.L.M., Guan, Y., Yam, L.Y.C., Lim, W., Nicholls, J., Yee, W.K.S., Yan, W.W., Cheung, M.T. and Cheng, V.C.C., 2003. Coronavirus as a possible cause of severe acute respiratory syndrome. *The Lancet*, 361(9366), pp.1319-1325.
6. Lu, G., Hu, Y., Wang, Q., Qi, J., Gao, F., Li, Y., Zhang, Y., Zhang, W., Yuan, Y., Bao, J. and Zhang, B., 2013. Molecular basis of binding between novel human coronavirus MERS-CoV and its receptor CD26. *Nature*, 500(7461), pp.227-231.
7. van der Hoek, L., Pyrc, K., Jebbink, M.F., Vermeulen-Oost, W., Berkhout, R.J., Wolthers, K.C., Wertheim-van Dillen, P.M., Kaandorp, J., Spaargaren, J. and Berkhout, B., 2004. Identification of a new human coronavirus. *Nature medicine*, 10(4), pp.368-373.
8. van der Hoek, L., Pyrc, K., Jebbink, M.F., Vermeulen-Oost, W., Berkhout, R.J., Wolthers, K.C., Wertheim-van Dillen, P.M., Kaandorp, J., Spaargaren, J. and Berkhout, B., 2004. Identification of a new human coronavirus. *Nature medicine*, 10(4), pp.368-373.
9. Woo, P.C., Lau, S.K., Chu, C.M., Chan, K.H., Tsoi, H.W., Huang, Y., Wong, B.H., Poon, R.W., Cai, J.J., Luk, W.K. and Poon, L.L., 2005. Characterization and complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia. *Journal of virology*, 79(2), pp.884-895.
10. Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R. and Niu, P., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *New England Journal of Medicine*.

11. <https://www.worldometers.info/coronavirus/>
12. Wang, Y. and Zhu, L.Q., 2020. Pharmaceutical care recommendations for antiviral treatments in children with coronavirus disease 2019. *World Journal of Pediatrics*, pp.1-4.
13. Sun, H., Charles, C.H., Lau, L.F. and Tonks, N.K., 1993. MKP-1 (3CH134), an immediate early gene product, is a dual specificity phosphatase that dephosphorylates MAP kinase in vivo. *Cell*, 75(3), pp.487-493.
14. Franklin, C.C. and Kraft, A.S., 1997. Conditional expression of the mitogen-activated protein kinase (MAPK) phosphatase MKP-1 preferentially inhibits p38 MAPK and stress-activated protein kinase in U937 cells. *Journal of Biological Chemistry*, 272(27), pp. 16917-16923.
15. Caceres, A., Perdiguero, B., Gomez, C.E., Cepeda, M.V., Caelles, C., Sorzano, C.O. and Esteban, M., 2013. Involvement of the cellular phosphatase DUSP1 in vaccinia virus infection. *PLoS pathogens*, 9(11).
16. Choi, J.E., Kwon, J.H., Kim, J.H., Hur, W., Sung, P.S., Choi, S.W. and Yoon, S.K., 2015. Suppression of dual specificity phosphatase I expression inhibits hepatitis C virus replication. *PloS one*, 10(3).
17. Robitaille, A.C., Caron, E., Zucchini, N., Mukawera, E., Adam, D., Mariani, M.K., Gélinas, A., Fortin, A., Brochiero, E. and Grandvaux, N., 2017. DUSP1 regulates apoptosis and cell migration, but not the JIP1-protected cytokine response, during Respiratory Syncytial Virus and Sendai Virus infection. *Scientific reports*, 7(1), pp.1-15.
18. Qin, Z., Dai, L., Defee, M., Findlay, V.J., Watson, D.K., Toole, B.P., Cameron, J., Peruzzi, F., Kirkwood, K. and Parsons, C., 2013. Kaposi's sarcoma-associated herpesvirus suppression of DUSP1 facilitates cellular pathogenesis following de novo infection. *Journal of virology*, 87(1), pp.621-635.
19. Liao, Ying, Xiaoxing Wang, Mei Huang, James P. Tam, and Ding Xiang Liu. "Regulation of the p38 mitogen-activated protein kinase and dual-specificity phosphatase 1 feedback loop modulates the induction of interleukin 6 and 8 in cells infected with coronavirus infectious bronchitis virus." *Virology* 420, no. 2 (2011): 106-116.
20. Mamoor, S., 2020. IL6 and IL1RN is differentially expressed and transcriptionally induced in models of SARS coronavirus infection.

- 1 21. Hammer, M., Mages, J., Dietrich, H., Servatius, A., Howells, N., Cato, A.C. and Lang, R.,
2 2006. Dual specificity phosphatase 1 (DUSP1) regulates a subset of LPS-induced genes
3 and protects mice from lethal endotoxin shock. *The Journal of experimental*
4 *medicine*, 203(1), pp.15-20.
- 5 22. Li, M., Zhou, J.Y., Ge, Y., Matherly, L.H. and Wu, G.S., 2003. The phosphatase MKP1 is a
6 transcriptional target of p53 involved in cell cycle regulation. *Journal of Biological*
7 *Chemistry*, 278(42), pp.41059-41068.
- 8 23. Sakaue, H., Ogawa, W., Nakamura, T., Mori, T., Nakamura, K. and Kasuga, M., 2004. Role
9 of MAPK phosphatase-1 (MKP-1) in adipocyte differentiation. *Journal of Biological*
10 *Chemistry*, 279(38), pp.39951-39957.
- 11 24. Perdiguero, E., Sousa-Victor, P., Ruiz-Bonilla, V., Jardí, M., Caelles, C., Serrano, A.L. and
12 Muñoz-Cánoves, P., 2011. p38/MKP-1-regulated AKT coordinates macrophage transitions
13 and resolution of inflammation during tissue repair. *Journal of Cell Biology*, 195(2), pp.
14 307-322.
- 15 25. Jeanneteau, F., Deinhardt, K., Miyoshi, G., Bennett, A.M. and Chao, M.V., 2010. The MAP
16 kinase phosphatase MKP-1 regulates BDNF-induced axon branching. *Nature*
17 *neuroscience*, 13(11), p.1373.
- 18 26. Franklin, C.C., Srikanth, S. and Kraft, A.S., 1998. Conditional expression of mitogen-
19 activated protein kinase phosphatase-1, MKP-1, is cytoprotective against UV-induced
20 apoptosis. *Proceedings of the National Academy of Sciences*, 95(6), pp.3014-3019.
- 21
22
23
24
25
26
27
28

Rank	ID	p-value	t	B	Fold change	Gene	Gene name
26	9093/ NM_004417	0.00004175	1.1E+01	2.194191	1.678	DUSP1	dual specificity phosphatase 1

Table 1: DUSP1 is differentially expressed after infection of a human liver cell line with human coronavirus 229E.

The rank of differential expression, the probe/transcript ID, the p-value of differential expression transcriptome-wide, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, the fold change in expression as compared to uninfected cells, the gene and gene name are listed in this chart.

Rank	ID	p-value	F	Fold change	Gene	Gene name
16	A_23_P110712 NM_004417	3.79E-15	362.7384	3 hr=0.848 7 hr=0.932 12 hr=1.017 18 hr=1.157 24 hr=1.344	DUSP1	dual specificity phosphatase 1

Table 2: DUSP1 is differentially expressed after infection of a human lung cancer cell line with Middle East Respiratory Syndrome (MERS) coronavirus London (MERS-CoV London).

The rank of differential expression, the probe/transcript ID, the p-value of differential expression transcriptome-wide, the F statistic, the fold change in expression as compared to uninfected cells, the gene and gene name are listed in this chart.

Figure 1: DUSP1 is transcriptionally induced in a human liver cell line after infection with human coronavirus 229E.

The mRNA expression level of DUSP1 in uninfected HuH-7 cells and in cells infected with HCoV 229E is graphically represented with mean mRNA levels marked and p-value, the result of a statistical test evaluating significance of difference in DUSP1 expression between groups, listed.

Figure 2: DUSP1 is transcriptionally induced in a human lung cancer cell line after infection with MERS-CoV London.

The mRNA expression level of DUSP1 in uninfected CALU3 2B4 cells and in cells infected with MERS-CoV London at each indicated timepoint post-infection is graphically represented with mean mRNA levels marked and p-values, the results of statistical tests evaluating significance of difference in DUSP1 expression at each time point compared to baseline, listed above.