Differential expression of centromere protein F in cancers of the breast.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY 11730

Breast cancer affects women at relatively high frequency¹. We mined published microarray datasets^{2,3} to determine in an unbiased fashion and at the systems level genes most differentially expressed in the primary tumors of patients with breast cancer. We report here significant differential expression of the gene encoding the centromere protein F, CENPF, when comparing primary tumors of the breast to the tissue of origin, the normal breast. CENPF was also differentially expressed in the tumor cells of patients with triple negative breast cancer. CENPF mRNA was present at significantly higher quantities in tumors of the breast as compared to normal breast tissue. Analysis of human survival data revealed that expression of CENPF in primary tumors of the breast was correlated with overall survival in patients with basal, luminal A, and luminal B subtype cancer, but in a contrary manner. CENPF may be of relevance to initiation, maintenance or progression of cancers of the female breast.

Keywords: breast cancer, CENPF, centromere protein F, systems biology of breast cancer, targeted therapeutics in breast cancer.

Invasive breast cancer is diagnosed in over a quarter of a million women in the United States each year¹ and in 2018, breast cancer was the leading cause of cancer death in women worldwide⁴. While patients with localized breast cancer are provided a 99% 5-year survival rate, patients with regional breast cancer, cancer that has spread to lymph nodes or nearby structures, are provided an 86% 5-year survival rate^{5,6}. Patients with metastasis to distant sites, like the brain, are provided a 27% 5-year survival rate^{5,6}. Understanding how primary tumors are most transcriptionally different from the tissue from which they originate, the breast, can facilitate development of novel diagnostic and therapeutics to promote early detection and enhanced treatment, and contribute to efforts to prevent progression to metastatic stages. We mined published microarray data^{2,3} to understand at the transcriptome level and in an unbiased fashion genes most differentially expressed in primary tumors of the breast as compared to normal breast tissue. The centromere protein F emerged as among the most differentially expressed genes in cancer of the female breast.

Methods

We utilized datasets GSE109169² and GSE38959³ for this differential gene expression analysis of female breast cancer. GSE109169 was generated using Affymetrix Human Exon 1.0 ST Array technology with $n=25$ normal breast tissue and $n=25$ tumors of the breast; analysis was performed using platform GPL5175. The tissues from this first dataset are paired tissues (25 tumors matching 25 breast tissues from 25 patients). GSE38959 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with $n=13$ samples of normal mammary gland ductal cells and $n=30$ samples of tumor cells from patients with triple negative breast cancer; analysis was performed using platform GPL4133. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether CENPF expression was significantly different between primary breast tumors and breast tissue using a two-tailed t -test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of CENPF mRNA expression levels with overall survival (OS) in $n=404$ patients with basal subtype cancer, $n=794$ patients with luminal A subtype cancer, $n=515$ patients with luminal B subtype cancer, and $n=166$ patients with HER2+ cancer.

Results

We performed discovery of genes associated with breast cancer in females by mining two independently published microarray datasets^{2,3}.

CENPF is differentially expressed in primary tumors of the breast.

Comparison of 25 normal breast tissues to 25 tumors of the breast² from patients with early-onset breast cancer revealed that the gene encoding the centromere protein F, encoded by the CENPF gene, was among the genes most differentially expressed in tumors of the breast in human breast cancer (Chart 1). When sorting each of the genes expressed in tumors of the breast based on significance of difference as compared to normal breast tissue, CENPF ranked 200 out of 19076 total transcripts, equating to 99.0% differential expression (Chart 1). Differential expression of CENPF in female breast cancer was statistically significant (Chart 1; $p=3.81e-13$).

Analysis of a second microarray dataset³, here in tumor cells of patients with triple negative breast cancer as compared to normal mammary gland ductal cells, again revealed significant differential expression of CENPF in human breast cancer (Chart 2). When sorting each of the genes expressed in the

tumor cells of patients with triple negative breast cancer based on significance of difference as compared to normal mammary gland ductal cells, CENPF ranked 122 out of 45015 total transcripts, equating to 99.7% differential expression (Chart 2). Differential expression of CENPF in the tumor cells of patients with triple negative breast cancer was statistically significant (Chart 2; $p=1.92\text{e-}12$). These data suggested that differential expression of CENPF was not an artifact of a single microarray dataset, nor was it associated with one type of breast cancer, rather a general feature of cancers of the breast.

CENPF is expressed at significantly higher levels in breast tumors as compared to the breast.

We obtained exact mRNA expression levels for CENPF from the breast and from breast tumors² to understand the magnitude and direction of CENPF expression change. CENPF was expressed at higher levels in tumors of the breast as compared to normal breast tissue (Figure 1). Increased expression of CENPF in primary breast tumors was statistically significant (Figure 1: $p<0.0001$). CENPF was expressed at 5.45 ± 0.42 arbitrary units (AU) in normal breast tissue, while it was expressed at 7.40 ± 0.95 AU in tumors of the breast. We calculated a mean fold change of 1.36 in CENPF mRNA levels when comparing primary tumors of the breast to normal breast tissues.

CENPF expression correlates with survival outcomes in basal and luminal A human breast cancer.

We performed Kaplan-Meier survival analysis in an attempt to correlate CENPF mRNA expression levels with survival outcomes in patients with breast cancer, evaluating correlations between molecular subtype and overall survival. We observed a statistically significant correlation between CENPF expression and overall survival (OS) in patients with luminal A subtype breast cancer (Figure 2; log rank p -value: $1.7\text{e-}05$ for overall survival, hazard ratio: 2.03 (1.46-2.83) (Fig. 2)). CENPF mRNA levels were a negative prognostic indicator in luminal A subtype breast cancer patients. Median OS was 179.01 months for luminal A patients with low tumor expression of CENPF while median OS was 89.04 months for luminal A patients with high tumor expression of CENPF (Chart 3). We also observed a statistically significant correlation between CENPF expression and overall survival (OS) in patients with basal subtype breast cancer (Figure 2; log rank p -value: 0.01 for overall survival, hazard ratio: 0.6 (0.41-0.89) (Fig. 2)) as well in patients with luminal B subtype breast cancer (Figure 2; log rank p -value: 0.022 for overall survival, hazard ratio: 1.51 (1.06-2.15) (Fig. 2)). CENPF mRNA levels were a positive prognostic indicator in basal subtype breast cancer patients. Median OS was 41.04 months for basal subtype patients with low tumor expression of CENPF while median OS was 98.14 months for basal subtype patients with high tumor expression of CENPF (Chart 3). As in luminal A breast cancer, CENPF mRNA levels were a negative prognostic indicator in luminal B subtype breast cancer patients. Median OS was 110.4 months for luminal B subtype patients with low tumor expression of CENPF while median OS was 66.6 months for luminal B subtype patients with high tumor expression of CENPF (Chart 3). CENPF primary tumor expression was not correlated with overall survival in HER2+ breast cancer (Figure 2; log rank p -value: 0.62 for OS, hazard ratio: 1.15 (0.66-2.03) (Fig. 2)).

Thus, through comparative transcriptome analysis of primary tumors of the breast and normal breast tissue, we found that differential expression and up-regulation of CENPF was among the most significant transcriptional features in primary tumors from patients with breast cancer. CENPF expression in primary tumors of the breast was correlated with overall survival in patients with basal, luminal A and luminal B subtype cancers, with high expression of CENPF associating with superior survival in basal subtype cancer but with low expression of CENPF associating with superior survival in luminal A and luminal B subtype cancers.

Discussion

Invasive breast cancer is a medical problem with a 27% 5-year survival rate for women whose disease has spread to distant sites^{5,6}. To facilitate understanding of the basic transcriptional differences between primary tumors of the breast and the tissues from which these tumors originate, normal breast tissues, we performed comparative transcriptome analysis using two independently published microarray datasets^{2,3}, providing evidence here that centromere protein F, encoded by the CENPF, is a defining transcriptional feature of human breast cancer: in early onset-breast cancer, and in triple negative breast cancer. CENPF was expressed at significantly higher levels in primary tumors from patients with breast cancer as compared to normal tissues of the breast. In patients with luminal A and luminal B subtype breast cancer, low expression of CENPF was associated with superior overall survival, while in patients with basal subtype breast cancer, high expression of CENPF was associated with superior overall survival. CENPF and the molecular processes to which it pertains to may be relevant to the initiation or progression of human breast cancer.

References

1. DeSantis, C.E., Ma, J., Goding Sauer, A., Newman, L.A. and Jemal, A., 2017. Breast cancer statistics, 2017, racial disparity in mortality by state. *CA: a cancer journal for clinicians*, 67(6), pp.439-448.
2. Chang, J.W., Kuo, W.H., Lin, C.M., Chen, W.L., Chan, S.H., Chiu, M.F., Chang, I.S., Jiang, S.S., Tsai, F.Y., Chen, C.H. and Huang, P.H., 2018. Wild-type p53 upregulates an early onset breast cancer-associated gene GAS7 to suppress metastasis via GAS7–CYFIP1- mediated signaling pathway. *Oncogene*, 37(30), pp.4137-4150.
3. Komatsu, M., Yoshimaru, T., Matsuo, T., Kiyotani, K., Miyoshi, Y., Tanahashi, T., Rokutan, K., Yamaguchi, R., Saito, A., Imoto, S. and Miyano, S., 2013. Molecular features of triple negative breast cancer cells by genome-wide gene expression profiling analysis. *International journal of oncology*, 42(2), pp.478-506.
4. Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A. and Jemal, A., 2018. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*, 68(6), pp.394-424.
5. ACS Cancer Facts & Figures 2019.
<https://www.cancer.net/cancer-types/breast-cancer/metastatic/statistics>.
6. Survival Rates for Breast Cancer. <https://www.cancer.org/cancer/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-survival-rates.html>
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 200
Probe ID: 2379863
p-value: 3.81e-13
t: -9.68
B: 19.743481
Gene: CENPF
Gene name: centromere protein F

Chart 1: CENPF is differentially expressed in the primary tumors of patients with early-onset human breast cancer.

Rank of differential expression, probe ID, p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Rank: 122
Probe ID: 28683
p-value: 1.92e-12
t: -9.5744056
B: 18.215789
Gene: CENPF
Gene name: centromere protein F

Chart 2: CENPF is differentially expressed in the tumor cells of patients with triple-negative breast cancer.

Rank of differential expression, probe ID, p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, gene and gene name are listed in this chart.

Figure 1: CENPF is expressed at significantly higher levels in primary breast tumors as compared to normal breast tissue.

The mRNA expression level of CENPF in normal breast tissue (left) and in primary tumors of the breast (right) is graphically depicted with the result of a statistical test evaluating significance of difference in CENPF expression between normal breast tissue and primary tumors of the breast, a *p*-value, is $p < 0.0001$.

Figure 2: CENPF expression correlates with overall survival in patients with basal subtype, luminal A subtype, and luminal B subtype breast cancers but not in patients with HER2+ cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival (OS) for $n=404$ patients with basal subtype breast cancer (upper left), $n=166$ patients with HER2+ breast cancer (upper right), $n=794$ patients with luminal A breast cancer (lower left), and $n=515$ patients with luminal B breast cancer (lower right), stratified into two groups, based on low or high expression of CENPF in patient primary tumors. The log rank p-value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on CENPF expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Basal

Low CENPF expression: 41.04 months

High CENPF expression: 98.14 months

Luminal A

Low CENPF expression: 179.01 months

High CENPF expression: 89.04 months

Luminal

Low CENPF expression: 110.4 months

High CENPF expression: 66.6 months

Chart 3: In basal subtype breast cancer, median overall survival is superior in patients with high primary tumor expression of CENPF, while in luminal A and luminal B subtype breast cancers, median overall survival is superior in patients with low primary tumor expression of CENPF.

The overall survival of $n=404$ basal subtype breast cancer patients, $n=794$ luminal A, and $n=515$ luminal B subtype breast cancer patients based on stratification into low or high tumor expression of CENPF is listed in this chart.