ALKBH6 is a differentially expressed gene in brain metastatic human breast cancer.

Shahan Mamoor, MS¹
¹shahanmamoor@gmail.com
East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with brain metastasis in humans with metastatic breast cancer. We found that the alkB, alkylation repair homolog 6, encoded by ALKBH6, was among the genes whose expression was most different in the brain metastases of patients with metastatic breast cancer as compared to primary tumors of the breast. ALKBH6 mRNA was present at increased quantities in brain metastatic tissues as compared to primary tumors of the breast. Importantly, expression of ALKBH6 in primary tumors was significantly correlated with patient recurrence-free survival. Modulation of ALKBH6 expression may be relevant to the biology by which tumor cells metastasize from the breast to the brain in humans with metastatic breast cancer.

Keywords: breast cancer, metastasis, brain metastases, central nervous system metastases, alkB alkylation repair homolog 6, ALKBH6, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with brain metastatic breast cancer^{4,5}. We discovered differential and increased expression of the gene encoding the alkB, alkylation repair homolog 6, ALKBH6, in brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE52604⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=11$ primary breast tumors and $n=3$ brain metastases from patients with breast cancer; analysis was performed using platform GPL885. GSE52604 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer; analysis was performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether ALKBH6 gene expression was significantly different between primary tumors of the breast and brain metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of ALKBH6 mRNA expression levels with recurrence-free survival in $n=1764$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of brain metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

ALKBH6 is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified the

1 alkB, alkylation repair homolog 6, encoded by ALKBH6, as a differentially expressed gene in the
2 breast metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes
3 expressed in brain metastases based on significance of difference as compared to primary tumors
4 of the breast in patients with breast cancer, ALKBH6 ranked 235 out of 17418 total transcripts
5 (Chart 1), equating to 98.7% differential expression. Differential expression of ALKBH6 in the
6 brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=$
7 6.95E-04).

8 Differential gene expression analyses that utilize primary and metastatic tumor tissues to
9 discover genes associated with metastasis can be challenging due to the relatively heterogenous
10 cellular composition and constitution of the tumor as compared to normal, untransformed
11 (benign) tissues. Thus, to attempt to validate differential transcriptome-wide differential
12 expression of ALKBH6 in brain metastases in human metastatic breast cancer, we queried a
13 second microarray dataset⁵, here comparing normal breast tissues rather than primary tumors of
14 the breast to brain metastases. Again, we identified ALKBH6 as a differentially expressed gene
15 in the brain metastatic tissues of patients with breast cancer (Chart 2). When sorting each of the
16 genes expressed in brain metastases based on significance of difference as compared to normal
17 breast tissues, ALKBH6 ranked 455 out of 41093 total transcripts (Chart 2), equating to 98.9%
18 differential expression. Differential expression of ALKBH6 in the brain metastases of patients
19 with metastatic breast cancer was statistically significant (Chart 2; $p=6.08E-09$). Thus,
20 differential expression of ALKBH6, transcriptome-wide, in the brain metastases of women with
21 metastatic breast cancer was conserved across two independent microarray datasets, both when
22 compared to benign and transformed breast tissues.

23 **ALKBH6 is expressed at higher levels in the brain metastases of patients with metastatic 24 breast cancer.**

25 We obtained exact mRNA expression levels for ALKBH6, in primary tumors of the breast
26 and in brain metastasis of patients with brain metastatic breast cancer to determine direction and
27 statistical significance of change in ALKBH6 expression in brain metastatic tissues. We observed
28 significantly higher expression of ALKBH6 in the brain metastases of patients with breast cancer
as compared to primary tumors of the breast: ALKBH6 was expressed at 0.25 ± 0.20 arbitrary
units (AU) in primary tumors of the breast, while it was expressed at 0.85 ± 0.14 AU in brain
metastatic tissues. The difference in ALKBH6 mRNA levels between primary tumors of the
breast and brain metastatic tissues was statistically significant (Figure 1; $p=0.000118$).

29 **ALKBH6 expression is significantly correlated with survival outcomes in human breast 30 cancer.**

31 We performed Kaplan-Meier survival analysis⁸ in $n=1764$ breast cancer patients in total, to
32 evaluate whether ALKBH6 tumor expression was correlated with survival outcomes in breast
33 cancer. We observed a statistically significant correlation between primary tumor expression of
34 ALKBH6 and recurrence-free survival (RFS) in patients with breast cancer (Figure 2). Patients
35 whose primary tumors expressed low levels of ALKBH6 possessed median RFS of 32 months,
36 while patients whose tumors expressed high levels of ALKBH6 possessed median RFS of 38.57
37 months. This difference in RFS based on ALKBH6 tumor expression in patients with breast
38 cancer was statistically significant (Figure 2, Chart 3; logrank p -value: 0.029; hazard ratio: 0.84
(0.72-0.98)).

1 Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we
2 identified the alkB, alkylation repair homolog 6, encoded by ALKBH6, as among the genes
3 whose expression was most different, transcriptome-wide, in the brain metastases of patients with
4 breast cancer, both when compared to primary tumors of the breast and to normal breast tissues;
5 we observed significantly increased expression of ALKBH6 in brain metastases as compared to
6 primary tumors of the breast. Further, we found a significant correlation between ALKBH6
expression and patient survival outcomes, as recurrence-free survival was significantly greater in
patients whose primary tumors expressed higher levels of ALKBH6 as compared to patients
whose primary tumors expressed lower levels of ALKBH6.

7 **Discussion**

8 We provided evidence here that the alkB, alkylation repair homolog 6, encoded by
9 ALKBH6, is among the genes whose expression is most different in the brain metastases of
10 patients with brain metastatic breast cancer, that ALKBH6 mRNA is present at significantly
11 increased quantities in brain metastatic tissues as compared to primary tumors of the breast, and
12 that primary tumor ALKBH6 expression is significantly correlated with patient survival
13 outcomes in human breast cancer. Evaluation of the effects of genetic depletion of ALKBH6 in
14 mouse models of metastatic breast cancer on metastasis to the central nervous system is merited.
15 Modulation of ALKBH6 expression may be relevant to the processes by which breast cancer cells
16 exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune
17 clearance, breach the blood-brain barrier and colonize the brain.
18
19
20
21
22
23
24
25
26
27
28

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Salhia, B., Kiefer, J., Ross, J.T., Metapally, R., Martinez, R.A., Johnson, K.N., DiPerna, D.M., Paquette, K.M., Jung, S., Nasser, S. and Wallstrom, G., 2014. Integrated genomic and epigenomic analysis of breast cancer brain metastasis. *Plos one*, 9(1), p.e85448.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 235
Probe ID: 9317
p-value: 6.95E-04
t: -4.2
B: -0.257717
Gene: ALKBH6
Gene name: alkB, alkylation repair homolog 6

Chart 1: ALKBH6 is differentially expressed in brain metastatic breast cancer when comparing brain metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 455
probe ID: A_23_P39402
p-value: 6.08E-09
t: -7.086969
B: 10.354159
Gene: ALKBH6
Gene name: alkB, alkylation repair homolog 6

Chart 2: ALKBH6 is differentially expressed in brain metastatic breast cancer when comparing brain metastases to normal breast tissues.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: ALKBH6 is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of ALKBH6 in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in ALKBH6 expression between primary tumors of the breast and brain metastases, a *p*-value, is *p*=0.001371.

Figure 2: Significant correlation between ALKBH6 primary tumor expression and recurrence-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=1764$ total patients stratified into two groups, based on low or high expression of ALKBH6 in patient primary tumors. The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on ALKBH6 expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low ALKBH6 expression: 32 months
High ALKBH6 expression: 38.57 months

Chart 3: Median recurrence-free survival is inferior in patients with low primary tumor expression of ALKBH6.

The median RFS (recurrence-free survival) of patients with low primary tumor expression of ALKBH6 and high primary tumor expression of ALKBH6 is listed in this chart, for $n=1764$ breast cancer patients.