1 **Antipsychotics modulate the expression of the 5' ecto-nucleotidase NT5E (CD73) in the**
2 **brains of patients with schizophrenia.**

3 Shahan Mamoor¹

4 ¹Thomas Jefferson School of Law

5 mamoorisk@tjssl.edu

6 San Diego, CA 92101

7 Antipsychotic medications are used in patients with schizophrenia and related psychotic
8 disorders to control positive symptoms (1, 2). Positive symptoms include hallucinations -
9 seeing or hearing things that are not present, and delusions - fixed false beliefs (3). The use of
10 antipsychotics was fortuitously arrived at in this patient population (4) and though it is
11 understood which receptor subtypes are pharmacologically modulated by these drugs (5, 6, 7)
12 there is still a lack of understanding regarding the specific therapeutic mechanism, at the level
13 of gene expression, by which they exert their function (7, 8, 9). By mining a published
14 microarray dataset (10), we found that the 5' ecto-nucleotidase NT5E, also known as CD73,
15 was among the genes whose expression was most different in the dorsolateral frontal cortex
16 (DLPFC) of patients with schizophrenia as compared to the DLPFC of control subjects.
17 Analysis of a separate public dataset (11) revealed that schizophrenic patients treated with
18 antipsychotics possessed significantly increased mRNA expression of NT5E in the deep
19 pyramidal neurons when compared to the deep pyramidal neurons of control subjects. This
20 phenomena appeared to be a specific effect of antipsychotic treatment as we could not detect
21 differential expression of NT5E in untreated psychotic patients. These data reveal significantly
22 increased expression of NT5E in the brains of patients with psychotic disorders and suggest
23 that NT5E is a direct target of antipsychotic medications.

24 **Keywords:** NT5E, CD73, schizophrenia, psychosis, systems biology of schizophrenia, targeted
25 therapeutics in psychiatry.

Psychosis is characterized by a disconnection with reality as evidenced by impairments in reality testing (12, 13). The symptoms experienced by psychotic patients such as schizophrenics include positive symptoms and negative symptoms (3). Positive symptoms include hallucinations, which can be either visual - seeing things that are not present - or auditory - hearing things that are not present (3). Negative symptoms include avolition, or loss of motivation and anhedonia, an inability to experience pleasure (3). The major class of therapeutics used to treat psychosis include a family of drugs known as the antipsychotics (4). The therapeutic function of "typical" antipsychotics such as chlorpromazine and haloperidol was arrived at fortuitously (4) and lauded as a major advance for a group of patients for which institutionalization was a reality (14, 15). Receptor blockade at dopaminergic D2 receptors is thought to be a major pharmacological mechanism by which these drugs modulate neurotransmission (5-8). Later, "atypical" antipsychotics, also known as second-generation antipsychotics were introduced, including the drugs risperidone, olanzapine, and quetiapine, and are generally characterized by relatively greater serotonergic receptor blockade and thus a lower ratio of blockade at dopaminergic receptors relative to blockade at serotonergic receptors as compared to first-generation or typical antipsychotics (16). Antipsychotics are powerful drugs with action at multiple receptor subtypes, and their use is accompanied by significant morbidities including metabolic side effects and movement disorders (17-19). A major issue in understanding how this class of drugs, as a group, exert their therapeutic function in ameliorating positive symptoms of patients with psychotic disorders, beyond dopamine and serotonin receptor blockade, is a lack of understanding as to the etiology of psychotic disorders (20). The development of enhanced pharmacological management tools for psychotic disorders, with an increase in therapeutic efficacy and a decrease in undesirable side effects such as akathisia (21) that drive non-compliance (22) in patients, can be significantly facilitated by an understanding of which genes and gene expression programs are specifically affected by antipsychotic medications. The question of which genes and gene

expression programs are modulated by antipsychotic medications is one in psychopharmacology that has yet to be fully answered (23).

To identify specific genes whose mRNAs were differentially expressed in psychosis, we mined published and public transcriptome datasets (10, 11) generated using the brain tissues of patients with psychotic disorders including schizophrenia. We found that the 5' ecto-nucleotidase NT5E, also known as CD73 (24), was among the genes whose expression was most significantly different in the dorsolateral frontal cortex of patients with schizophrenia (10). Analysis of a separate dataset revealed that NT5E was among the genes whose expression was most significantly different in the deep pyramidal neurons of patients treated with antipsychotic medications (11). NT5E is a gene whose expression is among the most significantly different in the brains of patients with psychotic disorders treated with antipsychotic medications and may be a specific therapeutic target of this class of drugs.

Methods

Differential Gene Expression Analyses

We utilized datasets GSE120340 and GSE145554 for this differential gene expression analysis in conjunction with GEO2R. GSE120340 (10) was generated using Affymetrix Human Genome U133 Plus 2.0 Array technology with post-mortem tissues from Brodmann's area 46 from control subjects (n=10 samples utilized in this analysis) and from patients with psychotic disorders (schizophrenia and patients with psychotic bipolar disorder) (n=16 samples utilized in this analysis). GSE145554 (11) was generated using Affymetrix Human Gene 1.0 ST Array technology from deep pyramidal neurons isolated by laser capture microdissection from control subjects (n=11 samples utilized in this analysis) and from patients with schizophrenia that had been treated with antipsychotic medications (n=8 samples utilized in this analysis). We also utilized GSE145554 to compare the transcriptomes of deep pyramidal neurons from control subjects (n=11 samples utilized in this analysis) and from the deep pyramidal neurons of patients with schizophrenia that had not been treated with antipsychotic medications (n=2

1 samples utilized in this analysis). The Benjamini and Hochberg method of p-value adjustment
2 was used for ranking of differential expression but raw p-values were used for assessment of
3 statistical significance of global differential expression. Log-transformation of data was auto-
4 detected, and the NCBI generated category of platform annotation was used.

5 **Statistical Analyses**

6
7 A statistical test was performed to evaluate whether NT5E expression was significantly
8 different between the dorsolateral prefrontal cortex of control subjects and the dorsolateral
9 prefrontal cortex of psychotic patients using a two-tailed, unpaired t-test with Welch's correction.
10 A statistical test was performed to evaluate whether NT5E expression was significantly different
11 between the deep pyramidal neurons of control subjects and the deep pyramidal neurons of
12 antipsychotic-treated schizophrenic patients using a two-tailed, unpaired t-test with Welch's
13 correction. A separate statistical test was performed to evaluate whether NT5E expression was
14 significantly different between the deep pyramidal neurons of control subjects and the deep
15 pyramidal neurons of schizophrenic patients who were not bring treated with antipsychotic
16 medications using a two-tailed, unpaired t-test with Welch's correction. Only p-values less than
17 0.05 were considered statistically significant. We used PRISM for all statistical analyses
18 (Version 8.4.0)(455).
19
20
21

22 **Results**

23 We mined two independent published (10) and public (11) microarray datasets
24 containing transcriptome data from the brain tissues of patients with psychosis and from the
25 brain tissues of patients with psychosis that were treated with anti-psychotic medications, as
26 compared to the same regions in control subjects, in order to identify the genes whose
27 expression was quantitatively most significantly different in patents with psychosis and
28 antipsychotic-treated psychotic patients.

NT5E is differentially expressed in the dorsolateral prefrontal cortex of patients with psychotic disorders when compared to control subjects.

We found that the 5' ecto-nucleotidase NT5E, also known as CD73, was among the genes most differentially expressed in the dorsolateral frontal cortex of patients with psychotic disorders as compared to the same brain region in control subjects. The psychotic disorders included in the patient cohort tested included schizophrenia and psychotic bipolar disorder. When sorting each of the transcripts detected by microarray based on change in expression between patients with psychotic disorders and control subjects, NT5E ranked 217 out of 19070 total transcripts. Differential expression of NT5E in the dorsolateral prefrontal cortex of patients with psychotic disorders was statistically significant (Table 1; $p=0.0025114$).

NT5E is differentially expressed in the deep pyramidal neurons of antipsychotic-treated schizophrenics when compared to control subjects.

We also identified NT5E as one of the most differentially expressed genes in the deep pyramidal neurons of schizophrenics when comparing patients treated with antipsychotic medications, with the deep pyramidal neurons of control subjects. When sorting each of the transcripts detected by microarray based on change in expression between patients with schizophrenia taking antipsychotic medications and in control subjects, NT5E ranked 153 out of 32321 total transcripts. Differential expression of NT5E in the deep pyramidal neurons of patients with schizophrenia treated with antipsychotic medications was statistically significant (Table 2; $p=0.022347$).

NT5E is transcriptionally induced in the dorsolateral prefrontal cortex of patients with psychotic disorders when compared to control subjects.

Next, we extracted mRNA expression levels for NT5E from each patient tissue sample and from each control subject to assess exactly how different NT5E levels were between the dorsolateral prefrontal cortex of patients with psychotic disorders and control subjects. We

5 OF 19

also performed a statistical test to determine if this difference in NT5E expression was statistically significant. NT5E levels were significantly higher in the dorsolateral prefrontal cortex of patients with psychotic disorders as compared to control subjects (Figure 1; $p=0.0054$). We calculated a 1.1117 ± 0.0668 fold change in NT5E expression in the dorsolateral prefrontal cortex of patients with psychotic disorders relative to the dorsolateral prefrontal cortex of control subjects (Table 1).

NT5E is transcriptionally induced in the deep pyramidal neurons of antipsychotic-treated schizophrenics when compared to control subjects.

We also extracted mRNA expression levels for NT5E from each deep neuron patient tissue sample and from each deep neuron control tissue sample to assess exactly how different NT5E levels were between the deep pyramidal neurons of antipsychotic-treated patients with schizophrenia and control subjects. We also performed a statistical test to determine if this difference in NT5E expression was statistically significant. NT5E levels were significantly higher in the deep pyramidal neurons of patients with schizophrenia that were treated with antipsychotic medications as compared to control subjects (Figure 2; $p=0.0099$). We calculated a 1.0448 ± 0.0320 fold change in NT5E expression in the deep pyramidal neurons of schizophrenic patients relative to the deep pyramidal neurons of control subjects (Table 2).

NT5E is not differentially expressed in the deep pyramidal neurons of schizophrenics that are not being treated with antipsychotics when compared to control subjects.

Finally, we sought to determine whether NT5E was inherently differentially expressed in the brains of patients with psychotic disorders or whether this was a specific effect of antipsychotic treatment. Using the same dataset, we compared the pyramidal deep neuron transcriptomes of two patients that were schizophrenic but not being treated with antipsychotic medications with the transcriptomes of deep pyramidal neurons from control subjects. When

1 sorting each of the transcripts detected by microarray based on change in expression between
2 patients with schizophrenia not treated with antipsychotics and control subjects, NT5E ranked
3 26197 out of 32321 total transcripts. Differential expression of NT5E in the deep pyramidal
4 neurons of patients with schizophrenia not treated with antipsychotic medications was not
5 statistically significant (Table 3; $p=0.82195286$). Thus NT5E was not differentially expressed in
6 the deep pyramidal neurons of patients with schizophrenia that were not being treated with
7 antipsychotics.
8

9
10 Together, we found that NT5E was among the genes whose expression was most
11 significantly different in the dorsolateral prefrontal cortex of patients with psychotic disorders
12 and in the deep pyramidal neurons of patients with schizophrenia treated with antipsychotic
13 medications when compared to control subjects. Importantly, the expression of NT5E was not
14 significantly different in the deep pyramidal neurons of patients with schizophrenia who were
15 not being treated with antipsychotics when compared to control subjects. NT5E mRNA
16 expression was significantly higher in the dorsolateral prefrontal cortex of patients with
17 psychotic disorders and in the deep pyramidal neurons of schizophrenic patients treated with
18 antipsychotics when compared to control subjects. We conclude that NT5E is a specific
19 transcriptional target of antipsychotic medications in the brains of patients with psychotic
20 disorders.
21

22 Discussion

23
24 The common thread that unifies schizophrenic patients and patients with psychotic
25 bipolar disorder is the presence of psychosis - a disconnection with reality characterized by
26 impairments in reality testing (12, 13). The symptoms of psychotic disorders - hallucinations
27 and paranoid delusions - are treated with antipsychotic medications which at the level of
28 neurotransmission generally share the ability to block dopaminergic signaling at D2 receptor
subtypes. However, it is not clear that a dopaminergic hypothesis can explain the totality of

1 schizophrenia pathogenesis as drugs with markedly different dopamine blockade capacity do
2 not differ in efficacy of symptom resolution (9). Moreover, current treatment regimens possess
3 undesirable side effect profiles and leave room for improvement in terms of efficacy and
4 compliance (17-19, 22). To assess the transcriptional landscape of the brain in psychotic
5 disorders and of the anti-psychotic treated brain in an unbiased manner, we mined two
6 microarray datasets, published and public (10, 11), to identify genes whose expression was
7 most significantly different. We identified the 5'-ecto-nucleotidase NT5E (CD73) as a
8 differentially expressed gene in the dorsolateral frontal cortex of patients with schizophrenia
9 and psychotic bipolar disorder (10). We also found that NT5E was differentially expressed in
10 the deep pyramidal neurons of patients with schizophrenia who were being treated with
11 antipsychotic medications, when compared to control subjects (11). Critically, NT5E was not
12 differentially expressed in the deep pyramidal neurons of schizophrenic patients who were not
13 being treated with antipsychotic medications, when compared to control subjects.
14

15
16 NT5E is a membrane-bound glycoprotein involved in the conversion of AMP to
17 adenosine, a molecule with pleiotropic roles in immunity, cancer and in the brain (24).
18 Adenosine is composed of a purine ring attached to a ribose sugar (24, 25). Adenosine can
19 signal in the extracellular milieu through binding to one of four G-protein couple receptors, A1,
20 A2A, A2B, and A3, or it can be internalized to the cytosol (25). Adenosine has been described
21 as important for adaptation in response to low levels of oxygen (hypoxia), for the integrity of
22 barrier tissues by the sealing of epithelial tissues after neutrophil transmigration, in the
23 inflammatory response and in the host response to pathogens (24). CD73-deficient mice
24 manifest abnormalities in platelet biology, as bleeding time after tail-tip resection is significantly
25 reduced (26). However, *ex vivo* analysis of platelet aggregate in response to ADP did not
26 demonstrate any significant difference between the platelets of wild-type and CD73-deficient
27 mice (24, 26).
28

1 Our findings are contrary to that of O'Donovan et al. who showed that there was no
2 significant difference in NT5E expression in the dorsolateral pre-frontal cortex of parents with
3 schizophrenia when compared to controls (27). One reason for this difference could be the
4 method of analysis used: the measurement of the expression of a single gene using
5 quantitative reverse transcriptase PCR, which is dependent on oligonucleotide primer design,
6 reaction conditions, and the housekeeping gene used to normalize, versus transcriptome
7 analysis by microarray, which has its own technical limitations. Nevertheless, we found that of
8 all the transcripts measured by microarray in two separate datasets generated using the brain
9 tissues of patients with psychotic disorders, NT5E was among the genes whose expression
10 changed most significantly. When directly comparing NT5E mRNA expression levels between
11 control and patient populations, this difference was significantly different. There is otherwise
12 limited information describing a role for NT5E in schizophrenia biology but one genome-wide
13 association study identified a single nucleotide polymorphism (rs217331) in the SNAP91 gene
14 that associated with the risk of developing schizophrenia; the promoter region of SNAP91
15 interacts with the promoter of NT5E at the structural level (28). An adenosine hypothesis for
16 schizophrenia has been described that integrates dysfunction in both dopaminergic and
17 glutamatergic signaling (29). These findings may be of relevance to this hypothesis to this
18 model which proposes reduced adenosine function in patients with schizophrenia; we found
19 that antipsychotic medications increased expression of a molecule that converts AMP to
20 adenosine. It remains unclear whether NT5E induction in the brain caused by antipsychotics is
21 associated with a therapeutic benefit; this could be tested by using stereotactic injection of
22 adeno-associated virus (30) to express NT5E in the mouse brain using mouse models of
23 psychosis induced by ketamine or apomorphine (31, 32) or in reeler heterozygous mice (33).

27 We used systems-level transcriptome analysis using published and public datasets to
28 identify differential expression of the 5' ecto-nucleotidase NT5E in the dorsolateral frontal
cortex of patients with schizophrenia and psychotic bipolar disorder and in the deep pyramidal

neurons of patients with schizophrenia who had been treated with antipsychotic medications. NT5E mRNA levels were significantly higher in both the dorsolateral prefrontal cortex and in the deep pyramidal neurons of antipsychotic-treated patients with psychotic disorders. NT5E was not differentially expressed in the deep pyramidal neurons of patients who were not being treated with antipsychotics, and thus we conclude that antipsychotic medications cause the transcriptional induction of NT5E. These findings highlight the power of systems-level transcriptome analysis to identify, in unbiased fashion fundamental differences in a complex genetic diseases where it has been historically difficult to draw conclusions on the molecular basis of disease due to the number of genes that contribute to a complex phenotype - psychosis.

References

1. Lieberman, J.A., Stroup, T.S., McEvoy, J.P., Swartz, M.S., Rosenheck, R.A., Perkins, D.O., Keefe, R.S., Davis, S.M., Davis, C.E., Lebowitz, B.D. and Severe, J., 2005. Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *New England journal of medicine*, 353(12), pp.1209-1223.
2. Leucht, S., Corves, C., Arbter, D., Engel, R.R., Li, C. and Davis, J.M., 2009. Second-generation versus first-generation antipsychotic drugs for schizophrenia: a meta-analysis. *The Lancet*, 373(9657), pp.31-41.
3. Andreasen, N.C., Nopoulos, P., Schultz, S., Miller, D., Gupta, S., Swayze, V. and Flaum, M., 1994. Positive and negative symptoms of schizophrenia: past, present, and future. *Acta Psychiatrica Scandinavica*, 90, pp.51-59.
4. Jašović-Gašić, M., Vuković, O., Pantović, M., Cvetić, T. and Marić Bojović, N., 2012. Antipsychotics—history of development and field of indication, new wine—old glasses. *Psychiatr Danub*, 24(Suppl 3), pp.342-4.
5. Meltzer, H.Y., 1991. The mechanism of action of novel antipsychotic drugs. *Schizophrenia bulletin*, 17(2), pp.263-287.
6. Seeman, P., Chau-Wong, M., Tedesco, J. and Wong, K., 1975. Brain receptors for antipsychotic drugs and dopamine: direct binding assays. *Proceedings of the National Academy of Sciences*, 72(11), pp.4376-4380.
7. Jibson, M.D., 2016. Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. Waltham, MA (accessed 12 January 2017).
8. Miller, R., 2009. Mechanisms of action of antipsychotic drugs of different classes, refractoriness to therapeutic effects of classical neuroleptics, and individual variation in sensitivity to their actions: Part I. *Current neuropharmacology*, 7(4), pp.302-314.
9. Insel, T.R., 2010. Rethinking schizophrenia. *Nature*, 468(7321), pp.187-193.
10. Abdolmaleky, H.M., Gower, A.C., Wong, C.K., Cox, J.W., Zhang, X., Thiagalingam, A., Shafa, R., Sivaraman, V., Zhou, J.R. and Thiagalingam, S., 2019. Aberrant transcriptomes and DNA methylomes define pathways that drive pathogenesis and loss of brain laterality/asymmetry in schizophrenia and bipolar disorder. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 180(2), pp.138-149.

11. GSE145554. Robert McCullumsmith, University of Toledo. Toledo, OH, USA. Wu X, Shukla R, Alganem K, Depasquale E, Reigle J, Simmons M, Hahn C, Haroutunian V, Meller J, Meador-Woodruff J, McCullumsmith R
12. Cannon, T.D., 2015. How schizophrenia develops: cognitive and brain mechanisms underlying onset of psychosis. *Trends in cognitive sciences*, 19(12), pp.744-756.
13. Arciniegas, D.B., 2015. Psychosis. *Continuum: Lifelong Learning in Neurology*, 21(3 Behavioral Neurology and Neuropsychiatry), p.715.
14. Ban, T.A., 2007. Fifty years chlorpromazine: a historical perspective. *Neuropsychiatric disease and treatment*, 3(4), p.495.
15. Pow, J.L., Baumeister, A.A., Hawkins, M.F., Cohen, A.S. and Garand, J.C., 2015. Deinstitutionalization of American public hospitals for the mentally ill before and after the introduction of antipsychotic medications. *Harvard review of psychiatry*, 23(3), pp.176-187.
16. Meltzer, H.Y., 2013. Update on typical and atypical antipsychotic drugs. *Annual review of medicine*, 64, pp.393-406.
17. Woods, S.W., Morgenstern, H., Saksa, J.R., Walsh, B.C., Sullivan, M.C., Money, R., Hawkins, K.A., Gueorguieva, R.V. and Glazer, W.M., 2010. Incidence of tardive dyskinesia with atypical and conventional antipsychotic medications: prospective cohort study. *The Journal of clinical psychiatry*, 71(4), p.463.
18. Ray, W.A., Chung, C.P., Murray, K.T., Hall, K. and Stein, C.M., 2009. Atypical antipsychotic drugs and the risk of sudden cardiac death. *New England Journal of Medicine*, 360(3), pp. 225-235.
19. De Hert, M., Detraux, J., Van Winkel, R., Yu, W. and Correll, C.U., 2012. Metabolic and cardiovascular adverse effects associated with antipsychotic drugs. *Nature Reviews Endocrinology*, 8(2), p.114.
20. Zwicker, A., Denovan-Wright, E.M. and Uher, R., 2018. Gene–environment interplay in the etiology of psychosis. *Psychological medicine*, 48(12), pp.1925-1936.
21. Kane, J.M., Fleischhacker, W.W., Hansen, L., Perlis, R., Pikalov III, A. and Assuncao-Talbott, S., 2009. Akathisia: an updated review focusing on second-generation antipsychotics. *Journal of Clinical Psychiatry*, 70(5), p.627.

22. Ww, F. and U, M., 1994. Compliance with antipsychotic drug treatment: influence of side effects. *Acta Psychiatrica Scandinavica*, 89, pp.11-15.
23. Voisey, J., Mehta, D., McLeay, R., Morris, C.P., Wockner, L.F., Noble, E.P., Lawford, B.R. and Young, R.M., 2017. Clinically proven drug targets differentially expressed in the prefrontal cortex of schizophrenia patients. *Brain, behavior, and immunity*, 61, pp.259-265.
24. Colgan, S.P., Eltzschig, H.K., Eckle, T. and Thompson, L.F., 2006. Physiological roles for ecto-5'-nucleotidase (CD73). *Purinergic signalling*, 2(2), p.351.
25. Klaasse, E.C., IJzerman, A.P., Willem, J. and Beukers, M.W., 2008. Internalization and desensitization of adenosine receptors. *Purinergic signalling*, 4(1), pp.21-37.
26. Koszalka, P., Özüyaman, B., Huo, Y., Zerneck, A., Flögel, U., Braun, N., Buchheiser, A., Decking, U.K., Smith, M.L., Sévigny, J. and Gear, A., 2004. Targeted disruption of cd73/ecto-5'-nucleotidase alters thromboregulation and augments vascular inflammatory response. *Circulation research*, 95(8), pp.814-821.
27. O'Donovan, S.M., Sullivan, C., Koene, R., Devine, E., Hasselfeld, K., Moody, C.L. and McCullumsmith, R.E., 2018. Cell-subtype-specific changes in adenosine pathways in schizophrenia. *Neuropsychopharmacology*, 43(8), pp.1667-1674.
28. Buxton, D.S., Batten, D.J., Crofts, J.J. and Chuzhanova, N., 2019. Predicting novel genomic regions linked to genetic disorders using GWAS and chromosome conformation data—a case study of schizophrenia. *Scientific Reports*, 9(1), pp.1-14.
29. Boison, D., Singer, P., Shen, H.Y., Feldon, J. and Yee, B.K., 2012. Adenosine hypothesis of schizophrenia—opportunities for pharmacotherapy. *Neuropharmacology*, 62(3), pp. 1527-1543.
30. Xu, R., Janson, C.G., Mastakov, M., Lawlor, P., Young, D., Mouravlev, A., Fitzsimons, H., Choi, K.L., Ma, H., Dragunow, M. and Leone, P., 2001. Quantitative comparison of expression with adeno-associated virus (AAV-2) brain-specific gene cassettes. *Gene therapy*, 8(17), pp.1323-1332.
31. Schobel, S.A., Chaudhury, N.H., Khan, U.A., Paniagua, B., Styner, M.A., Asllani, I., Inbar, B.P., Corcoran, C.M., Lieberman, J.A., Moore, H. and Small, S.A., 2013. Imaging patients with psychosis and a mouse model establishes a spreading pattern of hippocampal dysfunction and implicates glutamate as a driver. *Neuron*, 78(1), pp.81-93.

- 1 32. Dixit, T.S., Sharma, A.N., Lucot, J.B. and Elased, K.M., 2013. Antipsychotic-like effect of
2 GLP-1 agonist liraglutide but not DPP-IV inhibitor sitagliptin in mouse model for
3 psychosis. *Physiology & behavior*, 114, pp.38-41.
- 4 33. Tueting, P., Doueiri, M.S., Guidotti, A., Davis, J.M. and Costa, E., 2006. Reelin down-
5 regulation in mice and psychosis endophenotypes. *Neuroscience & Biobehavioral*
6 *Reviews*, 30(8), pp.1065-1077.
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28

Rank	ID	p-value	t	B	Fold change	Gene	Gene name
GSE120340							
217	4907_at	0.0025114	3.32437858	-1.49009	1.1117 ± 0.0668	NT5E	5'- nucleotidase, ecto (CD73)

Table 1: NT5E is differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to controls.

The rank of differential expression out of all transcripts globally, the transcript/probe ID, the p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Fold change	Gene	Gene name
GSE145554							
153	8120967	0.022347	2.4730769	-3.61	1.0448 ± 0.0320	NT5E	5'-nucleotidase, ecto (CD73)

Table 2: NT5E is differentially expressed in the deep pyramidal neurons of antipsychotic-treated schizophrenics when compared to controls.

The rank of differential expression out of all transcripts globally, the transcript/probe ID, the p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
GSE145554						
26197	8120967	0.82195286	0.2292045	-4.76	NT5E	5'-nucleotidase ecto

Table 3: NT5E is not differentially expressed in the deep pyramidal neurons of schizophrenics that are not treated with antipsychotics when compared to controls.

The rank of differential expression out of all transcripts globally, the transcript/probe ID, the p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Figure 1: NT5E is transcriptionally induced in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to controls.

The mRNA expression levels of NT5E in the dorsolateral frontal cortex of control patients and in psychotic patients are graphically represented on the left by individual patient with mean value marked and on the right by box and whisker plot, with bars extending from minimum to maximum value with median value marked. The result of a statistical test evaluating the significance of difference in NT5E expression between the dorsolateral frontal cortex of control patients and psychotic patients, a p-value, is denoted on each graph.

Figure 2: NT5E is transcriptionally induced in the deep pyramidal neurons of antipsychotic-treated schizophrenics when compared to controls.

The mRNA expression levels of NT5E in the deep neurons of control patients and in the deep neurons of antipsychotic-treated schizophrenics are graphically represented on the left by individual patient with mean value marked and on the right by box and whisker plot, with bars extending from minimum to maximum value with median value marked. The result of a statistical test evaluating the significance of difference in NT5E expression between the deep neurons of control patients and antipsychotic treated schizophrenics, a p-value, is denoted on each graph.