Fusarium graminearum Species Complex: A Bibliographic Analysis and Web-Accessible Database for Global Mapping of Species and Trichothecene Toxin chemotypes

Emerson M. Del Ponte^{1*}, Gláucia M. Moreira¹, Todd J. Ward², Kerry O'Donnell², Camila P. Nicolli¹, Franklin J. Machado¹, Máira R. Duffeck¹, Kaique S. Alves¹, Dauri J. Tessmann³, Cees Waalwijk⁴, Theo van der Lee⁴, Hao Zhang⁵, Sofia N. Chulze⁶, Sebastian A. Stenglein⁷, Dinorah Pan⁸, Silvana Vero⁸, Lisa J. Vaillancourt⁹, David G. Schmale III¹⁰, Paul D. Esker¹¹, Antonio Moretti¹², Antonio F. Logrieco¹², H. Corby Kistler¹³, Gary C. Bergstrom¹⁴, Altus Viljoen¹⁵, Lindy J. Rose¹⁵, Gert J. van Coller¹⁶, Theresa Lee¹⁷

¹ Departamento de Fitopatologia, Universidade Federal de Viçosa, Viçosa, MG, Brazil

² USDA, Agricultural Research Service, National Center for Agricultural Utilization Research, 1815 N. University St., Peoria, IL 61604, United States

³ Departamento de Agronomia, Universidade Estadual de Maringá, Maringá, PR, Brazil

⁴ Biointeractions & Plant Health, Wageningen Plant Research, Wageningen, Netherlands

⁵ State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China

⁶ Universidad Nacional de Río Cuarto, Río Cuarto, Argentina

⁷ Laboratorio de Biología Funcional y Biotecnología (BIOLAB)-INBIOTEC-CONICET-CICBA, Facultad de Agronomía, Universidad Nacional del Centro, Buenos Aires, Argentina

⁸ Universidad de la República, Facultad de Ciencias-Facultad de Ingeniería, Montevideo, Uruguay

⁹ Department of Plant Pathology, University of Kentucky, Lexington, KY, United States

¹⁰ School of Plant and Environmental Sciences, Virginia Tech, Blacksburg, VA, United States

¹¹ Department of Plant Pathology and Environmental Microbiology, Pennsylvania State University, University Park, PA, United States

¹² National Research Council of Research, Institute of Sciences of Food Production, 70126, Bari, Italy

¹³ USDA, Agricultural Research Service, Cereal Disease Laboratory, 1551 Lindig St., St. Paul, MN, 55108, United States

¹⁴ School of Integrative Plant Science, Plant Pathology and Plant-Microbe Biology Section, Cornell University, Ithaca, NY 14853-5904, United States

¹⁵ Department of Plant Pathology, Stellenbosch University, Stellenbosch, South Africa

¹⁶ Plant Science, Western Cape Department of Agriculture, Elsenburg, South Africa

¹⁷ Microbial Safety Team, National Institute of Agricultural Sciences, Wanju, Republic of Korea

*Corresponding author: Emerson M. Del Ponte, delponte@ufv.br

♠ These authors contributed equally to the work

Abstract

Fusarium graminearum is ranked among the five most destructive fungal pathogens that affect agroecosystems. It causes floral diseases in small grain cereals including wheat, barley and oats, as well as summer crops such as maize and rice. We conducted a systematic review of peer-reviewed studies reporting species within the *Fusarium graminearum* species complex (FGSC) and created two main data tables. The first The first contained summarized data from the articles including bibliographic, geographic, methodological (ID methods), host of origin and species, while the second data table contains information about the described strains such as publication, isolate code(s), host/substrate, year of isolation, geographical coordinates, species and trichothecene genotype. Analyses of the bibliographic data obtained from 123 publications from 2000 to 2021 by 498 unique authors and published in 40 journals are summarized. We describe the frequency of species and chemotypes for 16,274 strains for which geographical information was available, either provided as raw data or extracted from

the publications, and sampled across six continents and 32 countries. The database and interactive interface are publicly available allowing for searches, summarization and mapping of strains according to several criteria including article, country, host, species and trichothecene genotype. The database will be updated continuously and should be useful for guiding future surveys and exploring factors associated with species distribution such as climate and land use. Authors are encouraged to submit data at the strain level to the database, which is accessible at <https://fgsc.netlify.app/>.

Introduction

Members of the *Fusarium graminearum* species complex (FGSC) include some of the top fungal plant pathogens of global importance given the serious diseases they cause in most major cereal crops, including wheat, maize, rice, and barley (Dean et al. 2012; van der Lee et al. 2015). Crop yields are often severely reduced due to infection of roots, stalks, or ears/heads (Duffeck et al. 2020; McMullen et al. 1997; Munkvold 2003; Vaughan et al. 2016). Kernel infections are of particular concern, since members of the FGSC can synthesize a wide range of mycotoxins during plant infection and colonization (D'Mello et al. 1999). The most important mycotoxins are the trichothecenes, including deoxynivalenol, nivalenol, and their acetylated derivatives, which can be produced at high levels in cereal grains, and are highly toxic to humans and other animals. Ingestion of grain or grain products contaminated by these toxins can cause diarrhea, nausea, vomiting, and inhibition of protein synthesis (Chen et al. 2019; Pestka 2010).

In wheat and barley, members of the FGSC cause Fusarium Head Blight (FHB), which re-emerged as a major concern throughout the world in the early 1990s (Goswami and Kistler 2004; McMullen et al. 1997, 2012). Numerous surveys of disease incidence and severity, pathogen population diversity, and mycotoxin contamination have been conducted during the past three decades in the main cereal-growing

regions of the world where FHB epidemics have occurred (Cowger et al. 2019; Pereira et al. 2021; van der Lee et al. 2015; Vogelgsang et al. 2019).

Multilocus sequencing of a global collection of strains that were morphologically defined as a single panmictic species, previously known as “*Fusarium graminearum*,” revealed significant phylogeographic structure and reproductive isolation of seven lineages (i.e., phylopecies; O’Donnell et al. 2000). Employing the concept of genealogical concordance phylogenetic species recognition (GCPSR) (Taylor et al. 2000), a later study based on portions of 11 genes identified nine distinct phylogenetic species within the morphospecies *F. graminearum sensu lato*, and the eight newly discovered species were formally described (O’Donnell et al. 2004). Subsequently, six additional species were discovered and described; however a singleton phylospecies from soil in Umyaka, South Africa, represented by NRRL 34461 *Fusarium* sp., remains unnamed. These 16 FGSC cannot be distinguished from one another using morphological data (O’Donnell et al. 2008; Sarver et al. 2011; Starkey et al. 2007; Yli-Mattila et al. 2009). These and other studies have also investigated the toxigenic profile (chemotype) of isolates using genetic markers and chemical analyses. Within the FGSC, differences in trichothecene chemotype are encoded by interacting balanced polymorphisms at both ends of the TRI biosynthetic gene cluster (Alexander et al. 2011; Kimura et al. 2003; Lee et al. 2002; Ward et al. 2002) and by differences in the unlinked *Tri1* gene (Varga et al. 2015). This genetic variation has been exploited for the development of PCR primers (Lee et al. 2001; Liang et al. 2014; Nielsen et al. 2012; Pasquali and Migheli 2014; Ward et al. 2002) or Luminex probes (Garmendia et al. 2018; Sarver et al. 2011; Ward et al. 2008) for genotyping to predict the type of trichothecene produced by individual isolates.

Recently, two reviews were published focusing on the systematics, occurrence and distribution of species and chemotypes of the FGSC (Aoki et al. 2012; van der Lee et al. 2015). Two quantitative reviews on the distribution of FGSC have also been published, but these are limited in the geographic coverage and species included. The

first summarizes occurrence and spatial distribution within the FGSC and *F. culmorum*, with an associated database of chemotype diversity, but is restricted to strains isolated from cereals in several countries in Europe (Pasquali et al. 2016). In the second review, the focus was on how climate might affect the occurrence and distribution of three members of the FGSC members, *F. graminearum sensu stricto* (hereafter *F. graminearum*), *F. asiaticum*, and *F. boothii* (Backhouse 2014).

In this study, we conducted a systematic analysis of the peer-reviewed literature to summarize and map the global distribution of FGSC species and associated trichothecene chemotypes. The specific objectives were to: (i) screen the literature since 2000 and organize a database of peer-reviewed studies within which members of the FGSC were identified; (ii) conduct a bibliographic and authorship network analysis; (iii) collect strain ID data and metadata at both the strain and study (summary of frequency) level; (iv) summarize trends in the identification methods and distribution of species and chemotypes by geography, host and year; and (v) build an online dashboard for searching and visualizing publication and strain data from the database. The database and interactive interface are publicly available (<https://fgsc.netlify.app/>) allowing for searches, summarization and mapping of strains according to several criteria including article, country, host, species, and trichothecene genotype. The database will be updated continuously with the expectation that it will be useful for guiding future surveys and exploring factors associated with species distribution such as climate and land use. Authors are encouraged to submit data at the strain level to the database.

Material and methods

Literature search and data gathering for the databases

Since 2017, we have been conducting bibliographic searches across three major databases (Google Scholar, Scopus, and Web of Science) using the following

keywords: "FGSC" and "*Fusarium graminearum* species complex." To facilitate the task, a "Google Alert" was employed using these keywords, so that automatic messages were sent each time a study containing either of these keywords was published.

Titles and abstracts of the search results were examined, and when these suggested that the strains had been identified to the species level by sequence-based molecular methods, the main text was further scrutinized. Several associated metadata were collected from each selected publication, and each was added to one or two separate databases (maintained as online google spreadsheets): 1) a spreadsheet named **FGSC publication database**, which contained all metadata of bibliographic information as well as the number and identity of FGSC strains and hosts; and 2) a spreadsheet named **FGSC isolate database**, which included year, geographic location (municipality, state/province, country), latitude, longitude, and host sampled, and the species and trichothecene chemotype (toxin production using chemical methods) or genotype (toxigenic potential using molecular assays targeting *TRI* genes) together with the respective method utilized. The overall workflow for gathering of data for the two databases is presented in Fig. 1.

Data description and summary

Metadata were described with summary statistics, tables, graphs, and maps to display absolute numbers and relative frequencies for several categorical and numerical variables (see metadata of the two databases in Fig. 1). A multiple correspondence analysis (MCA) was performed, given that the nature of multiple variables describing the isolates was nominal, to detect and represent underlying structure in the strain database (Hjellbrekke 2018).

Scholarly and country network analysis of article database

A scholarly collaborative network analysis was used to detect connections among authors and identify research communities (Kolaczyk and Csárdi 2020). Authors on

the same publication (co-authors) were grouped together and linked to other publications whenever authorship was shared. Each author is considered a "node" in the network and the connections between them are the "edges" (or links). Several statistics were used in this analysis including community detection (group of authors), network transitivity (the odds of two scholars sharing a common collaborator), node degree (number of connections for a single author), and betweenness (individuals who influence the flow around a system) (Kolaczyk and Csárdi 2020). The same analysis was conducted using countries as nodes. The Walktrap method, in which short random walks tend to stay in the same community, was used to detect communities in the network graphs, which were represented by different colors. The network analysis was performed in R using the *igraph* package (Csárdi and Nepusz 2006).

Results

Peer-reviewed article database

Bibliographic analysis

Scrutiny of the literature yielded 123 peer-reviewed publications that reported phylogenetic species within the FGSC (Supplementary Table 1). The earliest publications included the foundational phylogeographic study by O'Donnell et al. (2000) identifying seven distinct species within the FGSC, identification of an eighth phylogenetic species within this complex and demonstration that trichothecene chemotype diversity had been maintained by selection throughout the evolution of the FGSC and related species (Ward et al. 2002), and the first formal description of seven unnamed members of the FGSC and assignment of species names with *F. graminearum* retained for the most common species within Europe and North America, the continents where *F. graminearum* and *Gibberella zeae*, respectively, were formally described (O'Donnell et al., 2004). From 2005 onward, the average number of

publications per year was seven, ranging from 0 (in 2006) to 12 (in 2011 and 2020). Eighty percent of the articles were published between 2011 and 2021.

The studies were published in 40 different journals, with the large majority belonging to the subject areas of plant science/plant pathology, mycology, food science and technology, microbiology, and toxicology. Nearly half of the articles (48%) were published in one of these six journals: Plant Disease (n = 14), Plant Pathology (n = 11), Toxins (n = 9), Phytopathology (n = 9), Fungal Genetics and Biology (n = 9), and the International Journal of Food Microbiology (n = 8) (Fig. 2B).

Article title text mining

A text analysis of the words in the title of all publications (n = 123) identified 389 words, after excluding “stop words” (a set of commonly used words in a language [e.g. “a”, “the”, “is”] that carry very little information) which were ordered by frequency and displayed in a wordcloud map (Fig 3).

Authorship analysis

The selected group of studies included 498 unique authors representing 37 countries across six continents. The four countries that contributed the largest numbers of authors were the United States (n= 46 authors), China (n = 31), Brazil (n = 16), and Argentina (n = 14). Sixteen countries each contributed a single author (Fig. 4A). Most studies were conducted collaboratively by authors from the same country (n = 62 studies), followed by two (n = 46), three (n = 11), or four and six countries (n = 1 for each).

Among studies that included authors representing at least two countries, there were two relatively large communities (subgraph defined by colors) composed of authors from at least six different countries (Fig 4B). The largest nodes (highest number of connections) in these two communities were based in the United States,

China and Netherlands. Two small communities were formed by researchers in Italy as the largest node with authors from Luxembourg and Spain. One community was represented by a single publication by researchers in Latvia and Lithuania.

The largest subgraph (community with vectors in black) was led by researchers from the United States, who have had direct collaborations with scientists from institutions in 19 other countries from six continents. In the second largest community (blue vectors), China and Netherlands interacted with researchers from eight and seven other countries, respectively. The number of authors per publication ranged from three to nine, with a median of eight authors. Twelve researchers authored at least six articles, while four researchers authored at least ten articles (data not shown).

The authorship network positioned the most prolific (degree of 190) and influential (betweenness = 0.31) researcher in the central position (largest node) (Ward, TJ; Fig. 5). Six additional highly influential researchers had betweenness values of at least 0.035. The authorship network highlighted 14 authors (largest nodes) who collaborated with at least 38 individuals (degree), and who are centrally located and highly connected to other larger communities, identified by color. The probability that two authors with a common co-author were themselves co-authors (transitivity) was 0.473. The subgraphs depict communities that are mostly grouped by country, as expected. The largest of these included the United States community (Ward, O'Donnell, Kistler, Gale); Dutch (Waalwijk, van der Lee) and Chinese researchers (Zhang, Feng); South Korea (Lee, Choi); China (Chen); Brazil (Del Ponte); and Argentina (Chulze). Twenty-three of 123 single studies formed isolated, small clusters (see the outer edge of the network) representing authors who did not publish with authors from the other studies.

FGSC species and chemotypes by publication

A total of 26,644 isolates identified to species level were reported by the 123 peer-reviewed studies. The number of strains per study ranged from one (Dong et al. 2020) to 4,062 from a continent-wide survey (Kelly et al. 2015). The mean number of strains per study was 218.4, but half of the studies included at least 62 strains (median).

The dominant species was *F. graminearum*, reported in 106 studies. Five other species appeared in at least eight studies: *F. asiaticum* (47 studies), *F. meridionale* (36), *F. boothii* (20), *F. cortaderiae* (18) and *F. austroamericanum* (9). Ten additional species were each reported in at least one study.

Trichothecene chemotype diversity varied across the FGSC, with chemotype polymorphism observed in some species while others appear to be fixed for a specific trichothecene type. For example, *F. graminearum* and *F. asiaticum* were segregating for all three type B trichothecene chemotypes. The most prevalent was DON/15-ADON (n = 84 studies), followed by DON/3-ADON (n = 41) and NIV (n = 23). Two studies reported the presence of a DON/NIV chemotype in *F. graminearum*. In *F. asiaticum*, the NIV type was reported in 35/47 studies, followed by DON/3-ADON (n = 21 studies), and DON/15-ADON (n = 10 studies). Five of the six FGSC are segregating for NIV and 3-ADON. *F. meridionale* was the only species segregating for NIV and 15-ADON, but the 15-ADON type was found in just one isolate (Del Ponte et al. 2015). However, the NIV chemotype was reported in 28 of the 35 *F. meridionale* studies. Despite being reported in 20 studies, *F. boothii* appears to be fixed for the 15-ADON chemotype. More recently the type A trichothecene NX-3 and its 3-acetyl derivative, NX-2 were reported (n = 4 studies) in *F. graminearum* from North America.

Cereal hosts by publication

A total of 45 hosts were included in the 123 studies. The top six hosts sampled were wheat (n = 71 studies), maize (n = 33), barley (n = 29), rice (n = 12), soybean (n = 8), and

oats (n = 6). Other hosts were included in fewer than three publications each, with thirty represented by a single report.

Summary of the FGSC strain database

Strains by country

Strains associated with geographical information (n = 16,274), either provided as raw data or extracted from the publications, were sampled across six continents and 32 countries (Fig. 6). The majority (> 10,000 strains) were obtained from cereal crops grown in the Americas (Canada, Brazil, and the United States) and in China. In Europe, the most sampled countries (> 100 strains) were France, Netherlands, and Norway. In Asia, aside from China, the most sampled countries (> 100 strains) were Japan and South Korea. The only study from Africa that included >100 strains was from South Africa (560 strains) (Boutigny et al. 2011).

Methodology for species and chemotype identification

Morphological characters have proven to be inadequate for the identification of individual species within the FGSC (Aoki et al. 2012). As such, species identifications are performed using a variety of DNA-based methods. Analyses of DNA sequences from individual genes represents the gold-standard for identifying individual isolates to species level within the FGSC. Unfortunately, many of the sequences available in the GenBank database are associated with inaccurate species information and may be misleading (O'Donnell et al. 2015). To address this, the FUSARIUM-ID (<http://isolate.fusariumdb.org/blast.php>) and *Fusarium* MLST (<https://fusarium.mycobank.org/>) databases of curated DNA sequence data, primarily from translation elongation factor 1- α (*TEF1*) and the DNA-directed RNA polymerase II largest (*RPB1*) and second largest subunit (*RPB2*), were developed to facilitate

identification of the FGSC and other fusaria via BLASTn queries. A primer for conducting and interpreting the results of such analyses was provided by O'Donnell et al. (2015). However, targeted sequencing of individual genes and associated analyses can be relatively laborious and has been used to identify 11% (n = 1,807) of the FGSC isolates included in the studies analyzed here.

Some studies employed multiple methods to identify species within the FGSC, but most of the isolates were identified using a multilocus genotyping assay (MLGT) employing a Luminex flow cytometer (n = 12,317 strains; 75.6%). This extensively validated and high-throughput method relies on probes targeting nucleotide variation within multiple genes to simultaneously determine the species identity and trichothecene chemotype of isolates (Ward et al. 2008). The MLGT method has also enabled the discovery of novel FGSC and *F. graminearum* Clade species (sensu Laraba et al. 2021) and interspecific hybrids (Boutigny et al. 2011, 2014; Pereira et al. 2018; Sarver et al. 2011). A smaller number of studies have employed fragment analysis of polymorphic markers (RFLP, AFLP or ISSR; n = 1,686; 10.3%), and polymerase chain reaction (PCR) with species-specific primers (e.g. Fg16; n = 1,427; 8.7%).

Genotype analyses was the most common method for assessing the trichothecene type of individual strains. The MLGT method was most frequently used (n = 11,032 strains; 67.8%), followed by chemotype-specific PCR primers targeting at least one of the TRI genes (n = 4,103 strains; 10.3%). The recent discovery of a relatively small number of *F. graminearum* isolates producing NX-2 and related Type A trichothecenes represents variation that was not accounted for in the original design of many genotype-based methods. A PCR-RFLP approach targeting *TRI1* was developed to distinguish NX-2 from the 3-ADON chemotype (Liang et al. 2014) and the MLGT assay was subsequently upgraded to differentiate these chemotypes as well (Garmendia et al. 2018).

Chemical analyses can be used to document the actual production of trichothecene metabolites or the relative amounts of different trichothecenes that may

be produced by an isolate under specific conditions. However, their application has been limited by cost and availability. Analytical chemistry was used to determine trichothecene toxin production in 1,286 strains (~8% of the strains). The most common method was chromatography (e.g. HPLC, GC-MS, LC-MS), which was used to chemotype 1,099 strains (85%).

Isolates by host

The most sampled hosts, with at least 801 isolates recovered from each, were four commercially grown cereal crops: wheat (n = 12,233 strains, 75.1%), maize (n = 1,706, 10.5%), barley (n = 980, 6%), and rice (n = 801 strains, 4.9%). Fewer than 90 isolates were obtained from oats (n = 89, 0.54%), and soybean (n = 77, 0.47%). Less than 26 isolates (0.15%) were recovered from the other hosts (cereal and non-cereals). The host of origin was not reported for 201 isolates.

Strains by FGSC species

Within the whole collection of characterized strains (n = 16,274), *F. graminearum* was dominant, with 11,760 strains (72.2%), followed by *F. asiaticum* (n = 3,052, 18.7%), *F. meridionale* (n = 816; 5%), *F. boothii* (n = 279; 1.7%), *F. cortaderiae* (n = 201; 1.3%) and *F. austroamericanum* (n = 50, 0.3%). The following species were represented by 10 to 50 strains each: *F. nepalense* (n = 41), *F. aethiopicum* (n = 22), *F. vorosii* (n = 16), *F. ussurianum* (n = 12), and *F. acaciae-mearnsii* (n = 10). Finally, species with less than 10 strains were *F. brasiliicum* (n = 5 strains), *F. boothii* x *F. graminearum* hybrid (n = 3), *F. gerlachii* (n = 3), *F. louisianense* (n = 2), and *F. mesoamericanum* (n = 2).

349

350 **Strains by chemotypes**

351 Among the three most frequent chemotypes, DON/15-ADON was dominant (n = 9,067
 352 strains) followed by DON/3-ADON (n = 3,475) and NIV (n = 2,437). Other less
 353 frequently chemotypes, as reported, were: DON (n = 39), NX2 (n = 36), DON/NIV (n =
 354 18), NX2/3ADON (n = 5), 4,15-diANIV (n = 2).

355

356 **Strains by host × species × chemotypes**

357 A MCA was conducted to depict associations among the four most sampled hosts, six
 358 most represented species, and three most frequent chemotypes, totaling 14,393 strains.
 359 The MCA map confirmed the close associations between these levels and factors. The
 360 first and second dimensions retained a modest proportion of 37.1% (16.7% + 20.4%) of
 361 the total inertia (variation) contained in the data. This was similar to the result when
 362 each factor was analyzed separately, but the MCA more clearly connected the host
 363 with the species, and the species with the chemotypes. For example, *F. graminearum*, a
 364 species composed primarily of the DON/15-ADON chemotype, clustered more with
 365 wheat and barley than with the other crops. *Fusarium asiaticum*, which was more
 366 commonly NIV or 3-ADON, was more closely related to rice. *Fusarium meridionale*
 367 (NIV chemotype) and *F. boothii* (15-ADON chemotype) were more closely associated
 368 with maize. *Fusarium austroamericanum*, found mainly in wheat and barley, was
 369 represented primarily by the 3-ADON chemotype.

370

371 **Global mapping of species and trichothecene toxin chemotypes**

372 The global species composition appeared to be structured by region and host of origin.
 373 For example, *F. asiaticum* was most prevalent in Asia but has also been reported on
 374 rice and wheat in South America and wheat in a major rice-growing region of North

America (Fig. 8). *Fusarium meridionale* was most prevalent in the lower latitudes of South America (associated with maize) and also in central-south region of China (Fig. 7A). *Fusarium graminearum* was the most frequent species globally and was recovered across a wide range of latitudes. *Fusarium boothii* was found primarily in South Africa, México, Argentina and north Asia, with scattered incidence across other continents.

Discussion

We conducted a systematic analysis of peer-reviewed studies reporting on species within the FGSC. After a comprehensive and quantitative analysis of the literature, we have summarized the past 20 years of research on FGSC strains isolated from cereal and non-cereal hosts worldwide. Our study updates and complements a recent review of the global composition and spatial distribution of FGSC (van der Lee et al. 2015). Collectively, the studies included here provide a broad geographic assessment of FGSC species and chemotype diversity from different crops/substrates. The extent to which FGSC diversity was incorporated into the design and validation of different typing methods should be considered when interpreting the results of individual studies.

FGSC research during the past two decades can be divided into two phases. The first phase was characterized by the discovery that the morphospecies *F. graminearum* comprises at least 16 phylogenetic species, and development of tools to facilitate identification of these species and of their trichothecene genotypes. It should be noted that four species basal to the FGSC, in what is now recognized as the Graminearum Clade (Laraba et al. 2021), were discovered and described during roughly the same period. These included *F. pseudograminearum*, which was previously reported as the Group 1 population of *F. graminearum* (Aoki and O'Donnell 1999), *F. dactylidis* from orchard grass in Oregon and New Zealand (Aoki et al. 2015), *F. praegraminearum* from maize field litter in New Zealand (Gräfenhan et al. 2016), and *F. subtropicale* from barley in Brazil (Pereira et al. 2018). During the second phase, these

advancements were widely adopted, including use of binomials for the phylospecies and genotyping tools for large-scale surveys, leading to a rapid rise in the number of studies published each year. Some of this increase was related to wider availability of cost-effective methods for species identification and chemotyping. More intensive within- and cross-continental scientific cooperation, led by research groups from North America and Europe that appear as major nodes in our network analysis, also played an important role. Strengthened global collaborations have facilitated the extension and expansion of surveys to crops other than wheat and barley, including maize and rice. There has also been an expansion in regional surveys in previously underrepresented areas, and these efforts have dramatically increased the number of peer-review studies assessing FGSC species and chemotype diversity over the past ten years.

Overall, the research included in the present study was published in over 40 journals with a broad range of impact factors and target audiences. This was likely driven by an interest in delivering information at more regional or local levels, and by close connections with a range of fields including mycotoxicology, plant pathology, mycology and food microbiology. Expanded collaboration among groups worldwide, particularly during the last decade, has increased the number of studies published in international journals with a more global audience, mainly in the field of plant pathology. A third of the papers appeared in four journals published by international plant pathology professional societies.

The present review highlights two interesting points regarding FGSC species diversity. First, no novel member of the FGSC has been discovered since the study by Sarver et al. (2011), despite the several thousands of strains that were molecularly identified. Second, 11 of the 16 species comprise only 1.3% of all strains in the database. The available data suggests that the 11 less commonly collected species may be restricted in their geographic distribution and/or host.

It is apparent that host is an important driver of the regional composition of FGSC species. Sampling to date has been conducted primarily in commercial cereal fields where a few species are globally dominant. The most common FGSC species in the surveys, *F. graminearum*, was distributed worldwide on wheat and other cereals. However, *F. asiaticum* appears to be better adapted to rice agroecosystems and is the most common member of the FGSC on rice in Asia (Lee et al. 2010), but also in other crops in some parts of Asia including barley (Yang et al. 2008) and wheat (Zhang et al. 2012), depending on the crop rotation. More recently, *F. asiaticum* was found to be the dominant species on rice in South America (Gomes et al. 2015) and has also been observed on wheat in rice production regions of the United States (Gale et al. 2011) and Uruguay (Umpiérrez-Failache et al. 2013). Similarly, *F. meridionale* was more prevalent than *F. graminearum* on maize and rice in South America, and *F. boothii* was dominant on South African maize (Boutigny et al. 2011). The reasons for the observed variation in species dominance are not entirely clear, but recent studies have suggested that differences in aggressiveness among the species may play a role (Machado et al. 2021).

Higher FGSC species diversity was found in wheat crops grown at lower latitudes, near to or within subtropical environments (Pereira et al. 2021; Umpiérrez-Failache et al. 2013). This suggests that there is also an effect of climate in shaping species distributions, as posited previously by a modelling study (Backhouse 2014). For example, although members of the FGSC can be found in the tropics, a drier environment during the season limits disease occurrence, which in turn decreases inoculum availability. However, this can be easily confounded with the effect of different host species, which is also dependent on climate. It appears that a small group of minor species including *F. meridionale*, *F. boothii*, and *F. cortaderiae* are more frequent in subtropical conditions where there is an abundance of other grasses year-round. A study in Brazil that surveyed a relatively small collection of FGSC in annual ryegrass showed an increased frequency of these minor species in comparison with

their composition on major crops like wheat (Machado et al. 2015). A recent study in the United States reported FGSC as endophytes in several wild grasses, but the phylogenetic species were not identified (Lofgren et al. 2018).

Among the three main chemotypes from global surveys, 15-ADON represented more than half of the strains, while 3-ADON and NIV were less frequent and restricted in distribution. The 3-ADON chemotype prevailed at higher latitudes, while NIV was more frequent at lower ones, as suggested by our global map. Whether these species have different requirements of temperature is not entirely clear, as it is difficult to disentangle a triple interaction effect that includes climate, host, and fungal species that shape chemotype composition. Additional global surveys are needed to address whether some of the rarely collected species are fixed for a specific chemotype. Relative chemotype frequencies in *F. graminearum* and *F. asiaticum*, which are segregating for all three trichothecene toxin chemotypes globally, appear to be shaped by region-specific factors. However, changes in chemotype frequency can change rapidly as evidenced by the introduction and spread of a non-native 3-ADON population in North America (Ward et al. 2008).

An important outcome of the present research is a website (<http://fgsc.netlify.app>) that facilitates searching, visualization and downloading of data and reports in both tabular and graphic format. The system is composed of three main views (grid, map and chart) all of which allow for interactive navigation, searching, and selection of data for display. For example, data filtered or searched for in the grid view can be downloaded in csv or xlsx format, and screenshots can be taken from the maps and plots.

All website content (data and summaries) can be downloaded and used freely according to a Creative Commons Attribution 4.0 International (CC BY 4.0). The database will be updated continuously, and we encourage all members of the research community to deposit raw data from their prior and future work on the topic

following instructions on the website, which is an online spreadsheet of the web-based Google Docs Editor Suite.

Acknowledgements

Emerson M. Del Ponte was supported by the National Council for Scientific and Technological Development (CNPq) through a Productivity Research Fellowship (PQ) project 310208/2019-0. Gláucia M. Moreira was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) through a PNPd scholarship. This work was supported in part by the U.S. Department of Agriculture, Agricultural Research Service. This work is also partly supported by from the National Key R&D Program of China (2017YFE0126700) and the European Union's Horizon 2020 research and innovation programme under grant agreement No 678781 (MycoKey). Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the United States Department of Agriculture (USDA). USDA is an equal opportunity provider and employer.

References

- Alexander, N. J., McCormick, S. P., Waalwijk, C., van der Lee, T., and Proctor, R. H. 2011. The genetic basis for 3-ADON and 15-ADON trichothecene chemotypes in *Fusarium*. *Fungal Genet. Biol.* 48:485–495.
- Aoki, T., and O'Donnell, K. 1999. Morphological and molecular characterization of *Fusarium pseudograminearum* sp. nov., formerly recognized as the Group 1 population of *F. graminearum*. *Mycologia* 91:597–609.
- Aoki, T., Vaughan, M. M., McCormick S. P., Busman, M., Ward, T. J., Kelly, A., and O'Donnell, K. 2015. *Fusarium dactylidis* sp. nov., a novel nivalenol toxin-producing species sister to *F. pseudograminearum* isolated from orchard grass (*Dactylis glomerata*) in Oregon and New Zealand. *Mycologia* 107:409–418.

- 512 Aoki, T., Ward, T. J., Kistler, H. C., and O'Donnell, K. 2012. Systematics, phylogeny
513 and trichothecene mycotoxin potential of *Fusarium* head blight cereal
514 pathogens. JSM Mycotoxins 62:91–102.
- 515 Backhouse, D. 2014. Global distribution of *Fusarium graminearum*, *F. asiaticum* and *F.*
516 *boothii* from wheat in relation to climate. Eur. J. Plant Pathol. 139:161–173.
- 517 Boutigny, A.-L., Ward, T. J., Ballois, N., Iancu, G., and Ioos, R. 2014. Diversity of the
518 *Fusarium graminearum* species complex on French cereals. Eur. J. Plant Pathol.
519 138:133–148.
- 520 Boutigny, A.-L., Ward, T. J., Van Coller, G. J., Flett, B., Lamprecht, S. C., O'Donnell,
521 K., and Viljoen, A. 2011. Analysis of the *Fusarium graminearum* species
522 complex from wheat, barley and maize in South Africa provides evidence of
523 species-specific differences in host preference. Fungal Genet. Biol. 48:914–920.
- 524 Chen, Y., Kistler, H. C., and Ma, Z. 2019. *Fusarium graminearum* trichothecene
525 mycotoxins: Biosynthesis, regulation, and management. Annu. Rev.
526 Phytopathol. 57:15–39.
- 527 Cowger, C., Smith, J., Boos, D., Bradley, C. A., Ransom, J., and Bergstrom, G. C. 2019.
528 Managing a destructive, episodic crop disease: A national survey of wheat
529 and barley growers' experience with *Fusarium* head blight. Plant Dis. 104:
530 634–648.
- 531 Csárdi, G., and Nepusz, T. 2006. The igraph software package for complex network
532 research. InterJournal, complex systems 1695:1–9.
- 533 Dean, R., Kan, J. A. L. V., Pretorius, Z. A., Hammond-Kosack, K. E., Pietro, A. D.,
534 Spanu, P. D., Rudd, J. J., Dickman, M., Kahmann, R., Ellis, J., and Foster, G. D.
535 2012. The top 10 fungal pathogens in molecular plant pathology. Mol. Plant
536 Pathol. 13:414–430.
- 537 D'Mello, J. P. F., Placinta, C. M., and Macdonald, A. M. C. 1999. *Fusarium*
538 mycotoxins: A review of global implications for animal health, welfare and
539 productivity. Anim. Feed Sci. Tech. 80:183–205.

- 540 Dong, F., Xu, J. H., Shi, J. R., Mokoena, M. P., Olaniran, A. O., Chen, X. Y., and Lee,
541 Y. W. 2020. First report of *Fusarium* head blight caused by *Fusarium*
542 *meridionale* in rice in China. *Plant Dis.* 104:2726.
- 543 Duffeck, M. R., Alves, K. S., Machado, F. J., Esker, P. D., and Del Ponte, E. M. 2020.
544 Modeling yield losses and fungicide profitability for managing *Fusarium*
545 head blight in Brazilian spring wheat. *Phytopathology* 110:370–378.
- 546 Gale, L. R., Harrison, S. A., Ward, T. J., O'Donnell, K., Milus, E. A., Gale, S. W., and
547 Kistler, H. C. 2011. Nivalenol-type populations of *Fusarium graminearum* and
548 *F. asiaticum* are prevalent on wheat in southern Louisiana. *Phytopathology*
549 101:124–134.
- 550 Garmendia, G., Pattarino, L., Negrin, C., Martínez-Silveria, A., Pereyra, S., Ward, T.
551 J., and Vero, S. 2018. Species composition, toxigenic potential and
552 aggressiveness of *Fusarium* isolates causing Head Blight of barley in Uruguay.
553 *Food Microbiol.* 76:426–433.
- 554 Gomes, L. B., Ward, T. J., Badiale-Furlong, E., and Del Ponte, E. M. 2015. Species
555 composition, toxigenic potential and pathogenicity of *Fusarium graminearum*
556 species complex isolates from southern Brazilian rice. *Plant Pathol.* 64:980–
557 987.
- 558 Goswami, R. S., and Kistler, H. C. 2004. Heading for disaster: *Fusarium graminearum*
559 on cereal crops. *Mol. Plant Pathol.* 5:515–525.
- 560 Gräfenhan, T., Johnston, P. R., Vaughan, M. M., McCormick, S. P., Proctor, R. H.,
561 Busman, M., Ward, T. J., and O'Donnell, K. 2016. *Fusarium praegraminearum*
562 sp. nov., a novel nivalenol mycotoxin-producing pathogen from New Zealand
563 can induce head blight on wheat. *Mycologia* 108:1229–1239.
- 564 Hjellbrekke, J. 2018. Multiple Correspondence Analysis for the Social Sciences.
565 Routledge.
- 566 Kelly, A. C., Clear, R. M., O'Donnell, K., McCormick, S., Turkington, T. K., Tekauz,
567 A., Gilbert, J., Kistler, H. C., Busman, M., and Ward, T. J. 2015. Diversity of

- 568 Fusarium head blight populations and trichothecene toxin types reveals
 569 regional differences in pathogen composition and temporal dynamics. Fungal
 570 Genet. Biol. 82:22–31.
- 571 Kimura, M., Tokai, T., O'Donnell, K., Ward, T. J., Fujimura, M., Hamamoto, H.,
 572 Shibata, T., and Yamaguchi, I. 2003. The trichothecene biosynthesis gene
 573 cluster of *Fusarium graminearum* F15 contains a limited number of essential
 574 pathway genes and expressed non-essential genes. FEBS Lett. 539:105–110.
- 575 Kolaczyk, E. D., and Csárdi, G. 2020. Statistical analysis of network data with R.
 576 Springer-Verlag, New York.
- 577 Laraba, I, McCormick, S. P., Vaughan, M. M., Geiser, D. M., and O'Donnell, K. 2021.
 578 Phylogenetic diversity, trichothecene potential, and pathogenicity within
 579 *Fusarium sambucinum* species complex. PLoS ONE 16:e0245037.
- 580 Lee, T., Han, Y. K., Kim, K. H., Yun, S. H., and Lee, Y. W. 2002. *Tri13* and *Tri7*
 581 determine deoxynivalenol-and nivalenol-producing chemotypes of *Gibberella*
 582 *zeae*. Appl. Environ. Microbiol. 68:2148–2154.
- 583 Lee, S.-H., Lee, J.-K., Nam, Y.-J., Lee, S.-H., Ryu, J.-G., and Lee, T. 2010. Population
 584 structure of *Fusarium graminearum* from maize and rice in 2009 in Korea. The
 585 Plant Pathol. J. 26:321–327.
- 586 Lee, T., Oh, D.-W., Kim, H.-S., Lee, J., Kim, Y.-H., Yun, S.-H., and Lee, Y.-W. 2001.
 587 Identification of deoxynivalenol- and nivalenol-producing chemotypes of
 588 *Gibberella zeae* by using PCR. Appl. Environ. Microbiol. 67:2966–2972.
- 589 Liang, J. M., Xayamongkhon, H., Broz, K., Dong, Y., McCormick, S. P., Abramova, S.,
 590 Ward, T. J., Ma, Z. H., and Kistler, H. C. 2014. Temporal dynamics and
 591 population genetic structure of *Fusarium graminearum* in the upper
 592 Midwestern United States. Fungal Genet. Biol. 73:83–92.
- 593 Lofgren, L. A., LeBlanc, N. R., Certano, A. K., Nachtigall, J., LaBine, K. M., Riddle, J.,
 594 Broz, K., Dong, Y., Bethan, B., Kafer, C. W., Kistler, H. C. 2018. *Fusarium*
 595 *graminearum*: Pathogen or endophyte of North American grasses? New

- Phytologist 217: 1203–1212 doi.org/10.1111/nph.14894.
- Machado, F. J., Kuhnem, P., Casa, R. T., McMaster, N., Schmale, D., Vaillancourt, L. J., and Del Ponte, E. 2021. The dominance of *Fusarium meridionale* over *F. graminearum* causing Gibberella ear rot in Brazil may be due to increased aggressiveness and competitiveness. *Phytopathology*, in press.
- Machado, F. J., Möller, P. A., Nicolli, C. P., Del Ponte, E. M., and Ward, T. J. 2015. First report of *Fusarium graminearum*, *F. asiaticum*, and *F. cortaderiae* as head blight pathogens of annual ryegrass in Brazil. *Plant Dis.* 99:1859.
- McMullen, M., Bergstrom, G., De Wolf, E., Dill-Macky, R., Hershman, D., Shaner, G., and Van Sanford, D. 2012. A unified effort to fight an enemy of wheat and barley: *Fusarium* Head Blight. *Plant Dis.* 96:1712–1728.
- McMullen, M., Jones, R., and Gallenberg, D. 1997. Scab of wheat and barley: A re-emerging disease of devastating impact. *Plant Dis.* 81:1340–1348.
- Munkvold, G. P. 2003. Epidemiology of *Fusarium* diseases and their mycotoxins in maize ears. In: Xu, X., Bailey, J. A., and Cooke, B. M. (Eds.), *Epidemiology of Mycotoxin Producing Fungi*, Springer, Netherlands, pp. 705–713.
- Nielsen, L. K., Jensen, J. D., Rodríguez, A., Jørgensen, L. N., and Justesen, A. F. 2012. *TRI12* based quantitative real-time PCR assays reveal the distribution of trichothecene genotypes of *F. graminearum* and *F. culmorum* isolates in Danish small grain cereals. *Int. J. Food Microbiol.* 157:384–392.
- O'Donnell, K., Kistler, H. C., Tacke, B. K., and Casper, H. H. 2000. Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proc. Natl. Acad. Sci. U.S.A.* 97:7905–7910.
- O'Donnell, K., Ward, T. J., Aberra, D., Kistler, H. C., Aoki, T., Orwig, N., Kimura, M., Bjørnstad, Å., and Klemsdal, S. S. 2008. Multilocus genotyping and molecular phylogenetics resolve a novel head blight pathogen within the *Fusarium graminearum* species complex from Ethiopia. *Fungal Genet. Biol.* 45:1514–

- 624 1522.
- 625 O'Donnell, K., Ward, T. J., Geiser, D. M., Kistler, H. C., and Aoki, T. 2004.
- 626 Genealogical concordance between the mating type locus and seven other
- 627 nuclear genes supports formal recognition of nine phylogenetically distinct
- 628 species within the *Fusarium graminearum* clade. *Fungal Genet. Biol.* 41:600–
- 629 623.
- 630 O'Donnell, K., Ward, T. J., Robert, V. A. R. G., Crous, P. W., Geiser, D. M., and Kang,
- 631 S. 2015. DNA sequence-based identification of *Fusarium*: current status and
- 632 future directions. *Phytoparasitica* 43:583–595.
- 633 Pasquali, M., Beyer, M., Logrieco, A., Audenaert, K., Balmas, V., Basler, R., Boutigny,
- 634 A.-L., Chrpová, J., Czembor, E., Gagkaeva, T., González-Jaén, M. T.,
- 635 Hofgaard, I. S., Köycü, N. D., Hoffmann, L., Lević, J., Marin, P., Miedaner, T.,
- 636 Migheli, Q., Moretti, A., Müller, M. E. H., Munaut, F., Parikka, P., Pallez-
- 637 Barthel, M., Piec, J., Scauflaire, J., Scherm, B., Stanković, S., Thrane, U., Uhlig,
- 638 S., Vanheule, A., Yli-Mattila, T., and Vogelgsang, S. 2016. A European
- 639 database of *Fusarium graminearum* and *F. culmorum* trichothecene genotypes.
- 640 *Front. Microbiol.* 7:406.
- 641 Pasquali, M., and Migheli, Q. 2014. Genetic approaches to chemotype determination
- 642 in type B-trichothecene producing Fusaria. *Int. J. Food Microbiol.* 189:164–182.
- 643 Pereira, C. B., Ward, T. J., Ponte, E. M. D., Moreira, G. M., Busman, M., McCormick,
- 644 S. P., Feksa, H. R., Almeida, J. L. D., and Tessmann, D. J. 2021. Five-year
- 645 survey uncovers extensive diversity and temporal fluctuations among
- 646 fusarium head blight pathogens of wheat and barley in Brazil. *Plant Pathol.*
- 647 70:426–435.
- 648 Pereira, C. B., Ward, T. J., Tessmann, D. J., Del Ponte, E. M., Laraba, I., Vaughan, M.
- 649 M., McCormick, S. P., Busman M., Kelly, A., Proctor, R. H., and O'Donnell, K.
- 650 2018. *Fusarium subtropicale* sp. nov., a novel nivalenol mycotoxin-producing
- 651 species isolated from barley (*Hordeum vulgare*) in Brazil and sister to *F.*

- 652 *praegraminearum*. Mycologia 110:860–871.
- 653 Pestka, J. J. 2010. Deoxynivalenol: Mechanisms of action, human exposure, and
654 toxicological relevance. Arch. Toxicol. 84:663–679.
- 655 Sarver, B. A. J., Ward, T. J., Gale, L. R., Broz, K., Kistler, H. C., Aoki, T., Nicholson, P.,
656 Carter, J., and O'Donnell, K. 2011. Novel Fusarium head blight pathogens
657 from Nepal and Louisiana revealed by multilocus genealogical concordance.
658 Fungal Genet. Biol. 48:1096–1107.
- 659 Starkey, D. E., Ward, T. J., Aoki, T., Gale, L. R., Kistler, H. C., Geiser, D. M., Suga, H.,
660 Tóth, B., Varga, J., and O'Donnell, K. 2007. Global molecular surveillance
661 reveals novel Fusarium head blight species and trichothecene toxin diversity.
662 Fungal Genet. Biol. 44:1191–1204.
- 663 Taylor, J. W., Jacobson, D. J., Kroken, S., Kasuga, T., Geiser, D. M., Hibbett, D. S., and
664 Fisher, M. C. 2000. Phylogenetic species recognition and species concepts in
665 fungi. Fungal Genet. Biol. 31:21–32.
- 666 Umpiérrez-Failache, M., Garmendia, G., Pereyra, S., Rodríguez-Haralambides, A.,
667 Ward, T. J., and Vero, S. 2013. Regional differences in species composition and
668 toxigenic potential among Fusarium head blight isolates from Uruguay
669 indicate a risk of nivalenol contamination in new wheat production areas. Int.
670 J. Food Microbiol. 166:135–140.
- 671 van der Lee, T., Zhang, H., van Diepeningen, A., and Waalwijk, C. 2015.
672 Biogeography of *Fusarium graminearum* species complex and chemotypes: A
673 review. Food Addit. Contam. Part A Chem. Anal. Control Expo. Risk Assess.
674 32:453–460.
- 675 Varga, E., Wiesenberger, G., Hametner, C., Ward, T. J., Dong, Y., Schöffbeck, D.,
676 McCormick, S., Broz, K., Stückler, R., Schuhmacher, R., Krska, R., Kistler, H.
677 C., Berthiller, F., and Adam, G. 2015. New tricks of an old enemy: isolates of
678 *Fusarium graminearum* produce a type A trichothecene mycotoxin. Environ.
679 Microbiol. 17:2588–2600.

- 680 Vaughan, M., Backhouse, D., and Ponte, E. M. D. 2016. Climate change impacts on
 681 the ecology of *Fusarium graminearum* species complex and susceptibility of
 682 wheat to *Fusarium* head blight: A review. *World Mycotoxin J.* 9:685–700.
- 683 Vogelgsang, S., Beyer, M., Pasquali, M., Jenny, E., Musa, T., Bucheli, T. D., Wettstein,
 684 F. E., and Forrer, H.-R. 2019. An eight-year survey of wheat shows distinctive
 685 effects of cropping factors on different *Fusarium* species and associated
 686 mycotoxins. *Eur. J. Agron.* 105:62–77.
- 687 Ward, T. J., Bielawski, J. P., Kistler, H. C., Sullivan, E., and O'Donnell, K. 2002.
 688 Ancestral polymorphism and adaptive evolution in the trichothecene
 689 mycotoxin gene cluster of phytopathogenic *Fusarium*. *Proc. Natl. Acad. Sci.*
 690 U.S.A. 99:9278–9283.
- 691 Ward, T. J., Clear, R. M., Rooney, A. P., O'Donnell, K., Gaba, D., Patrick, S., Starkey,
 692 D. E., Gilbert, J., Geiser, D. M., and Nowicki, T. W. 2008. An adaptive
 693 evolutionary shift in *Fusarium* head blight pathogen populations is driving
 694 the rapid spread of more toxigenic *Fusarium graminearum* in North America.
 695 *Fungal Genet. Biol.* 45:473–484.
- 696 Yang, L., van der Lee, T., Yang, X., Yu, D., and Waalwijk, C. 2008. *Fusarium*
 697 populations on Chinese barley show a dramatic gradient in mycotoxin
 698 profiles. *Phytopathology* 98:719-727.
- 699 Yli-Mattila, T., Gagkaeva, T., Ward, T. J., Aoki, T., Kistler, H. C., and O'Donnell, K.
 700 2009. A novel Asian clade within the *Fusarium graminearum* species complex
 701 includes a newly discovered cereal head blight pathogen from the Russian Far
 702 East. *Mycologia* 101:841–852.
- 703 Zhang, H., van der Lee, T., Waalwijk, C., Chen, W., Xu, J., Xu, J., Zhang, Y., and
 704 Feng, J. 2012. Population analysis of the *Fusarium graminearum* species
 705 complex from wheat in China show a shift to more aggressive isolates. *PLoS*
 706 *ONE* 7:e31722.

710

714

715

Figure 2. A. Number of peer-reviewed publications per year reporting species and chemotypes identified using molecular methods in the *Fusarium graminearum* species complex (FGSC) isolated from several hosts. The fill colors represent whether geographic location of the strain was obtained from the main text or supplementary files of the publication or directly from the authors, as well as cases in which this information was not available. B. Twelve scientific journals with the largest number (three or more) of peer-reviewed studies on FGSC published between 2000 and 2021.

Figure 4. A. Global map depicting the number of authors per country of origin. B. Network graph depicting countries as nodes and subgraphs (represented by different colors of the nodes) as communities of countries. These were defined using a Walktrap method (in which short random walks tend to stay in the same community). The data represents 121 peer-reviewed studies reporting species and chemotypes of the *Fusarium graminearum* species complex sampled across several hosts, and published between 2000 and 2021.

Figure 5. Network graph depicting authors as nodes and subgraphs (represented by different colors of the nodes) as communities of authors. These were defined using a Walktrap method in which short random walks tend to stay in the same community. The communities are highly connected due to shared authorship of 123 studies reporting species and chemotypes of *Fusarium graminearum* species complex (FGSC) isolates sampled across several hosts published between 2000 and 2021. Larger nodes are labelled with the names of authors with more than 37 connections (co-authors).

A

Figure 6. A. Global distribution of 16,274 *Fusarium graminearum* complex strains identified to species using molecular methods. These were reported in 123 peer-reviewed studies published in 30 journals from 2000 to 2021. B. The inset map represents the number of strains in the top 10 most sampled countries.

Figure 7. A. Multiple correspondence analysis plot depicting association among variables representing cereal hosts, *Fusarium graminearum* species complex species and trichothecene chemotypes (sample size = 14,393 strains). The number of strains per host and species is shown in B and C, respectively.

Figure 8. Global distribution of the six most prevalent species of the *Fusarium graminearum* species complex. The geolocations of 16,158 out of 16,274 strains of these six species were obtained from authors or extracted from studies published in 121 scientific journals between 2000 and 2021.

Figure 9. Global distribution of the three most prevalent trichothecene chemotypes (15-ADON = 15-acetyl deoxynivalenol; 3-ADON = 3-acetyl deoxynivalenol; and NIV = nivalenol) identified for 14,979 strains out of 15,274 with geolocation data. The chemotype data was obtained from authors or extracted from studies) of *Fusarium graminearum* complex species isolated from a range of hosts reported in studies published between 2000 and 2021.